

DANGEROUS POISON

KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

KENSO AGCARE

BROMOXYNIL M

SELECTIVE HERBICIDE

ACTIVE CONSTITUENT: 200 g/L BROMOXYNIL present as the OCTANOATE
200 g/L MCPA present as the ETHYL HEXYL ESTER
SOLVENT: 324 g/L LIQUID HYDROCARBON

GROUP	6	4	HERBICIDE
-------	---	---	-----------

For control of broadleaf weeds in cereals, linseed and grass pastures as specified in the
DIRECTIONS FOR USE Table.

Kenso Corporation (M) Sdn Bhd
Level 1, 98 Commercial Road,
Teneriffe QLD 4005
Phone (07) 3216 1188
www.kenso.com.au



IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USE

CONTENTS: 20 Litres APVMA Approval No.: 66163/ RV2024

KENSO AGCARE BROMOXYNIL M SELECTIVE HERBICIDE

STORAGE AND DISPOSAL

This product must be stored in a locked room or place away from children, animals, food, feedstuffs, seed and fertilisers. Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt. Do not use empty container for any other purpose.

110L Mini Bulk Returnable Container (110L only)

This product must be stored in a locked room or place away from children, animals, food, feedstuffs, seed and fertilisers. Store the original sealed drum in a cool well-ventilated area. DO NOT store for prolonged periods in direct sunlight. DO NOT tamper with the non return valve or the security seal. DO NOT contaminate the drum with water or any foreign matter. After each use of the product, please ensure that the non-return valve, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the drum have been used, please return the empty drum to the point of purchase.

Refillable containers (1000L only):

This product must be stored in a locked room or place away from children, animals, food, feedstuffs, seed and fertilisers. Store in the closed, original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight. Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Product is poisonous if inhaled or swallowed. Attacks the eyes. Will irritate the skin. Avoid contact with eyes and skin. Do not inhale spray mist. When preparing the spray and using the prepared spray wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow-length PVC gloves and face shield or goggles. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (telephone 13 11 26). If swallowed, do NOT induce vomiting. Give glass of water.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet, which can be obtained from www.kenso.com.au.

CONDITIONS OF SALE

Kenso Corporation (M) Sdn. Bhd. will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Kenso Corporation (M) Sdn. Bhd. is limited to the replacement of the goods or (at the option of Kenso Corporation (M) Sdn. Bhd.) the refund of the price paid and where possible sufficient part of the goods to enable proper examination being returned to Kenso Corporation (M) Sdn. Bhd. within thirty days of delivery.

In a Transport Emergency
Dial **000**
Police or Fire Brigade



Additional statements required by Globally Harmonised System of Classification and Labelling of Chemicals (GHS) and Safe Work Australia: **COMBUSTIBLE LIQUID. HARMFUL IF SWALLOWED. MAY BE FATAL IF SWALLOWED AND ENTERS AIRWAYS. HARMFUL IN CONTACT WITH SKIN. MAY CAUSE AN ALLERGIC SKIN REACTION. TOXIC IF INHALED. SUSPECTED OF DAMAGING FERTILITY OR THE UNBORN CHILD.**
Precautionary: Obtain special instructions before use. Do not handle until all safety precautions have been read and understood. Keep away from heat/hot surfaces/sparks/open flames and other ignition sources. No smoking. Avoid breathing dust/fume/gas/mist/vapours/spray. Wash contacted area thoroughly after handling. Do not eat, drink or smoke when using this product. Use only outdoors or in a well-ventilated area. Contaminated work clothing should not be allowed out of the workplace. Wear protective gloves/protective clothing/eye protection/face protection. IF SWALLOWED: Immediately call a POISON CENTER or doctor/physician. IF ON SKIN: Wash with plenty of soap and water. IF INHALED: Remove person to fresh air and keep comfortable for breathing. IF exposed or concerned: Get medical advice/attention. Call a POISON CENTER or doctor/physician. Specific treatment (see FIRST AID on this label). Rinse mouth. Do NOT induce vomiting. If skin irritation or rash occurs: Get medical advice/attention. Take off contaminated clothing and wash it before reuse. In case of fire: Use water fog, foam, dry agent (carbon dioxide, dry chemical powder) to extinguish. Store in a well-ventilated place. Keep container tightly closed. Store locked up.

Batch No.:

Date of Manufacture:

DANGEROUS POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

KENSO
agcare

BROMOXYNIL
M

Selective Herbicide

ACTIVE CONSTITUENT: 200 g/L BROMOXYNIL present as the OCTANOATE

200 g/L MCPA present as the ETHYL HEXYL ESTER
SOLVENT: 324 g/L LIQUID HYDROCARBON

GROUP 6 | 4 HERBICIDE

For control of broadleaf weeds in cereals, linseed and grass pastures as specified in the DIRECTIONS FOR USE Table.

READ THIS LEAFLET BEFORE USE

APVMA Approval No.: 66163/RV2024



Kenso Corporation (M) Sdn. Bhd.
Level 1, 98 Commercial Road,
Teneriffe QLD 4005
Phone (07) 3216 1188

GENERAL INSTRUCTIONS

Resistant Weeds Warning

GROUP 6 | 4 HERBICIDE

Bromoxynil M Selective Herbicide is a member of the nitrile and phenoxy groups of herbicides. Bromoxynil M has both the inhibitor of photosynthesis at photosystem II and disruptor of plant cell growth modes of action. For weed resistance management Bromoxynil M is a Group 6 and 4 herbicide. Some naturally occurring weed biotypes resistant to Bromoxynil M and other Group 6 and 4 herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Bromoxynil M or other Group 6 and 4 herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Kenso Corporation (M) Sdn Bhd accepts no liability for any losses that may result from the failure of Bromoxynil M to control resistant weeds.

Note

Maximum daily temperatures at or above 20°C at application, or on days following treatment, may result in seedling mortality and established plants may be damaged.

Application

Mixing

To ensure even mixing, half fill the spray tank with clean water and add the required amount of product. Agitate thoroughly, then add the remainder of the water. Agitate again before spraying commences.

Boom Sprayer

Calibrate the machine before use to ensure that the correct application rate is obtained. Check height of boom above the crop to ensure that the spray is evenly distributed. Wash the spray equipment thoroughly after use.

Cereals

Apply 50 to 200 litres of water per hectare. A minimum of 160 litres and preferably 200 litres of water per hectare is recommended where the weed infestation is heavy or the crop cover is dense. Complete coverage of weeds is essential.

Linseed

Apply a minimum of 220 litres of water per hectare.

STORAGE AND DISPOSAL

This product must be stored in a locked room or place away from children, animals, food, feedstuffs, seed and fertilisers. Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt. Do not use empty container for any other purpose.

110L Mini Bulk Returnable Container

This product must be stored in a locked room or place away from children, animals, food, feedstuffs, seed and fertilisers. Store the original sealed drum in a cool well-ventilated area. DO NOT store for prolonged periods in direct sunlight. DO NOT tamper with the non return valve or the security seal. DO NOT contaminate the drum with water or any foreign matter. After each use of the product, please ensure that the non-return valve, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the drum have been used, please return the empty drum to the point of purchase.

Refillable containers (1000L only):

This product must be stored in a locked room or place away from children, animals, food, feedstuffs, seed and fertilisers. Store in the closed, original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight. Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Product is poisonous if inhaled or swallowed. Attacks the eyes. Will irritate the skin. Avoid contact with eyes and skin. Do not inhale spray mist. When preparing the spray and using the prepared spray wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow-length PVC gloves and face shield or goggles. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (telephone 13 11 26). If swallowed, do NOT induce vomiting. Give glass of water.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet, which can be obtained from www.kenso.com.au.

Aerial Application

Apply a minimum of 22 litres of water per hectare. Markers should always move upwind before aircraft passes overhead. Application by microneaire equipment may give variable results.

Compatibility

This product is physically compatible in the spray tank with one of the following herbicides and insecticides in wheat and barley only: metsulfuron, chlorpyrifos, dicamba, dimethoate, endosulfan, diclofop-methyl, MCPA and omethoate. When mixed with metsulfuron or dicamba, do not apply to cereals undersown with lucerne. When mixed with diclofop-methyl, do not exceed 1.4 litres of this product per hectare. Observe all instructions on the mixture partner label. As formulations of other manufacturers' products are beyond the control of Kenso Corporation (M) Sdn Bhd, all mixtures should be tested prior to mixing commercial quantities.

PROTECTION OF LIVESTOCK

Grazing Precaution

Sprayed weeds may become more palatable to stock and a higher intake of some weeds may result in stock poisoning and death from causes such as nitrate poisoning. Care should be taken especially where capeweed, Paterson's curse and variegated thistles predominate in the pasture. Avoid grazing with young or breeding stock. Do not graze horses or pigs on Paterson's curse. If in doubt, contact your nearest Department of Agriculture.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions or from spraying equipment which could be expected to cause spray to drift onto nearby plants, adjacent crops, crop lands or pastures. Avoid spray drift and vapour movement onto susceptible crops such as cotton, tobacco, tomatoes, vines, lupins, fruit trees and ornamentals. DO NOT apply by misting machines.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Dangerous to fish. DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

CONDITIONS OF SALE

Kenso Corporation (M) Sdn. Bhd. will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Kenso Corporation (M) Sdn. Bhd. is limited to the replacement of the goods or (at the option of Kenso Corporation (M) Sdn. Bhd.) the refund of the price paid and where possible sufficient part of the goods to enable proper examination being returned to Kenso Corporation (M) Sdn. Bhd. within thirty days of delivery.

* is not a registered trademarks of Kenso Corporation (M) Sdn. Bhd.

In a Transport Emergency
Dial **000**
Police of Fire Brigade



Kenso Corporation (M) Sdn. Bhd.
Level 1, 98 Commercial Road,
Teneriffe QLD 4005
Phone (07) 3216 1188

APVMA Approval No.: 66163/RV2024

DIRECTIONS FOR USE:
RESTRAINTS:

DO NOT apply if crop, pasture or weeds are stressed due to dry or excessively moist conditions.
DO NOT apply to diseased or frost-affected crops or if frosts are imminent.
DO NOT apply if rain is expected within three hours.

CROP / SITUATION	WEEDS CONTROLLED		STAGE OF WEED GROWTH	STATE	RATE	CRITICAL COMMENTS
Wheat, Barley, Cereal Rye, Oats and Triticale	Amsinckia	(<i>Amsinckia spp.</i>)	Plants up to the 6 leaf stage but not more than 50 mm in diameter	WA only	750 mL/ha	CROP STAGE: Wheat, Barley, Cereal Rye, Oats and Triticale: 3 leaf to fully tillered stage (Zadok Scale Z13-30) for rates of 750 mL to 1.4 L/ha; 5 leaf to fully tillered stage (Z15-30) for rate of 2.0 L/ha (NSW, Vic, SA, WA, Tas only); or early tillering to fully tillered stage (Z21-30) (Qld only). Linseed: 50 to 150 mm high. DO NOT apply to linseed by aircraft. * If spraying Black Bindweed when wheat, barley, cereal rye, oats, triticale or linseed is more than 300 mm high, increase spray volume to 200 L/ha. For optimal control, apply only when the weeds are actively growing. In the early stages following spraying, temporary wilting of crop and pasture may occur. Crop scorch may occur, particularly on oats, if treatment is delayed to later stages of growth. Refer to the General Instructions. Avoid application when maximum daily temperatures above 20°C occur, or are likely to occur for a few days after application. DO NOT apply to crops or pastures undersown with lucerne, clover or medics unless some damage is acceptable.
	Capeweed	(<i>Arctotheca calendula</i>)		Qld, NSW, Vic, Tas, SA, NT, ACT only	1.4 L/ha	
	Chamomile	(<i>Matricaria matricarioides</i>)			2.0 L/ha	
	Charlock	(<i>Sinapis arvensis</i>)				
Linseed	Corn Gromwell	(<i>Buglossoides arvensis</i>)	Plants up to the 8 leaf stage but not more than 75 mm in diameter	WA only		1.4 L/ha
	(Ironweed, Sheepweed)				2.0 L/ha	
Grass Pastures	Fat Hen	(<i>Chenopodium album</i>)	Plants up to the 4 leaf stage but not more than 35 mm in diameter	All States	1.4 L/ha	
	Field Madder	(<i>Sherardia arvensis</i>)				
	Hexham Scent	(<i>Melilotus indica</i>)				
	(King Island Melilot)					
	Horned Poppy	(<i>Glaucium flavum</i>)				
	Indian Hedge	(<i>Sisymbrium orientale</i>)				
	Mustard					
	Lesser	(<i>Coronopus didymus</i>)				
	Swinecress					
	Mexican Poppy	(<i>Argemone ochroleuca</i>)				
	Mintweed	(<i>Salvia reflexa</i>)				
	Paterson's	(<i>Echium plantagineum</i>)				
	Curse					
	(Salvation Jane)					
	Rough Poppy	(<i>Papaver hybridum</i>)				
	Saffron Thistle	(<i>Carthamus lanatus</i>)				
Shepherd's	(<i>Capsella bursa-pastoris</i>)	Plants up to the 8 leaf stage but not more than 75 mm in diameter		2.0 L/ha		
Purse						
Slender	(<i>Carduus tenuiflorus</i> ,					
Thistles	<i>C.pycnocephalus</i>)					
Tree Hogweed	(<i>Polygonum patulum</i>)					
	(<i>Rapistrum rugosum</i>)					
Turnip Weed	(<i>Rapistrum rugosum</i>)					
(Giant Mustard)						
Wild Radish	(<i>Raphanus raphanistrum</i>)					
Wild Turnip	(<i>Brassica tournefortii</i>)					
	Black Bindweed*	(<i>Fallopia convolvulus</i>)	Plants up to the 4 leaf stage but not more than 35 mm in diameter		1.0 L/ha	
	(Climbing Buckwheat)					
	Common	(<i>Lepidium africanum</i>)	Plants up to the 6 leaf stage but not more than 50 mm in diameter	All States	1.4 L/ha	
	Peppergrass					
	Mustards	(<i>Sisymbrium spp.</i>)	Plants up to the 8 leaf stage but not more than 75 mm in diameter		2.0 L/ha	
	Purple Calandrinia	(<i>Calandrinia menziesii</i>)				

CROP / SITUATION	WEEDS CONTROLLED	STAGE OF WEED GROWTH	STATE	RATE	CRITICAL COMMENTS	
Wheat, Barley, Cereal Rye, Oats and Triticale Linseed Grass Pastures	Fumitory (<i>Fumaria spp.</i>) Threecornered Jack (Doublegee, Spiny Emex)	Plants up to the 3 leaf stage but not more than 50 mm in diameter	Qld, NSW, Vic, SA, Tas only	1.4 L/ha	CROP STAGE: Wheat, Barley, Cereal Rye, Oats and Triticale: 3 leaf to fully tillered stage (Zadok Scale Z13-30) for rates of 750 mL to 1.4 L/ha; 5 leaf to fully tillered stage (Z15-30) for rate of 2.0 L/ha (NSW, Vic, SA, WA, Tas only); or early tillering to fully tillered stage (Z21-30) (Qld only). Linseed: 50 to 150 mm high. DO NOT apply to linseed by aircraft. For optimal control, apply only when the weeds are actively growing. In the early stages following spraying, temporary wilting of crop and pasture may occur. Crop scorch may occur, particularly on oats, if treatment is delayed to later stages of growth. Refer to the General Instructions. Avoid application when maximum daily temperatures above 20°C occur, or are likely to occur for a few days after application. DO NOT apply to crops or pastures undersown with lucerne, clover or medics unless some damage is acceptable. ** Where Wireweed occurs in red soils of low fertility, it has been found to be less susceptible. In these circumstances use 2.0L/ha.	
		Plants up to the 4 leaf stage but not more than 60 mm in diameter	WA only	750 mL/ha		
		Plants up to the 5 leaf stage but not more than 75 mm in diameter	Qld, NSW, Vic, Tas, SA, NT, ACT only	2.0 L/ha		
		Plants up to the 6 leaf stage but not more than 85 mm in diameter	WA only	1.4 L/ha		
	Variegated Thistle (<i>Silybum marianum</i>) Wireweed (<i>Polygonum aviculare</i>) (Hogweed)**	Plants up to the 4 leaf stage but not more than 35 mm in diameter	Qld, NSW, Vic, Tas, SA, NT, ACT only	750 mL/ha		
		Plants up to the 6 leaf stage but not more than 50 mm in diameter		1.4 L/ha		
		Plants up to the 6 leaf stage		2.0 L/ha		
	Mountain Sorrel (<i>Oxalis acetosella</i>) Three-horned Bedstraw (<i>Galium tricornutum</i>)	Plants up to the 6 leaf stage	WA only	1.4 L/ha		
		Plants up to the 8 leaf stage		750 mL/ha		
	Cleavers (<i>Galium aparine</i>)	2-4 stem stage and 1-3 whorls of leaves per stem		750 mL/ha		

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WITHHOLDING PERIOD

Harvest

Cereals, linseed: NOT REQUIRED WHEN USED AS DIRECTED

Grazing

Cereals, pasture: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 8 WEEKS AFTER APPLICATION.