

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

KENSO AGCARE

OX 240

HERBICIDE

ACTIVE CONSTITUENT: 240 g/L OXYFLUORFEN

SOLVENTS: 608 g/L LIQUID HYDROCARBON

100 g/L N-METHYL PYRROLIDONE

GROUP 14 HERBICIDE

For the selective control of certain broadleaf and grass weeds as per
Directions for Use table.

Kenso Corporation (M) Sdn Bhd
Level 1, 98 Commercial Road,
Teneriffe QLD 4005
Phone (07) 3216 1188
www.kenso.com.au



IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USE

CONTENTS: 10 Litres APVMA Approval No.: 61270/ 107912

KENSO AGCARE OX 240 HERBICIDE

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. DO NOT burn empty container or product. For refillable containers, empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Will damage eyes and irritate the skin. Avoid contact with eyes and skin. When opening the container and preparing spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and face shield or goggles. If product on skin, immediately wash area with soap and water. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use wash gloves, goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126; New Zealand 0800 764 766. If swallowed, do NOT induce vomiting. Give a glass of water. If in eyes wash out immediately with water.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet, which can be obtained from www.kenso.com.au.

CONDITIONS OF SALE

Kenso Corporation (M) Sdn. Bhd. will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection

with the supply of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Kenso Corporation (M) Sdn. Bhd. is limited to the replacement of the goods or (at the option of Kenso Corporation (M) Sdn. Bhd.) the refund of the price paid and where possible sufficient part of the goods to enable proper examination being returned to Kenso Corporation (M) Sdn. Bhd. within thirty days of delivery.

In a Transport Emergency
Dial **000**
Police or Fire Brigade



Additional statements required by Globally Harmonised System of Classification and Labelling of Chemicals (GHS) and Safe Work Australia: **MAY BE FATAL IF SWALLOWED AND ENTERS AIRWAYS. CAUSES SKIN IRRITATION. CAUSES SERIOUS EYE IRRITATION. MAY DAMAGE FERTILITY OR THE UNBORN CHILD.** Precautionary: Obtain special instruction before use. Do not handle until all safety precautions have been read and understood. Wash contacted areas thoroughly after handling. Wear protective gloves, protective clothing and eye or face protection. IF SWALLOWED: Immediately call a POISON CENTER or doctor/physician. IF ON SKIN: Wash with plenty of soap and water. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. IF exposed or concerned: Get medical advice/attention. Specific treatment (see FIRST AID on this label). DO NOT induce vomiting. If skin irritation occurs: Get medical advice/attention. If eye irritation persists: Get medical advice/attention. Take off contaminated clothing and wash it before reuse. Store locked up.

Batch No.:

Date of Manufacture:

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

KENSO
ag-gear

OX 240

Herbicide

ACTIVE CONSTITUENT: 240 g/L OXYFLUORFEN
SOLVENT: 608 g/L LIQUID HYDROCARBON
100 g/L N-METHYL PYRROLIDONE

GROUP 14 HERBICIDE

For the selective control of certain broadleaf and grass weeds as per the Directions for Use table.

READ THIS LEAFLET BEFORE USE

APVMA Approval No.: 61270/107912



Kenso Corporation (M) Sdn. Bhd.
Level 1, 98 Commercial Road,
Teneriffe QLD 4005
Phone (07) 3216 1188

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. DO NOT burn empty container or product.

For refillable containers, empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Will damage eyes and irritate the skin. Avoid contact with eyes and skin. When opening the container and preparing spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and face shield or goggles. If product on skin, immediately wash area with soap and water. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use wash gloves, goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126; New Zealand 0800 764 766. If swallowed, do NOT induce vomiting. Give a glass of water. If in eyes wash out immediately with water.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet, which can be obtained deom www.kenso.com.au.

CONDITIONS OF SALE

Kenso Corporation (M) Sdn. Bhd. will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Kenso Corporation (M) Sdn. Bhd. is limited to the replacement of the goods or (at the option of Kenso Corporation (M) Sdn. Bhd.) the refund of the price paid and where possible sufficient part of the goods to enable proper examination being returned to Kenso Corporation (M) Sdn. Bhd. within thirty days of delivery.

In a Transport Emergency
Dial **000**
Police of Fire Brigade

* Not a registered trademark of Kenso Corporation (M) Sdn. Bhd.



Kenso Corporation (M) Sdn. Bhd.
Level 1, 98 Commercial Road,
Teneriffe QLD 4005
Phone (07) 3216 1188

APVMA Approval No.: 61270/107912

Crop	Weed Controlled	Time of Application	Rate	Critical Comment
Tree fruit, Nuts and Vines not less than 3 years old DORMANT APPLICATION Almonds, Apples, Apricots, Grapevine, Macadamias, Peaches, Plums, Pears, Pecans, Walnuts	Refer to Weeds Controlled list	Weed free soil	3 or 4 L/ha	DO NOT apply once budswell has occurred. Apply to freshly cultivated weed free soil. Use higher rate for longer residual control (up to 4 months). Where grass weeds are expected to be the major problem, or when control of a wider weed spectrum is needed, mix the lower rate with 4.5L Surflan* 500 (500 g/L) or 4.5kg Devrinol* WG (500 g/kg) per treated hectare. Refer to individual product labels for crops, rates, states and weeds controlled.
		Weeds at 4-6 true leaf stage		DO NOT apply once budswell has occurred. Use higher rate for longer residual control (up to 4 months). When young seedling grasses and/or broadleaf weeds are present, apply as a tank mix with Para-Ken 250, Spray Seed*, Diquat 200, Speedy 250, Ken-Up®, Ken-Up® 450 CT, Max Out 540, Weedmaster* Duo, Credit* + Bonus* to give both knockdown and residual control. Refer to individual product labels for rates. Addition of a non-ionic surfactant such as Activator Surfactant should be used in the spray mixture at 125 mL/100L. Mature, established weeds must be eliminated prior to application by mechanical or chemical means. Macadamias: Apply in 250 to 500L water/ha. Apply after harvest to prevent spray contacting nuts. Avoid spray contact with the macadamia tree foliage and stem. DO NOT apply to nuts on the ground.
Almonds, Apples, Apricots, Grapevine, Macadamias, Peaches, Plums, Pears, Pecans, Walnuts	Refer to Speedy 250 label	Emerged weeds present	250 mL/ha plus Speedy 250 at recommended label rate OR 25 to 30 mL/100L plus Speedy 250 at recommended label rates	The addition of Speedy 250 will improve control of small flowered mallow, evening primrose and other weeds sensitive to Ox 240. Apply as a tank mix with Speedy 250 refer to the Speedy 250 label and follow all label directions.
Almonds, Apples, Apricots, Grapevine, Macadamias, Peaches, Pears, Pecans, Walnuts	Refer to Ken-Up®, Ken-Up® 450 CT, Max Out 540, Weedmaster* Duo, Credit* + Bonus* label	Apply as a tank mix with glyphosate products	75 mL/ha plus Ken-Up® 450 CT, Max Out 540, Weedmaster* Duo or Credit* + Bonus* at recommended label rates	Addition of Ox 240 to glyphosate products will improve knockdown and increase the speed at which treated weeds develop visible symptoms of phytotoxicity (compared to results achieved with glyphosate applied alone) and give control of annual nettles <i>Urtica spp.</i> , <i>Marshmallow</i> , <i>Malva parviflora</i> . For rates refer to the appropriate label. Read and follow all label directions. DO NOT apply the tank mix of glyphosate and Ox 240 near trees or vines less than 3 years old unless they are effectively shielded from spray and spray drift.
Pyrethrum – Greater than 4 leaf	Blackberry nightshade (<i>Solanum nigrum</i>) Fumitory (<i>Fumaria spp.</i>) Hogweed (<i>Polygonum aviculare</i>) Sorrel (<i>Rumex acetosella</i>) Volunteer potato (<i>Solanum tuberosum</i>) Cleavers (<i>Galium aparine</i>) Field Bindweed (<i>Convolvulus vulgaris</i>) Groundsel (<i>Senecio vulgaris</i>)	Emerged weeds present	100 to 150 mL/ha	Apply when pyrethrum growth stage is greater than 4 true leaf.
Pyrethrum – established crops, greater than 1 year old	As above plus Sow Thistle (<i>Sonchus spp.</i>) Spear Thistle (<i>Cirsium vulgare</i>)	Emerged weeds present	200 mL to 4 L/ha	Apply when pyrethrum is greater than 100 mm rosettes. Apply rates of greater than 1.0 L/ha ONLY between 1 st of February and 31 st of March. DO NOT apply later than 25 weeks before harvest.

Crop	Weed Controlled	Time of Application	Rate	Critical Comment
Duboisia	Refer to Weeds Controlled list	Weed free soil	4 to 8 L/ha	Apply after transplantation of young seedlings or after harvest of mature plants. Use higher rate where longer residual control (up to 6 months) is required. “Over-the-top” application will be tolerated.
		Weeds at 4 to 6 true leaf stage		Best results are achieved when applied to moist soil free of weeds. However, some newly germinated small seedling grasses and broadleaf weeds (4 to 6 true leaves) will be controlled at these rates. Addition of a non-ionic surfactant such as Ken-Wett 1000 or Activator Surfactant at recommended rates will improve activity. Established weeds must be eliminated prior to application by mechanical means.
Tobacco	Refer to Weeds Controlled list	Weed free soil	4 L/ha	Use to control weeds along spray lines only. DO NOT apply to tobacco crop. Apply to soil after solid-set irrigation system has been laid out in the field. Ox 240 should be applied to moist soil. Where very small weeds (2 to 3 leaf) emerge prior to spraying, the addition of a wetting agent to the spray mixture is necessary for effective control. Should the weeds be more advanced, the addition of 2 L/ha of a registered product containing both 75g/L diquat & 125g/L paraquat. Avoid spray drift.
Coffee	Refer to Weeds Controlled list	Weed free soil	2 to 4 L/ha	Apply after transplanting coffee plants or under established bushes. Use the higher rate where longer residual activity (up to 4 months) is required. Best results are achieved when applied to moist soil, free of weeds with follow-up irrigation.
		Weeds at 2 to 4 true leaf stage		When seedlings are present, apply as a tank mix with Para-Ken 250 to produce both knockdown and residual control. Addition of a non-ionic surfactant such as Ken-Wett 1000 at 150 mL/100L or Activator Surfactant at 125 mL/100L should be used in the spray mixture. Apply as directed spray to avoid contact with coffee plants. Mature established weeds must be eliminated by mechanical or chemical means prior to application.
Onions (seeded)	Blackberry Nightshade (<i>Solanum nigrum</i>)	Hook leaf stage	50 to 100 mL/ha	Do not exceed 100 mL/ha at the hook leaf stage as excessive crop damage may occur. Best results are obtained when the hook leaf stage treatment is followed by a further application of Ox 240 when onions are at the 1 ½ to 2 ½ leaf stage. Apply Ox 240 in at least 100 L/ha water at pressure below 150 kPa. DO NOT exceed 150 kPa because excessive crop damage could result. Multiple treatments may be applied provided the total dose does not exceed 2 L/ha in one season. For control of other weeds, use in a program with other registered herbicides. Note: Can cause flecking, twisting or stunting of onion plants although such symptoms are normally short-lived. DO NOT apply in conditions of very high humidity. DO NOT apply to weeds or crops which are stressed by prevailing weather conditions, disease or mechanical damage. DO NOT use wetting agents or other adjuvants.
	Common Cotula (<i>Cotula australis</i>) Crowsfoot / Storksbill (<i>Erodium spp.</i>) Deadnettle (<i>Lamium amplexicaule</i>) Docks (<i>Rumex spp.</i>) Fumitory (<i>Pink and white flowered</i>) (<i>Fumaria spp.</i>) Groundsel (<i>Senecio vulgaris</i>) Hogweed / Wireweed (<i>Polygonum aviculare</i>) Milk Thistle (<i>Sonchus oleraceus</i>) Plantain (<i>Plantago spp.</i>) Sorrel (<i>Rumex acetosella</i>) Potato weed (<i>Galinsoga parviflora</i>) Volunteer potato (<i>Solanum tuberosum</i>)	1 ½ to 2 ½ leaf stage	350-500 mL/ha	

Crop	Weed Controlled	Time of Application	Rate	Critical comment
Tropical and Subtropical fruit <i>(inedible peel)</i> Custard Apple, Kiwi Fruit, Longan, Lychees, Mango, Passionfruit, Pawpaw, Rambutan	Refer to Weeds Controlled list	Weed free soil	4 L/ha	Best results are obtained when applied to most soil free of weeds.
		Emerged weeds present		If weeds are present Ox 240 should be applied as a tank mix with the recommended rates of Ken-Up ® 450 CT, Max Out 540, Weedmaster* Duo, Credit* + Bonus*, Speedy 250 or Para-Ken 250.
Brassica Crops, Broccoli, Cabbages, Cauliflower	Refer to Weeds Controlled list	Weed free soil	1.5 to 2 L/ha	Apply Ox 240 to prepared ground 4 to 7 days prior to transplanting. If soil is dry irrigation or rainfall is required prior to transplanting for activation of Ox 240. Utilise transplanting techniques which cause minimal soil disturbance. Excessive soil disturbance will lessen herbicide activity. Use the higher rate in situations where weed pressure is known to be heavy.
Forestry – <i>Eucalyptus</i> spp. <i>Pinus</i> spp.	Refer to Weeds Controlled list	Weed free soil	3 or 4 L/ha	Apply to weed free soil either before or preferably soon after transplanting. Disturbance of the herbicidal barrier on the soil surface by the transplantation process may reduce the length of weed control. Use higher rate for longer residual control. Apply as a directed or over-the-top spray. Ox 240 may be tank mixed with Simazine 900 DF to extend the spectrum and length of weed control. Application to foliage that is soft and has not hardened off and/or if temperatures exceed 20°C may increase the likelihood of foliar damage to trees eucalypts especially Ox 240 has been found to be safe to commonly planted species used in forestry. Sensitivity of less common species should be checked prior to large scale application.
		Weeds at 4-6 true leaf stage		Apply as an over-the-top spray to eucalypts and pines. Best results are achieved when applied to moist soil free of weeds. However, some newly germinated small seedling grasses and broadleaf weeds (4 to 6 true leaves) will be controlled at these rates. Addition of a non-ionic surfactant such as Activator Surfactant at recommended rates will improve activity. Established weeds must be eliminated prior to application. Ox 240 may be tank mixed with Simazine 900 DF to extend the spectrum and length of weed control. Application to foliage that is soft and has not hardened off and/or if temperatures exceed 20°C may increase the likelihood of foliar damage to trees, eucalypts especially. Ox 240 has been found to be safe to commonly planted species used in forestry. Sensitivity of less common species should be checked prior to large scale application.
Forestry Trees	Broadleaf weeds and grasses	Weeds at 4-6 true leaf stage	4 L/ha or 4 mL/10m ²	For the establishment of trees of approved farm practices such as wind breaks, erosion control, woodlots and forestry plantings. When applying as a post-plant spray, ensure spray is directed to the base of seedlings, or that seedlings are protected. Do not apply under hot or windy conditions.
Prior to sowing winter cereals	Refer to Ken-Up [®] , Ken-Up [®] 450 CT, Max Out 540, Weedmaster* Duo, Credit* + Bonus* label	Apply as a tank mix with glyphosate products	75 mL/ha plus Ken-Up [®] 450 CT, Max Out 540, Weedmaster* Duo or Credit* + Bonus* at recommended label rates	Addition of Ox 240 to glyphosate products will improve knockdown and increase the speed at which treated weeds develop visible symptoms of phytotoxicity (compared to results achieved with glyphosate applied alone) and give control of annual nettles <i>Urtica spp.</i> , and <i>Marshmallow, Malva parviflora</i> . For rates refer to the appropriate label. Read and follow all label directions.
Prior to sowing cotton				

When using Ox 240 in combination with other herbicides, refer to the appropriate label and reads follow all label directions.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

THIS PRODUCT IS TOO HAZARDOUS FOR USE IN THE HOME GARDEN.

WITHHOLDING PERIODS:

GRAZING - DO NOT GRAZE TREATED WEEDS

HARVEST - NOT REQUIRED WHEN USED AS DIRECTED

WEEDS CONTROLLED		
BEFORE GERMINATION	WEEDLINGS	
Amsinckia <i>Amsinckia</i> spp.	Prickly lettuce <i>Lactuca</i> spp.	Amsinckia <i>Amsinckia</i> spp.
Barley grass <i>Hordeum leporinum</i>	Red natal grass <i>Rhynchelytrum repens</i>	Belline <i>Ipomoea</i> spp.
Barnyard grass <i>Echinochloa</i> spp.	Redshank <i>Amaranthus cruentus</i>	Capeweed <i>Arctotheca calendula</i>
Blackberry nightshade <i>Solanum nigrum</i>	Ryegrass <i>Lolium</i> spp.	Common cotula <i>Cotula australis</i>
Bladder ketmia <i>Hibiscus trionum</i>	Sesbania pea <i>Sesbania cannabina</i>	Crowfoot grass <i>Eleusine indica</i>
Burrgrass <i>Cenchrus australis</i>	Shepherd’s purse <i>Capsella bursa-pastoris</i>	Deadnettle <i>Lamium amplexicaule</i>
Caltrop <i>Tribulus terrestris</i>	Smallflower mallow <i>Malva parviflora</i>	Groundsel <i>Senecio vulgaris</i>
Capeweed <i>Arctotheca calendula</i>	Soursob <i>Oxalis pes-caprae</i>	Liverseed grass <i>Urochloa panicoides</i>
Chickweed <i>Stellaria media</i>	Sowthistle <i>Sonchus oleraceus</i>	Marshmallow <i>Malva parviflora</i>
Crowsfoot grass <i>Eleusine indica</i>	Starburr <i>Acanthospermum hispidum</i>	Pigweed <i>Portulaca oleracea</i>
Deadnettle <i>Lamium amplexicaule</i>	Stinkgrass <i>Eragrostis cilianensis</i>	Potato weed <i>Galinsoga parviflora</i>
Fat hen <i>Chenopodium album</i>	Summer grass <i>Digitaria</i> spp.	Redshank <i>Amaranthus cruentus</i>
Giant pigweed <i>Trianthema portulacastrum</i>	Thornapple <i>Datura stramonium</i>	Shepherd's purse <i>Capsella bursa-pastoris</i>
Liverseed grass <i>Urochloa panicoides</i>	White eye <i>Richardia brasiliensis</i>	Sowthistle <i>Sonchus olearaceus</i>
Lovergrass <i>Eragrostis</i> spp.	Wild mustard <i>Sisymbrium</i> spp.	Stinging nettle <i>Urtica urens</i>
Pigeon grass <i>Setaria</i> spp.	Wild radish <i>Raphanus raphanistrum</i>	Stinkgrass <i>Eragrostis cilianensis</i>
Pigweed <i>Portulaca oleracea</i>	Wireweed <i>Polygonum aviculare</i>	Wild radish <i>Raphanus raphanistrum</i>

GENERAL INSTRUCTIONS

Ox 240 is a selective herbicide for the control of certain annual grasses and broadleaf weeds in dormant apples, grapes, peaches, pears, plums, apricots, almonds, macadamia, duboisia, tobacco, coffee, pyrethrum, winter cereals, tropical/sub tropical crops, brassicas, onions, Eucalyptus and Pinus spp. and other forestry trees used in forestry applied either to a weed free soil or to seedling weeds at the 4 to 6 true leaf stage.

Ox 240 applied to well prepared weed free soil should not be disturbed or incorporated after application. Weed control for up to 6 months can be expected, but spot treatment with knockdown herbicides, of escape weeds and perennial grasses may be necessary.

When Ox 240 is applied to seedling weeds at the 4 to 6 true leaf stage a non-ionic surfactant such as non-ionic surfactant or Ken-Wett 1000 should be added at recommended rate to improve activity.

Ox 240 can also be utilized at low rates to improve the weed spectrum of knockdown herbicides.

Duboisia seedlings and mature plants will tolerate “over the top” applications. Eucalypt and pine transplants and established trees can tolerate “over the top” applications.

TIMING

For optimum residual weed control, Ox 240 should be applied to the soil surface prior to weed emergence after all other agricultural operations have been completed, such as mechanical cultivation and reshaping of irrigation furrows. The area should be left undisturbed during the period of desired weed control. When applied to seedling weeds, they should be young and actively growing. Weed control for up to 6 months is expected but spot treatment, with knockdown herbicides, for escape weeds and perennial grasses may be necessary.

MIXING

Shake well before use. When using Ox 240 alone, fill the spray tank, at least one-third full with clean water, add the recommended amount of Ox 240 while the pump and agitator are running, then complete filling the spray tank.

A non-ionic surfactant, if required by label directions, should be added near the end of the filling process to minimise foaming.

When tank mixing with Surflan* 500 or Devrinol, add to one-third filled tank, then the Ox 240 during the filling operation. Maintain agitation during mixing and until spraying is completed.

When tank mixing with Ken-Up® 450 CT, Max Out 540, Weedmaster* Duo, Credit* and Bonus*, Spray Seed* or Para-Ken 250, add these after Ox 240 during the filling operation.

When using Ox 240 in mixtures always refer to the appropriate label and read and follow all label directions.

APPLICATION

Ox 240 should be applied as a ground spray directed at the base of the crop plants. The spray mixture should be applied evenly with a carefully calibrated, standard low pressure herbicide sprayer with a flat fan or hollow cone nozzle. When seedling weeds are present ensure thorough coverage of their foliage is required to obtain the maximum knockdown effect. Ensure both the weed foliage and the soil surface are sprayed. When applying to bare soil use 250 to 500 L of water/ha. For application to seedling weeds (4 to 6 leaf stage) use 250 to 1350 L/ha. Use higher volumes for high weed density.

CROP SAFETY

Ox 240 may be applied as directed around dormant peach, plum, apricot, almond, apple and pear trees and grape vines of all ages when applied at rates of less than 1.0 L/ha. When applied at 3.0 l/ha and above, the trees and grape vines should be at least 3 years of age. DO NOT apply Ox 240 once bud swell has occurred when using rates greater than 1.0 L/ha. Duboisia seedlings and mature plants will tolerate “over-the-top” applications of Ox 240. When using the 75 mL/ha rate, Ox 240 may be applied as directed prior to sowing winter cereals.

COMPATIBILITY

Ox is compatible with Credit* + Bonus*, Naproamide, Speedy 250, Para-Ken 250, Diquat 200, Ken-Up®, Ken-Up® 450 CT, Max Out 540, Spray Seed*, Oryxalin, Glyphosate Trimesium, Weedmaster* Duo and Tryquat*.

RESISTANT WEEDS WARNING

GROUP 14**HERBICIDE**

Ox 240 is a member of the diphenyl ether group of herbicides. Ox 240 has the inhibitor of protoporphyrinogen oxidase mode of action. For weed resistance management Ox 240 is a Group 14 herbicide. Some naturally occurring weed biotypes resistant to Ox 240 and other Group 14 herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Ox 240 or other Group 14 herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, Kenso Corporation (M) Sdn Bhd accepts no liability for any losses that may result from the failure of Ox 240 to control resistant weeds.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby water bodies, susceptible plants/crops, cropping lands or pastures.

PROTECTION OF LIVESTOCK

Use with care when applying in areas frequented by stock.

PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND ENVIRONMENT

This product is highly toxic to wildlife and fish. DO NOT contaminate lakes, ponds, streams, rivers or waterways with the chemical or used containers. Use care when applying in areas frequented by wildlife or adjacent to any body of water. DO NOT apply when weather conditions favour drift from target areas.