

CAUTION

KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

KENSO AGCARE

PADDOM SC HERBICIDE

ACTIVE CONSTITUENT: 275 g/L TERBUTRYN
160 g/L MCPA present as the POTASSIUM SALT

GROUP **5 | 4** HERBICIDE

For the control of seedling broadleaf weeds in wheat, barley, some varieties of
oats, pasture and sugar cane as specified in the directions for use



Kenso Corporation (M) Sdn Bhd
Level 1, 98 Commercial Road,
Teneriffe QLD 4005
Phone (07) 3216 1188
www.kenso.com.au

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USE

CONTENTS: 20 Litres APVMA Approval No.: 86112/ 114704

KENSO AGCARE PADDO M SC HERBICIDE

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated locked area away from children, animals, food, feedstuffs, seed and fertilisers. Do not store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Harmful if swallowed. Attacks eyes. Will irritate the skin. Avoid contact with skin. Repeated exposure may cause allergic disorders. When preparing the spray and using the prepared spray wear cotton overalls buttoned to the neck and wrist and washable hat, elbow-length PVC gloves and goggles. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, goggles and contaminated clothing.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet, which can be obtained from www.kenso.com.au.

CONDITIONS OF SALE

Kenso Corporation (M) Sdn. Bhd. will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or

supply of these goods. To the extent allowed by such Statutes the liability of Kenso Corporation (M) Sdn. Bhd. is limited to the replacement of the goods or (at the option of Kenso Corporation (M) Sdn. Bhd.) the refund of the price paid and where possible sufficient part of the goods to enable proper examination being returned to Kenso Corporation (M) Sdn. Bhd. within thirty days of delivery.

In a Transport Emergency
Dial **000**
Police or Fire Brigade



Additional statements required by Globally Harmonised System of Classification and Labelling of Chemicals (GHS) and Safe Work Australia: **HARMFUL IF SWALLOWED. CAUSES SKIN IRRITATION. CAUSES SERIOUS EYE DAMAGE.** Precautionary: Wash contacted areas thoroughly after handling. Do not eat, drink or smoke when using this product. Wear protective gloves, protective clothing and eye or face protection. IF SWALLOWED: Call a POISON CENTER or doctor/physician if you feel unwell. IF ON SKIN: Wash with plenty of soap and water. Immediately call a POISON CENTER or doctor/physician. Specific treatment (see FIRST AID on this label). Rinse mouth. If skin irritation occurs: Get medical advice/attention. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. Take off contaminated clothing and wash before reuse.

Batch No.:

Date of Manufacture:

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

KENSO
agcare

**PADDO M
SC**

Herbicide

**ACTIVE CONSTITUENT: 275 g/L TERBUTRYN
160 g/L MCPA present as the POTASSIUM SALT**

GROUP 5 | 4 HERBICIDE

For the control of seedling broadleaf weeds in wheat,
barley, some varieties of oats, pasture and sugar
cane as specified in the directions for use

IMPORTANT: READ THIS LEAFLET BEFORE USE

APVMA Approval No.: 86112/114704



Kenso Corporation (M) Sdn. Bhd.
Level 1, 98 Commercial Road,
Teneriffe QLD 4005
Phone (07) 3216 1188

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND THE ENVIRONMENT

Dangerous to fish. DO NOT contaminate streams, rivers,
or waterways with the chemical or used containers.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions or from spraying
equipment that may cause spray to drift onto nearby
susceptible plants/crops, cropping lands or pastures.
Avoid spray drift onto susceptible crops such as cotton,
tobacco, tomatoes, vines, lupins, fruit trees and
ornamentals.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool,
well-ventilated locked area away from children, animals,
food, feedstuffs, seed and fertilisers. Do not store for
prolonged periods in direct sunlight. Triple or preferably
pressure rinse containers before disposal. Add rinsings to
spray tank. Do not dispose of undiluted chemicals on-site.
If recycling, replace cap and return clean containers to
recycler or designated collection point.
If not recycling, break, crush or puncture and bury empty
containers in a local authority landfill. If no landfill is
available, bury the containers below 500 mm in a disposal

pit specifically marked and set up for this purpose clear of
waterways, desirable vegetation and tree roots. Empty
containers and product should not be burnt.

SAFETY DIRECTIONS

Harmful if swallowed. Attacks eyes. Will irritate the skin.
Avoid contact with skin. Repeated exposure may cause
allergic disorders. When preparing the spray and using the
prepared spray wear cotton overalls buttoned to the neck
and wrist and washable hat, elbow-length PVC gloves and
goggles. If product in eyes, wash it out immediately with
water. After use and before eating, drinking or smoking,
wash hands, arms and face thoroughly with soap and
water. After each day's use, wash gloves, goggles and
contaminated clothing.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information
Centre. Phone Australia 13 11 26.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet,
which can be obtained from www.kenso.com.au.

CONDITIONS OF SALE

Kenso Corporation (M) Sdn. Bhd. will not accept any
responsibility whatsoever and howsoever arising and whether
for consequential loss or otherwise in connection with the
supply of these goods other than responsibility for the
merchantable quality of the goods and such responsibilities
mandatorily imposed by Statutes applicable to the sale or
supply of these goods. To the extent allowed by such Statutes
the liability of Kenso Corporation (M) Sdn. Bhd. is limited to the
replacement of the goods or (at the option of Kenso
Corporation (M) Sdn. Bhd.) the refund of the price paid and
where possible sufficient part of the goods to enable proper
examination being returned to Kenso Corporation (M) Sdn.
Bhd. within thirty days of delivery.

* Not a registered trademark of Kenso

In a Transport Emergency
Dial **000**
Police of Fire Brigade



Kenso Corporation (M) Sdn. Bhd.
Level 1, 98 Commercial Road,
Teneriffe QLD 4005
Phone (07) 3216 1188

APVMA Approval No.: 86112/114704

WITHHOLDING PERIOD:

**DO NOT GRAZE BARLEY, WHEAT, OATS OR PASTURE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER
APPLICATION.**

**DO NOT APPLY THE TANK MIX OF PADDO M SC AND AMETRYN (500 g/L) LATER THAN 9 MONTHS BEFORE
HARVEST**

GENERAL INSTRUCTIONS Resistant Weeds Warning

GROUP 5 | 4 HERBICIDE

Kenso Agcare Paddo M SC Herbicide is a member of both the triazines and phenoxy's groups of herbicides. Paddo M SC
has the inhibitor of photosynthesis at photosystem II and disruptor of plant cell growth modes of action. For weed resistance
management Paddo M SC is a group 5 and 4 herbicide.
Some naturally occurring weed biotypes resistant to Paddo M SC and other group 5 or 4 herbicides may exist through
normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if
these herbicides are used repeatedly. These resistant weeds will not be controlled by Paddo M SC or other inhibitors of
photosynthesis at photosystem II, or disruptor of plant cell growth herbicides.
Since the occurrence of resistant weeds is difficult to detect prior to use, Kenso Agcare Pty Ltd accepts no liability for any
losses that may result from the failure of Paddo M SC to control resistant weeds.

MIXING

Paddo M SC mixes readily with water. Half fill the spray tank with water. Shake the container several times before use and
add the recommended quantity to the tank. Keep the spray mixture well agitated during mixing and spraying using either
mechanical or venturi type agitation that has sufficient water movement for the solution to remain in suspension. Reseal
part-used containers immediately. Wash sprayer thoroughly with clean water after use.

APPLICATION

Boom Sprayer: Calibrate the sprayer before use to ensure the correct application rate is applied per hectare. Check boom
height for even application.

Cereal and pastures: Apply 50 to 100 L of water/ha using no smaller than 80 mesh pump and jet filters. Use the higher
volume if the weed infestation is heavy or the crop cover is dense.

Sugar cane: Apply 200 to 400 L of water/ha using no smaller than 80 mesh pump and jet filters. Use the higher volume if the
weed density is high or on more advanced weeds. Thorough coverage is necessary. Always add the required wetter.

Aircraft: Apply 20 to 30 L of water/ha. The higher volume is preferred due to more reliable results. Weed control following
aerial application may not be as good as from ground application.

COMPATIBILITY

This product can be tank mixed with Ken-Met 600 WG for the control of specific weeds such as capeweed, sheepweed and
toadrush. DO NOT tank mix with diclofop methyl or tralkoxydim as significant crop injury and reduced weed control will
occur. In pastures this product can be mixed with Kenso Agcare MCPA 750 and the insecticides dimethoate, phosmet,
Karate*, Ken-Tac 100 and omethoate (ensure good agitation is maintained with this product and avoid use of cold water). In
sugar cane this product can be tank mixed with Agral*, BS*1000 wetter, Ken-Amine 625 and ametryn (500 g/L). When tank
mixing read both labels and adhere to the crop recommendations and directions that are most restrictive. The compatibilities
given are based on experience and are considered correct. However, as changes in climatic conditions can alter the
sensitivity of plants to mixtures of sprays, Kenso Agcare Pty Ltd accepts no responsibility for the behaviour of such mixtures.

TIMING

Weeds should be sprayed while actively growing and at the 2 to 6 true leaf stage when they are most susceptible. As a
general guide Paddo M SC should be applied approximately 5 to 8 weeks after planting. Extremes of cold or drought are
unfavourable. The most favourable conditions are when soil is moist, weather fine and rain unlikely within 6 hours following
application. In pasture, weeds should be up to 20 cm diameter but refer to the directions for use table for specific details. In
sugar cane, apply Paddo M SC alone to broadleaf weeds up to the 8 leaf or 30 cm diameter stage, vines to 1m length.

CROP SAFETY

Research has shown that although some early growth check and leaf margin scorch can occur in oats, if grain yield is
measured, the following varieties are tolerant or moderately tolerant to Paddo M SC.

Tolerant: (apply up to 1.5 L/ha) Bandicoot, Coolabah, Mortlock, Potaroo, Swan, Wallaroo, West.

Moderately Tolerant: (DO NOT exceed 1 L/ha) Dalyup, Windjardie.

DO NOT spray other oat varieties unless you seek advice before spraying.

In pastures, sub clover growth may be reduced for a period after application but plants usually recover and spring growth is
not normally affected.

Sugar cane varieties such as Q122 may show yellowing and some leaf scorch immediately after spraying but plants
normally recover within 4 weeks.

DIRECTIONS FOR USE:

RESTRAINTS:

DO NOT apply to crops undersown with legumes.
DO NOT spray if weeds are wilted by dry or cold weather.
DO NOT spray if rain is expected within 6 hours.
DO NOT spray cereals or pastures on days when the temperature is greater than 22°C.
DO NOT add a wetting agent or a spraying oil for use in cereals or pasture.

Crop	Weeds Controlled	State	Rate /ha	Critical Comments
Sugar cane (plant and ratoon)	Bellvine Morning glory Red convolvulus Pink convolvulus Star of Bethlehem Blackberry nightshade Red pigweed Giant (black) pigweed Pink burr Square weed Thornapple Sowthistle Native jute Calopo Rattlepod Annual ground cherry Suppression only Bluetop ♣ Common sensitive plant	Qld, NSW, WA, NT only	2L to 4L #	Apply post-emergent to broadleaf weeds up to 8 leaf or 30cm diameter stage, vines to 1 m. Application should be made when weeds are growing actively and not suffering from moisture stress or waterlogging. Under cool conditions (<22° C) control may be slower and less effective. Can be applied over the top of cane. Some initial leaf yellowing and scorch may result but cane normally recovers well within 4 weeks. Caution should be exercised on new varieties of cane which may not have been tested. Use the lower rates on small weeds (less than 4 leaf) and the higher rates for larger weeds. Apply in a spray volume of 200-400 L/ha and ensure thorough coverage. Use the higher spray volumes where weed density is high or weeds are more advanced. ♣ Bluetop will only be suppressed. Use a minimum of 3 L/ha, on weeds less than 10 cm. Common sensitive plant - use a minimum of 3 L/ha
Sugar cane (plant and ratoon)	Green summer grass Crowsfoot grass Barnyard grass Guinea grass Summer grass Bluetop Centro Wild rose Bellvine Morning glory Red convolvulus Pink convolvulus Star of Bethlehem Giant sensitive plant Common sensitive plant Stinking passion vine Square weed Cinderella weed Giant (Black) pigweed Pink burr Thornapple Sowthistle Native jute Calopo Rattlepod Annual ground cherry Blackberry nightshade	Qld, NSW, WA, NT only	3L (+ 3L ametryn 500g/L) to 4L (+ 3.7L ametryn 500g/L) #	Use in a tank mix with ametryn (500 g/L) Apply between the early (1-4 leaf) and late (20 cm for grasses [Guinea grass and summer grass max. 15 cm] and 30 cm for broadleaf weeds) post-emergent stages of weeds. Use the lower rates on small weeds (less than 4 leaf) and for short residual control and the higher rates for large weeds and medium residual control. Where residual control is required the tank mix should be applied to moist soil and when rain or irrigation is expected within 10 days. Application should be made when weeds are growing actively and not suffering from moisture stress or waterlogging. Under cool conditions (<22°C) control may be slower and less effective. Apply in a spray volume of 200-400 L/ha and ensure thorough coverage. Use the higher volumes where weed density is high or weeds are more advanced. DO NOT apply this mixture by air. DO NOT apply over the top of cane. Use a directed spray to minimise contact with sugar cane foliage and shading of weeds. CAUTION: Contact with the foliage of ametryn sensitive cane varieties such as Q119, Q122 and H56-752 should be avoided as they are susceptible to phytotoxicity from the tank mix. Caution should be exercised on new varieties of cane, which may not have been tested.
	<i>Ipomoea plebeian</i> <i>Ipomoea purpurea</i> <i>Ipomoea hederifolia</i> <i>Ipomoea triloba</i> <i>Ipomoea quamoclit</i> <i>Solanum nigrum</i> <i>Portulaca oleracea</i> <i>Trianthema portulacastrum</i> <i>Urena lobata</i> <i>Spermacoce latifolia</i> <i>Datura stramonium</i> <i>Sonchus oleraceus</i> <i>Corchorus trilocularis</i> <i>Calapogonium mucunoides</i> <i>Crotalaria spp</i> <i>Physalis ixiocarpa</i> <i>Ageratum houstonianum</i> <i>Mimosa pudica</i> <i>Brachiaria miliiformis</i> <i>Eleusine indica</i> <i>Echinochloa colona</i> <i>Panicum maximum</i> var Hamil <i>Digitaria ciliaris</i> <i>Ageratum houstonianum</i> <i>Centrosema pubescens</i> <i>Cleome aculeata</i> <i>Ipomoea plebeia</i> <i>Ipomoea purpurea</i> <i>Ipomoea hederifolia</i> <i>Ipomoea triloba</i> <i>Ipomoea quamoclit</i> <i>Mimosa invisa</i> <i>Mimosa pudica</i> <i>Passiflora foetida</i> <i>Spermacoce latifolia</i> <i>Synedrella nodiflora</i> <i>Trianthema portulacastrum</i> <i>Urena lobata</i> <i>Datura stramonium</i> <i>Sonchus oleraceus</i> <i>Corchorus trilocularis</i> <i>Calapogonium mucunoides</i> <i>Crotalaria spp</i> <i>Physalis ixiocarpa</i> <i>Solanum nigrum</i>			

♣ * Suppression only at 1.5L/ha.
† Tolerant oat varieties are listed under the Crop Safety sub heading in the General Instructions section.
Add BS1000 120 mL/100L or Agral 200 mL/100L (sugar cane only)

Crop	Weeds Controlled		State	Rate/ha	Critical Comments
Wheat, barley and certain oat varieties †	Australian Crassula	<i>Crassula sieberana</i>	NSW, Vic, SA & WA only	1L	Spray actively growing weeds at the 2 to 6 leaf stage when cereals have 3 to 5 leaves on the main stem, i.e. Zadok scale code Z 13- 15. Ensure yellow burr weed plants are at the 2 to 6 leaf stage. If plants are older than this use the higher rate. DO NOT spray this weed after the 12 leaf stage.
	Cape Weed	<i>Arctotheca calendula</i>			
	Charlock	<i>Sinapis arvensis</i>			
	Fumitory-Red	<i>Fumaria densiflora</i>			
	Fumitory-White	<i>Fumaria parviflora</i>			
Indian Hedge Mustard	<i>Sisymbrium orientale</i>				
Mellilotus/Hexam scent	<i>Melilotus indica</i>				
Paterson's Curse	<i>Echium plantagineum</i>				
Rough Poppy	<i>Papaver hybridum</i>				
Rounded Chamomile	<i>Matricaria matricarioides</i>				
Sheepweed/Corn Gromwell	<i>Buglossoides arvensis</i>				
Turnip Weed	<i>Rapistrum rugosum</i>				
Wards weed	<i>Carrichtera annua</i>				
Wild Turnip	<i>Brassica tournefortii</i>				
Yellow Burr Weed	<i>Amsinckia spp.</i>				
Fat-hen	<i>Chenopodium album</i>	SA &			
Scented Woolly Sage	<i>Salvia lanigera</i>	WA only			
Ball Mustard *♣	<i>Neslia paniculata</i>	NSW, Vic, SA & WA only	1.5 L	Spray actively growing weeds at the 2 to 6 leaf stage when cereals have 3 to 5 leaves on the main stem, i.e. Zadok scale code Z 13- 15. For Three-cornered Jack apply at the 1 to 4 true leaf stage. Control of Ivyleaf Speedwell may at times be variable.	
Deadnettle	<i>Lamium amplexicaule</i>				
Mallee Catchfly	<i>Silene apetala</i>				
Pimpernel	<i>Anagallis arvensis</i>				
Speedwell-Ivyleaf	<i>Veronica hederifolia</i>				
Three-cornered Jack/ Spiny Emex/ Doublegee	<i>Emex australis</i>				
Toadrush	<i>Juncus bufonis</i>				
Wireweed	<i>Polygonum aviculare</i>				
Saffron Thistle	<i>Carthamus lanatus</i>	SA & WA only			
Stemless Thistle	<i>Onopordum acaulon</i>				

Crop	Weeds Controlled		State	Rate /ha	Critical Comments
Pastures Use on Sub. clover (min 1 trifoliolate leaves) based pasture with either/or Ryegrass, Phalaris, Cocksfoot (grasses min. 2 leaves)	Paterson's Curse (cotyledon to 2 leaf)	<i>Echium plantagineum</i>	NSW, Vic, SA, and WA only	300 mL to 500 mL	Apply soon after the break to the season when the weeds are up to the 2 leaf stage. Apply by boom spray or aircraft. Use the highest rate when the majority of plants are at the 2 leaf stage. Some reduction in clover growth may occur for a period after application. Plants usually recover and grow normally. Pasture should be growing well with adequate fertiliser and mite/insect control to help reduce the establishment of later germinating seedlings. Larger weeds present at spraying may not be controlled. DO NOT use on medics. DO NOT apply in less than 50 L/ha water. DO NOT use on seed crops. DO NOT spray after mid-August as damage may occur.
Pastures Use on mixtures of either/or Sub. Clover (min. 3 trifoliolate leaves), Balansa clover (established plants), White clover (min. 5 trifoliolate leaves) with either/or Ryegrass, Phalaris, Cocksfoot (grasses min. 3 leaves)	Capeweed (seedlings up to 20 cm diam.) Long Storksbill (seedlings up to 8 cm diam.) Paterson's Curse (seedlings up to 20 cm diam.) Toadrush (1-5 leaves) Crassula (1-5 cm diam.)	<i>Arctotheca calendula</i> <i>Erodium botrys</i> <i>Echium plantagineum</i> <i>Juncus bufonius</i> <i>Crassula spp</i>	NSW, Vic, SA & WA only	1 L to 1.5 L	Apply by boom spray or aircraft. Use the highest rate when weeds are greater than 7 cm in diameter. Where a selective grass herbicide is to be used, apply Paddo M SC prior to the application of the selective grass herbicide. Some reduction in clover growth may occur for a period after application. Plants usually recover and grow normally. DO NOT use on medics. DO NOT apply in less than 50 L/ha water. DO NOT use on seed crops. DO NOT spray after mid-August as damage may occur.

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION