

**KENSO AGCARE**

# **PREVAKEN**

## **200 SC**

### **INSECTICIDE**

**ACTIVE CONSTITUENTS:**  
**200 g/L CHLORANTRANILIPROLE**

|              |           |                    |
|--------------|-----------|--------------------|
| <b>GROUP</b> | <b>28</b> | <b>INSECTICIDE</b> |
|--------------|-----------|--------------------|

For the control of Lepidopteran species of insect pests  
in certain vegetables, as per the Directions for Use.

Kenso Corporation (M) Sdn Bhd  
Level 1, 98 Commercial Road,  
Teneriffe QLD 4005  
Phone (07) 3216 1188  
[www.kenso.com.au](http://www.kenso.com.au)



**IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USE**

**CONTENTS: 5 Litres    APVMA Approval No.: 93440/ 139159**

# KENSO AGCARE PREVAKEN 200 SC INSECTICIDE

## STORAGE AND DISPOSAL

Store in the closed, original container in cool, well ventilated area. Do not store for prolonged periods in direct sunlight. Triple-rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. Do not burn empty containers or product. Do not re-use empty container for any other purpose.

## FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

## SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet, which can be obtained from [www.kenso.com.au](http://www.kenso.com.au).

## CONDITIONS OF SALE

Kenso Corporation (M) Sdn. Bhd. will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Kenso Corporation (M) Sdn. Bhd. is limited to the replacement of the goods or (at the option of Kenso Corporation (M) Sdn. Bhd.) the refund of the price paid and where possible sufficient part of the goods to enable proper examination being returned to Kenso Corporation (M) Sdn. Bhd. within thirty days of delivery.

In a Transport Emergency  
Dial **000**  
Police or Fire Brigade



Batch No.:  
Date of Manufacture:

**KENSO**  
agcare

# PREVAKEN 200 SC

**Insecticide**

**ACTIVE CONSTITUENT:**  
**200 g/L CHLORANTRANILIPROLE**

**GROUP 28 INSECTICIDE**

For the control of Lepidopteran species of insect pests  
in certain vegetables, as per the Directions for Use.

**IMPORTANT: READ THE LEAFLET BEFORE USE**

**APVMA Approval No.: 93440/139159**



**Kenso Corporation (M) Sdn. Bhd.**  
Level 1, 98 Commercial Road,  
Teneriffe QLD 4005  
Phone (07) 3216 1188

## RESTRAINTS

DO NOT apply if rainfall is expected within 2 hours of application.  
DO NOT use on container or hydroponic grown crops.

## SPRAY DRIFT RESTRAINTS

Specific definitions for terms used in this section of the label can be found  
at [www.apvma.gov.au/spraydrift](http://www.apvma.gov.au/spraydrift)

DO NOT allow bystanders to come into contact with the spray cloud.  
DO NOT apply in a manner that may cause an unacceptable impact to  
native vegetation, agricultural crops, landscaped gardens and  
aquaculture production, or cause contamination of plant or livestock  
commodities, outside the application site from spray drift. The buffer  
zones in the relevant buffer zone table/s below provide guidance but may  
not be sufficient in all situations. Wherever possible, correctly use  
application equipment designed to reduce spray drift and apply when the  
wind direction is away from these sensitive areas.

DO NOT apply unless the wind speed is between 3 and 20 kilometres per  
hour at the application site during the time of application.

DO NOT apply if there are hazardous surface temperature inversion  
conditions present at the application site during the time of application.  
Surface temperature inversion conditions exist most evenings one to two  
hours before sunset and persist until one to two hours after sunrise.

DO NOT apply by a boom sprayer unless the following requirements are  
met:

- spray droplets not smaller than a MEDIUM spray droplet size category
- minimum distances between the application site and downwind sensitive  
areas (see 'Mandatory no-spray zones' section) are observed.

DO NOT apply by aircraft unless the following requirements are met:

- spray droplets not smaller than a MEDIUM spray droplet size category
- for release heights 25% of wingspan or 25% of rotor diameter or lower  
above the target canopy, minimum distances between the application  
site and downwind sensitive areas (see 'Mandatory no-spray zones'  
section) are observed.

## MANDATORY NO-SPRAY ZONES

DO NOT apply if there are aquatic or wetland areas including aquacultural  
ponds downwind from the application area and within the mandatory  
no-spray zones shown in Table 1 below.

## RESISTANCE WARNING

**GROUP 28 INSECTICIDE**

For insecticide resistance management Kenso Agcare Prevaken 200 SC  
Insecticide is a Group 28 insecticide. Some naturally occurring insect  
biotypes resistant to Kenso Agcare Prevaken 200 SC Insecticide and  
other Group 28 insecticides may exist through normal genetic variability in  
any insect population. The resistant individuals can eventually dominate  
the insect population if Kenso Agcare Prevaken 200 SC Insecticide and  
other Group 28 insecticides are used repeatedly. The effectiveness of  
Kenso Agcare Prevaken 200 SC Insecticide on resistant individuals could  
be significantly reduced. Since the occurrence of resistant individuals is  
difficult to detect prior to use Kenso Agcare Pty Ltd accepts no liability for  
any losses that may result from the failure of Kenso Agcare Prevaken 200  
SC Insecticide to control resistant insects. Kenso Agcare Prevaken 200  
SC Insecticide may be subject to specific resistance management  
strategies. For further information contact our local supplier, Kenso  
Agcare Pty Ltd representative or local agricultural department agronomist.

## INSECTICIDE RESISTANCE MANAGEMENT (IRM)

Kenso Agcare Prevaken 200 SC Insecticide contains chlorantraniliprole, a  
Group 28 Insecticide. Unless directed otherwise in the specific crop/insect  
sections of the label, the following practices are recommended to prevent  
or delay the development of insecticide resistance to Kenso Agcare  
Prevaken 200 SC Insecticide and to Group 28 insecticides:

- Apply Kenso Agcare Prevaken 200 SC Insecticide or other Group 28  
insecticides using a "window" approach to avoid exposure of  
consecutive insect pest generations to the same mode of action. Multiple  
successive applications of Kenso Agcare Prevaken 200 SC Insecticide  
or other Group 28 insecticides are acceptable if they are used to treat a  
single insect generation.
- Following a "window" of Kenso Agcare Prevaken 200 SC Insecticide or  
other Group 28 insecticides, rotate to a "window" of applications of  
effective insecticides with a different mode of action.
- The total exposure period of all "Group 28-active windows" applied  
throughout the crop cycle (from seedling to harvest) should not exceed  
50% of the crop cycle.
- Incorporate IPM techniques into the overall pest management program.
- Monitor insect populations for loss of field efficacy.

For additional information on insect resistance, modes of action and  
monitoring visit the Insecticide Resistance Action Committee (IRAC) on  
the web at <http://www.irac-online.org>

Table 1 – No-Spray Zones for Protection of the Aquatic Environment

| FOR AERIAL APPLICATION                  |                                  |                                  |
|-----------------------------------------|----------------------------------|----------------------------------|
| Wind Speed Range at Time of Application | Downwind Mandatory No-Spray Zone |                                  |
|                                         | Fixed-Wing                       | Helicopter                       |
|                                         | 200 metres                       | 200 metres                       |
|                                         |                                  |                                  |
|                                         |                                  |                                  |
| from 3 to 20 kilometres per hour        |                                  |                                  |
| from 8 to 14 kilometres per hour        |                                  |                                  |
| from 14 to 20 kilometres per hour       |                                  |                                  |
| FOR GROUND APPLICATION                  |                                  |                                  |
| Usage                                   | Applications per season          | Downwind Mandatory No-Spray Zone |
| Vegetables                              | 3                                | 15 metres                        |

## GENERAL INSTRUCTIONS

Kenso Agcare Prevaken 200 SC Insecticide has been specifically  
designed for use in Integrated Pest Management (IPM) schemes. Kenso  
Agcare Prevaken 200 SC Insecticide is an anthranilic diamide insecticide  
in the form of a suspension concentrate. Kenso Agcare Prevaken 200 SC  
Insecticide is particularly active on Lepidopteran insect pests, primarily as  
a larvicide. Before application monitor insect populations to determine  
whether or not there is a need for application Kenso Agcare Prevaken 200  
SC Insecticide based on locally determined economic thresholds. More  
than one treatment of Kenso Agcare Prevaken 200 SC Insecticide may be  
required to control a population of pests.

## MIXING

Fill spray tank to ¼ to ½ full of water. Measure the amount of Prevaken  
200 SC Insecticide required for the area to be sprayed. Add Kenso Agcare  
Prevaken 200 SC Insecticide directly to the spray tank with the agitation  
engaged. Mix thoroughly to disperse the insecticide. Once dispersed, the  
material must be kept in suspension at all times by continuous agitation.  
Use mechanical or hydraulic means, DO NOT use air agitation, premix or  
slurry. If spray solution is left standing, ensure thorough re-agitation of the  
spray mix until fully resuspended. DO NOT allow spray mix to sit  
overnight, as resuspension may be difficult.

## FIRST AID INSTRUCTIONS

If poisoning occurs, contact a doctor or Poisons Information Centre.  
Phone Australia 13 11 26.

## SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet, which can be  
obtained from [www.kenso.com.au](http://www.kenso.com.au)

## CONDITIONS OF SALE

Kenso Corporation (M) Sdn. Bhd. will not accept any responsibility  
whatsoever and howsoever arising and whether for consequential loss or  
otherwise in connection with the supply of these goods other than  
responsibility for the merchantable quality of the goods and such  
responsibilities mandatorily imposed by Statutes applicable to the sale or  
supply of these goods. To the extent allowed by such Statutes the liability  
of Kenso Corporation (M) Sdn. Bhd. is limited to the replacement of the  
goods or (at the option of Kenso Corporation (M) Sdn. Bhd.) the refund of  
the price paid and where possible sufficient part of the goods to enable  
proper examination being returned to Kenso Corporation (M) Sdn. Bhd.  
within thirty days of delivery.

In a Transport Emergency  
Dial **000**  
Police of Fire Brigade

\* Not a registered trademark of Kenso



**Kenso Corporation (M) Sdn. Bhd.**  
Level 1, 98 Commercial Road,  
Teneriffe QLD 4005  
Phone (07) 3216 1188

**APVMA Approval No.: 93440/139159**

## SURFACTANT/WETTING AGENT

For Brassica vegetable, Brassica leafy vegetable, Stalk & Stem  
vegetables, Leafy vegetables, Lettuce, Fruiting vegetables (Cucurbits),  
Legume vegetables, Sweet corn use a non-ionic surfactant/ wetting agent  
at 15 g active/100 L, (e.g. Agral\* 600 @ 25 mL/100 L). DO NOT use  
BS1000\* or Activator\* as it may cause crop phytotoxicity.

DO NOT add a non-ionic surfactant/wetting agent if:

- mixing with another product which already contains a surfactant and/or  
the product label advises not to add a surfactant.
- mixing with a liquid fertiliser.

## APPLICATION

Application equipment should be calibrated to apply at least sixty (60)  
droplets per cm<sup>2</sup> of target foliage. Droplet VMD should be of medium  
spray quality.

## GROUND APPLICATION

Use a boom sprayer fitted with high flow rate nozzles to apply the highest  
practical spray volume. Nozzles with higher rated flows produce larger  
droplets. Use the lower spray pressures recommended for the nozzle.  
Higher pressure reduces droplet size, DOES NOT improve canopy  
penetration and may increase drift potential. WHEN HIGHER FLOW  
RATES ARE NEEDED, USE A HIGHER-CAPACITY NOZZLE INSTEAD  
OF INCREASING PRESSURE. Use a nozzle type that is designed for the  
intended application. With most nozzle types, narrower spray angles  
produce larger droplets. Consider using low-drift nozzles. When applying  
Kenso Agcare Prevaken 200 SC Insecticide by ground application, keep  
the boom low to avoid spray drift.

## AERIAL APPLICATION (BY FIXED WING AIRCRAFT OR HELICOPTER)

Kenso Agcare Prevaken 200 SC Insecticide must only be applied with  
aircraft fitted with accurately calibrated equipment. Apply a minimum total  
spray volume of 40 L/ha with nozzles (e.g. Micronaire rotary atomisers,  
CP nozzles or conventional hydraulic nozzles) set to medium spray  
quality. A spray drift minimisation strategy should be employed at all times  
when applying this product. DO NOT apply Kenso Agcare Prevaken 200  
SC Insecticide using Ultra Low Volume (ULV) methods.

DIRECTIONS FOR USE

For use in all States where appropriate for the crop and/or insect pest.

| CROP                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          | PEST                                                                                                                                                                                                                                                                                                                                                             | RATE/HA                                                                                                                   | WHP          | CRITICAL COMMENTS                                                                                                                                                                                    |  |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------|--------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--|
| <b>CRITICAL COMMENTS - ALL CROPS</b><br>Regularly scout crops to monitor for eggs and larvae. Target sprays against eggs and newly hatched larvae before they become entrenched. Apply as egg and larvae reach threshold numbers.<br>A maximum of three (3) applications are to be applied to any one crop. No more than two (2) consecutive sprays per crop,with a minimum spray interval of 7 days (unless stated otherwise). Further treatments should be made with alternative mode of action insecticides.<br>Use enough water to ensure thorough coverage of the crop. Adjust water volumes to crop stage (200 – 1000 L/ha).Refer to Surfactant/Wetting agent section.<br>Use in accordance with AIRAC Insecticide Resistance Management Strategy guidelines. As part of an Insecticide Resistance Management programme for Cotton bollworm, it is important to plough crops immediately after harvest. |                                                                                                                                                                                                                                                                                                                                                                  |                                                                                                                           |              |                                                                                                                                                                                                      |  |
| Brassica vegetables including:<br>Broccoli<br>Brussels sprout<br>Cabbage<br>Cauliflower                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       | Cabbage-centre grub ( <i>Hellula hydralis</i> )<br>Cabbage cluster caterpillar ( <i>Crociodolomia pavonana</i> )                                                                                                                                                                                                                                                 | 100 mL<br>+<br>15 gai/100L of non-ionic surfactant                                                                        | 7 days       |                                                                                                                                                                                                      |  |
| Brassica leafy vegetables including:<br>Buk choy<br>Chinese broccoli (Gailum/ Gai lan/Kai lan),<br>Chinese cabbage (Pet sai/ Wombok/ Haksukai)<br>Choy sum<br>Gai choy/Am soy<br>Kai choy<br>Kale<br>Mibuna<br>Leafy mustard including Indian mustard and Mustard spinach (Komatsuma)<br>Pak choy Tat soy                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     | Cabbage leafminer ( <i>Liriomyza brassicae</i> )<br>Cabbage white butterfly ( <i>Pieris rapae</i> )<br>Cluster caterpillar ( <i>Spodoptera litura</i> )<br>Cotton bollworm ( <i>Helicoverpa armigera</i> )<br>Diamondback moth ( <i>Plutella xylostella</i> )Native budworm ( <i>Helicoverpa punctigera</i> ) Soybean looper ( <i>Thysanoplusia orichalcea</i> ) |                                                                                                                           | 3 days       |                                                                                                                                                                                                      |  |
| Stalk & Stem vegetables, including: Celery Rhubarb                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            | Cotton bollworm ( <i>Helicoverpa armigera</i> )<br>Native budworm ( <i>Helicoverpa punctigera</i> )                                                                                                                                                                                                                                                              | 100 mL<br>+<br>15 gai/100 L of non-ionic surfactant                                                                       |              |                                                                                                                                                                                                      |  |
| Leafy vegetables (excluding lettuce),including:<br>Cress<br>Endive<br>Silverbeet<br>Spinach                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |                                                                                                                                                                                                                                                                                                                                                                  |                                                                                                                           |              |                                                                                                                                                                                                      |  |
| Lettuce (leaf and closed head varieties)                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |                                                                                                                                                                                                                                                                                                                                                                  | 150 mL<br>+<br>15 gai/100 L of non-ionic surfactant                                                                       |              |                                                                                                                                                                                                      |  |
| Fruiting vegetables (excluding Cucurbits),including: Capsicum<br>Egg plant Peppers Tomato (trellis and field)                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                 | Cotton bollworm ( <i>Helicoverpa armigera</i> )<br>Native budworm ( <i>Helicoverpa punctigera</i> )<br>Tomato leaf miner ( <i>Phthorimaea operculella</i> )                                                                                                                                                                                                      | 100 mL<br>or<br>10 mL/100 L (dilute)                                                                                      | 3 days       | Ensure spray timing coincides with egg laying/hatching. The pest hatches from the egg and burrows directly into fruit. Larvae entrenched in the fruit at the time of sprayingwill not be controlled. |  |
|                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                               | Eggfruit caterpillar ( <i>Sceliodes cordalis</i> )                                                                                                                                                                                                                                                                                                               |                                                                                                                           |              |                                                                                                                                                                                                      |  |
| Fruiting vegetables (Cucurbits), including: Cucumbers Melons<br>Pumpkin Squash Zucchini                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       | Cotton bollworm ( <i>Helicoverpa armigera</i> )<br>Native budworm ( <i>Helicoverpa punctigera</i> )<br>Cucumber moth ( <i>Diaphania indica</i> )                                                                                                                                                                                                                 | 100 mL<br>+<br>15 gai/100 Lof non-ionic surfactant or<br>10 mL/100 L (dilute)<br>+<br>15 gai/100 Lof non-ionic surfactant | 1 day        | Apply with a minimum spray interval of 5 days.                                                                                                                                                       |  |
| Legume vegetables including:<br>Green beans Green peas<br>Processing peasSnow peas<br>Sugar snap peas                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         | Cotton bollworm ( <i>Helicoverpa armigera</i> )<br>Native budworm ( <i>Helicoverpa punctigera</i> )                                                                                                                                                                                                                                                              | 100 mL<br>+<br>15 gai/100 Lof non-ionic surfactant)                                                                       | 1 day        |                                                                                                                                                                                                      |  |
| Potatoes                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      | Cotton bollworm ( <i>Helicoverpa armigera</i> )<br>Native budworm ( <i>Helicoverpa punctigera</i> )<br>Potato moth ( <i>Phthorimaea operculella</i> )                                                                                                                                                                                                            | 100 mL                                                                                                                    | Not required | Only target foliar infestations of Potato moth. Moth larvae in the soil or within stems will notbe controlled. Apply with a spray interval of 10 - 14 days.                                          |  |
| Sweet corn                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    | Cotton bollworm ( <i>Helicoverpa armigera</i> )                                                                                                                                                                                                                                                                                                                  | 100 mL<br>+<br>15 gai/100 Lof non-ionic surfactant                                                                        | 7 days       | Ensure spray timing coincides with egg laying/hatching. Larvae entrenched in cobs at the time ofspraying will not be controlled.                                                                     |  |

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WITHHOLDING PERIODS

HARVEST

POTATOES: WITHHOLDING PERIOD NOT REQUIRED WHEN USED AS DIRECTED.  
FRUITING VEGETABLES (CUCURBITS INCLUDING CUCUMBERS, MELONS, PUMPKIN, SQUASH, ZUCCHINI), LEGUME VEGETABLES: DO NOT HARVEST FOR 1 DAY AFTER APPLICATION  
BRASSICA LEAFY VEGETABLES, FRUITING VEGETABLES (excluding CUCURBITS), LEAFY VEGETABLES (including LETTUCE), STALK & STEM VEGETABLES: DO NOT HARVEST FOR 3 DAYS AFTER APPLICATION  
BRASSICA VEGETABLES (INCLUDING BROCCOLI, BRUSSELS SPROUT, CABBAGE, CAULIFLOWER), SWEET CORN: DO NOT HARVEST FOR 7 DAYS AFTER APPLICATION.

GRAZING

LEGUME VEGETABLES: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 1 DAY AFTER APPLICATION.  
SWEET CORN: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION.  
OTHER CROPS: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION.

TRADE ADVICE EXPORT STATEMENT

Import tolerances for produce treated with Kenso Agcare Prevaken 200 SC Insecticide may be pending in some countries. Consult with your exporter or Kenso Agcare Pty Ltd before applying Kenso Agcare Prevaken 200 SC Insecticide to export crops.

COMPATIBILITY

Since formulations may be changed and new ones introduced, it is recommended that users premix a small quantity of the desired tank mix and observe possible adverse changes (settling out, flocculation etc). Avoid complex tank mixtures of several products or very concentrated spray mixtures.  
The mixing sequence recommended is: water soluble bags, dry flowable or water dispersible granules, wettable powders, water based suspension concentrates (Kenso Agcare Prevaken 200 SC Insecticide insecticide), water soluble concentrates, oil based suspension concentrates, emulsifiable concentrates, adjuvants and surfactants, soluble fertilisers.

SPRAY EQUIPMENT CLEANOUT

Prior to application, start with clean, well-maintained application equipment. Immediately following application, thoroughly clean all spray equipment to reduce the risk of forming hardened deposits which might become difficult to remove. Drain spray equipment.  
Thoroughly rinse sprayer and flush hoses, boom, and nozzles with clean water. Clean all other associated application equipment. Take all necessary safety precautions when cleaning equipment. DO NOT clean near wells, water sources or desirable vegetation. Dispose of waste rinse water in accordance with local regulations.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Very toxic to aquatic life. DO NOT contaminate wetlands or watercourses with this product or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in cool, well ventilated area. Do not store for prolonged periods in direct sunlight. Triple-rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.  
If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. Do not burn empty containers or product. Do not re-use empty container for any other purpose.