

# CAUTION

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

*KENSO AGCARE*

# AVIOR 250 SC FUNGICIDE

ACTIVE CONSTITUENT: 250 g/L AZOXYSTROBIN

**GROUP 11 FUNGICIDE**

For the control of various diseases of Grapes, Potatoes, Tomatoes, Cucurbits, Avocados, Mangoes, Passionfruit, Poppies, Pyrethrum, Olives, Ribberies, Anise Myrtle, Lemon Myrtle, Carrot, Nursery Stock, Ornamentals, Cut Flower/Foliage and *Rubus* crop as per the Directions for Use.



Kenso Corporation (M) Sdn Bhd  
Level 1, 98 Commercial Road,  
Teneriffe QLD 4005  
Phone (07) 3216 1188  
[www.kenso.com.au](http://www.kenso.com.au)

**IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USE**

**CONTENTS: 20 Litres APVMA Approval No.: 70285/ 118936**

# KENSO AGCARE AVIOR 250 SC FUNGICIDE

## STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. DO NOT store in direct sunlight. Triple rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush, or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

## SAFETY DIRECTIONS

Will irritate the eyes. Avoid contact with eyes. Wash hands after use. When opening the container and preparing spray and using the prepared spray wear cotton overalls buttoned to the neck and wrist, or equivalent clothing. After each day's use, wash contaminated clothing.

## FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

## SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet, which can be obtained from [www.kenso.com.au](http://www.kenso.com.au).

## CONDITIONS OF SALE

Kenso Corporation (M) Sdn. Bhd. will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply of these goods

other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Kenso Corporation (M) Sdn. Bhd. is limited to the replacement of the goods or (at the option of Kenso Corporation (M) Sdn. Bhd.) the refund of the price paid and where possible sufficient part of the goods to enable proper examination being returned to Kenso Corporation (M) Sdn. Bhd. within thirty days of delivery.

In a Transport Emergency  
Dial **000** Police or Fire Brigade



9 328666 003250



Additional statements required by Globally Harmonised System of Classification and Labelling of Chemicals (GHS) and Safe Work Australia: **MAY CAUSE AN ALLERGIC SKIN REACTION. CAUSES EYE IRRITATION.** *Precautionary:* Avoid breathing dust/ fume/ gas/ mist/ vapours/ spray. Wash contacted areas thoroughly after handling. Contaminated work clothing should not be allowed out of the workplace. Wear protective gloves, protective clothing and eye or face protection. IF ON SKIN: Wash with plenty of soap and water. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. Specific treatment (see FIRST AID) on this label. If skin irritation occurs: Get medical advice/attention. If eye irritation persists: Get medical advice/attention. If skin irritation or rash occurs: Get medical advice/attention. Wash contaminated clothing before reuse.

Batch No.:

Date of Manufacture:

**CAUTION**  
KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

**KENSO**  
*agcare*  
**AVIOR**  
**250 SC**  
Fungicide

ACTIVE CONSTITUENT: 250 g/L AZOXYSTROBIN

GROUP **11** FUNGICIDE

For the control of various diseases of Grapes, Potatoes, Tomatoes, Cucurbits, Avocados, Mangoes, Passionfruit, Poppies, Pyrethrum, Olives, Ribberries, Anise Myrtle, Lemon Myrtle, Carrot, Nursery Stock, Ornamentals, Cut Flower/Foliage and *Rubus* crop as per the Directions for Use.

READ THIS LEAFLET BEFORE USE

APVMA Approval No.: 70285/118936



**Kenso Corporation (M) Sdn. Bhd.**  
Level 1, Commercial Road,  
Teneriffe QLD 4005  
Phone (07) 3216 1188

**PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS**

**WARNING:** Kenso Agcare Avior 250 SC Fungicide is extremely phytotoxic to certain apple varieties.

**AVOID SPRAY DRIFT.** Extreme care must be used to prevent injury to apple trees.

**DO NOT** spray Kenso Agcare Avior 250 SC Fungicide where spray drift may reach apple trees.

**DO NOT** spray when conditions favour drift beyond the area intended for application. Conditions that may contribute to drift include thermal inversions, excessive wind speed, certain sprayer nozzle/pressure combinations, small spray droplet size etc.

**DO NOT** use spray equipment that has been previously used to apply Kenso Agcare Avior 250 SC Fungicide to spray apple trees. Even trace amounts can cause unacceptable phytotoxicity.

**PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT**

HIGHLY TOXIC TO AQUATIC LIFE. DO NOT contaminate dams, waterways or drains with the chemical or used containers.

DO NOT apply under weather conditions or from spraying equipment which could be expected to cause spray drift on adjacent areas, particularly wetlands, waterbodies or watercourses.

**STORAGE AND DISPOSAL**

Store in the closed, original container in a cool, well-ventilated area. DO NOT store in direct sunlight. Triple rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush, or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

**SAFETY DIRECTIONS**

Will irritate the eyes. Avoid contact with eyes. Wash hands after use. When opening the container and preparing spray and using the prepared spray wear cotton overalls buttoned to the neck and wrist, or equivalent clothing. After each day's use, wash contaminated clothing.

**FIRST AID**

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

**SAFETY DATA SHEET**

Additional information is listed in the Safety Data Sheet, which can be obtained from [www.kenso.com.au](http://www.kenso.com.au).

**CONDITIONS OF SALE**

Kenso Corporation (M) Sdn. Bhd. will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Kenso Corporation (M) Sdn. Bhd. is limited to the replacement of the goods or (at the option of Kenso Corporation (M) Sdn. Bhd.) the refund of the price paid and where possible sufficient part of the goods to enable proper examination being returned to Kenso Corporation (M) Sdn. Bhd. within thirty days of delivery.

In a Transport Emergency  
Dial **000**  
Police of Fire Brigade



**Kenso Corporation (M) Sdn. Bhd.**  
Level 1, 98 Commercial Road,  
Teneriffe QLD 4005  
Phone (07) 3216 1188

APVMA Approval No.: 70285/118936

**DIRECTIONS FOR USE**

**Restraint:** DO NOT apply by air, except on potatoes.

TREE AND VINE CROPS				
Rate				Critical comments
In the following table, Tree and Vine Crops, all rates given are for dilute spraying. For concentrate spraying, refer to the Application section.				For all uses in the table Tree and Vine Crops: apply by dilute or concentrate spraying equipment. Apply the same total amount of product to the target crop whether applying this product by dilute or concentrate spraying methods. When applying Kenso Agcare Avior 250 SC Fungicide through low volume application equipment, DO NOT use a concentrate factor greater than 4X. In these cases adequate coverage of all plant surfaces is still required to achieve control of diseases.
Crop	Disease	Rate /100L	WHP	
Avocados	Stem End Rot, Anthracnose	80ml	7 days	Apply Kenso Agcare Avior 250 SC Fungicide in a preventative fungicide program containing fungicides from a different chemical group. For best results commence the disease control program with an approved fungicide from an alternative chemical group, then apply 1 application of Kenso Agcare Avior 250 SC Fungicide during early fruit set. Follow with applications of an approved fungicide from a different chemical group. Apply 2 final applications of Kenso Agcare Avior 250 SC Fungicide at 14 to 28 day intervals with the final spray applied 7 days prior to harvest. Ensure thorough spray coverage.
Grapes table, wine, dried	Powdery Mildew ( <i>Uncinular necator</i> ), Downy Mildew ( <i>Plasmopara viticola</i> ), Botrytis Bunch Rot † ( <i>Botrytis cinerea</i> )	75ml to 100ml	14 days	<p><b>Application Method and Rate</b> Apply in a sufficient volume of water to achieve thorough coverage of all foliage and fruit. The volume of water required to achieve this will depend on the stage of vine growth and vigour. Ensure thorough coverage. Adjust spray nozzles to direct spray droplets to the canopy present. Apply the higher rate of application in the following circumstances:</p> <ol style="list-style-type: none"> <li>Where humid conditions favour Powdery Mildew infection, particularly on susceptible varieties.</li> <li>At the start of the season when there has been a heavy carry over of Powdery Mildew infection (flag shoots are present).</li> </ol> <p><b>Spray Timing and Interval</b> Apply 2 or 3 consecutive applications at 10 to 16 day intervals at any time between early shoot growth and 14 days before harvest. Use the recommended shorter interval during periods when climatic conditions are favourable for disease infection.</p> <p>† <b>Botrytis Bunch Rot</b> Kenso Agcare Avior 250 SC Fungicide must not be used alone for Botrytis control at critical times such as 80 to 100% capfall and preharvest. It must be tank mixed with or substituted by a specific botryticide at these critical times. When Kenso Agcare Avior 250 SC Fungicide is used in a seasonal spray programme it will provide control of Botrytis additional to that of specific botryticides such as Kenso Agcare Castor Fungicide.</p>
Mangoes	Stem End Rot, Anthracnose	80ml	3 days	For best results apply 1 to 2 applications of Kenso Agcare Avior 250 SC Fungicide at flowering and early fruit set, at no less than 14 day intervals. Follow with applications of an approved fungicide from a different chemical group. Further applications of Kenso Agcare Avior 250 SC Fungicide may be applied at 21 days and 3 to 7 days prior to harvest. Ensure thorough spray coverage.
Passion Fruit	Alternaria, Cladosporium	80ml	1 day	Apply Kenso Agcare Avior 250 SC Fungicide in a preventative fungicide program containing fungicides from a different chemical group. For best results apply 2 to 3 applications of Kenso Agcare Avior 250 SC Fungicide at 14 day intervals over flowering. Follow with applications of an approved fungicide from a different chemical group. Apply a further 1 to 2 applications of Kenso Agcare Avior 250 SC Fungicide finishing 1 day prior to harvest. Ensure thorough spray coverage.
Olives	Anthracnose	80 mL /100L	21 days	Apply sufficient quantity of the mixed chemical to wet all surfaces of leaves and fruit. Apply by air blast or boomspray. The use of an appropriate wetting agent is recommended to improve the spread of the chemical over the leaves and fruit. DO NOT apply more than 2 applications per season. Allow a minimum of 21 days between consecutive applications. Apply the treatment, preferably before the disease infects the trees. Fungicides are best applied prior to the onset of conditions conducive to this disease (warm, humid rainy weather. This will depend upon whether the olive grove is in a susceptible area (e.g. summer rains), and the season (unseasonal humid and moist conditions). Spraying prior to flowering is a good guide, and again just after fruit set. Protect the remaining periods with other approved fungicides if required.

OTHER CROPS				
Crop	Disease	Rate /100L	WHP	Critical Comments
Carrots	Powdery Mildew ( <i>Erysiphe heraclei</i> )	1 L/ha	21 days	Apply in a preventative program commencing before disease infection occurs, particularly during weather conditions that favour disease development, or (at the latest) when first signs of the disease are observed. Apply a maximum of three (3) foliar applications in total per crop per season, with a maximum two (2) consecutive applications. Apply foliar spray at 10 - 14 day interval. Use shorter interval when weather conditions are highly conducive to disease infection. Apply in sufficient water volume to achieve thorough coverage of all foliage using ground boom spray equipment or equivalent only as a foliar spray. Good coverage of foliage is essential. Apply between 500 - 1,500 L of spray mix to adequately treat a hectare, depending on crop stage and foliage density. Use a higher volume in dense or well grown crops. If treating for Black Rot, irrigate thoroughly (at least 20,000 L/ha) to water the product into the soil.
	Sclerotinia rot/ White mould ( <i>Sclerotinia sclerotiorum</i> )	400 mL/ha		
	Black Rot ( <i>Alternaria radicina</i> ) SUPPRESSION ONLY	400 mL/ha		
Cucurbits	Powdery Mildew ( <i>Sphaerotheca fuliginea</i> ), Downy Mildew ( <i>Pseudoperonospora cubensis</i> )	80 to 120 ml	1 day	Apply the higher rate when climatic conditions favour Powdery or Downy Mildew infection and in crops with large canopies. <b>Application</b> Apply in a sufficient volume of water to achieve thorough coverage of all foliage. The volume of water required to achieve this will depend on the stage of growth of the cucurbits. For dilute spraying (g/100 L), an application volume of 300 L/ha is suggested where sprays are banded in the early part of the season, increasing to 1000 L/ha as a broadcast spray in a vigorous crop at full canopy. <b>Spray Interval</b> Consecutive applications should be applied at 7 to 14 day intervals, commencing soon after transplanting and continuing up to fruit maturity. Use the recommended shorter application interval in the following circumstances: <ol style="list-style-type: none"> <li>Under humid weather conditions which are favourable for Powdery Mildew, Downy Mildew or Gummy Stem Blight infection.</li> <li>When there is rapid vegetative growth during the early part of the crop cycle.</li> </ol>
Nursery stock and ornamentals: Including (non-food) - seedlings, plugs, potted colour, trees, shrubs, foliage plants, palms, grasses, fruit trees (non-bearing*) and ornamentals	Downy mildew ( <i>Peronospora</i> spp., <i>Pseudoperonospora</i> spp., <i>Bremia lactucae</i> ) Grey mould ( <i>Botrytis</i> spp.) Leaf spots ( <i>Colletotrichum</i> spp. & <i>Alternaria</i> spp.) Powdery mildew ( <i>Erysiphe</i> spp., <i>Leveillula</i> spp., <i>Microsphaera</i> spp., <i>Oidium</i> spp. & <i>Sphaerotheca</i> spp.) Rusts ( <i>Puccinia</i> spp., <i>Phragmidium</i> spp., <i>Uromyces</i> spp.)	80 to 120 mL	-	Apply in sufficient volume to ensure adequate coverage of all plant surfaces. DO NOT use Kenso Agcare Avior 250 SC Fungicide curatively. Apply a maximum of 2 consecutive applications of Kenso Agcare Avior 250 SC Fungicide with a minimum re-treatment interval of 14 days. <b>TO AVOID CROP DAMAGE:</b> Nursery stock, ornamentals and cut flowers/ foliage (other than certain apple varieties) are not known to be sensitive to Kenso Agcare Avior 250 SC Fungicide when used in strict accordance with the rate, conditions of use and other warnings. However, due to the large number of species and varieties of ornamentals and nursery stock it is impossible to test every one for tolerance to Kenso Agcare Avior 250 SC Fungicide. The user should conduct small-scale testing to ensure plant safety prior to large-scale commercial use.

OTHER CROPS				
Crop	Disease	Rate /100L	WHP	Critical Comments
Nursery stock and ornamentals including nursery stock (non-food and forestry), non-bearing fruit trees*, ornamentals and cut flowers/foilage *At least 6 months prior to first harvest	Myrtle rust ( <i>Uredo rangellii</i> )	40 mL	-	Apply by knapsack, powered hand-gun, boom or air-assisted spray. Apply in sufficient volume to ensure thorough coverage of all plant surfaces. Treat a sample area and assess appropriately prior to whole crop treatment to help minimise potential for phytotoxic damage. This is particularly important for crops in bloom. Apply a maximum of 2 consecutive applications of Kenso Agcare Avior 250 SC Fungicide with a minimum re-treatment interval of 14 days. <b>TO AVOID CROP DAMAGE:</b> Nursery stock, ornamentals and cut flowers/ foliage (other than certain apple varieties) are not known to be sensitive to Kenso Agcare Avior 250 SC Fungicide when used in strict accordance with the rate, conditions of use and other warnings. However, due to the large number of species and varieties of ornamentals and nursery stock it is impossible to test every one for tolerance to Kenso Agcare Avior 250 SC Fungicide. The user should conduct small-scale testing to ensure plant safety prior to large-scale commercial use.
Poppies	Downy Mildew	750 ml/ha	6 weeks	Apply Kenso Agcare Avior 250 SC Fungicide in a preventative fungicide program containing fungicides from a different chemical group. Apply Kenso Agcare Avior 250 SC Fungicide preventatively before disease symptoms appear. Ensure thorough spray coverage.
Potatoes	Early blight (Target Spot) ( <i>Alternaria solani</i> )	300 to 400 ml/ha	-	Apply the higher rates when climatic conditions favour Early Blight or Late Blight infection and in crops with large canopies. <b>Application</b> Kenso Agcare Avior 250 SC Fungicide may be applied by ground or aerial application equipment in potatoes. Aerial application may be used only for Early Blight (Target Spot) control. Apply in a sufficient volume of water to achieve thorough coverage of all foliage. The volume of water required to achieve this will depend on the stage of growth of the potatoes. <b>Ground Application:</b> A volume of 200 to 300 L/ha is suggested at the start of the season, increasing to 500 to 600 L/ha in a vigorous crop at full canopy. <b>Aerial Application (Early Blight only):</b> A volume of 30 to 40 L/ha is recommended. <b>Spray Interval</b> Consecutive applications should be applied at 7 to 14 day intervals at any time between early shoot growth and 14 days before harvest. Use the recommended shorter application interval in the following circumstances: 1. Under humid weather conditions which are favourable for Early or Late Blight infection. 2. When there is rapid vegetative growth during the early part of the crop cycle. 3. At the first sign of Late Blight infection.
	Late Blight ( <i>Phytophthora infestans</i> )	500 to 600 ml/ha		
	<b>Soil borne:</b> Black Scurf ( <i>Rhizoctonia solani</i> )  Silver Scurf* ( <i>Helminthosporium solani</i> )  *Suppression only.	5 to 10 ml/100m of row		
Pyrethrum	Ray Blight Disease ( <i>Phoma ligulicola</i> )	600 mL/ ha	DO NOT graze or cut treated area for stock food	DO NOT apply fungicides from the same chemical group more than 3 times in a season. Apply in sufficient water volume to achieve thorough coverage of all foliage.
Ribberries ( <i>Syzygium leuhamnii</i> and <i>S. fibrosum</i> ) Anise myrtle ( <i>S. anisatum</i> ) Lemon myrtle ( <i>Backhousia citriodora</i> )	Myrtle rust ( <i>Uredo rangellii</i> )	200 to 300 mL/ha	Fruit - 14 days Leaf - 4 months Grazing: 21 days	Apply 2 sprays with a minimum re-treatment interval of 14 days. Apply via ground based equipment on appearance of myrtle rust in a plantation or when conditions favour development of the disease. Use a maximum spray volume of 400 L/ha. <b>TO AVOID CROP DAMAGE</b> Ribberries, Anise myrtle and lemon myrtle are not known to be sensitive to this product when used in accordance with the label directions. The sensitivity of some species/varieties/cultivars, however, has not been fully evaluated under all growing conditions. It is advisable to only treat a small number of plants to ascertain their reaction before treating a larger area. Discontinue applications when any adverse symptoms post-treatment are observed.
<i>Rubus</i> (including: Raspberries, Blackberries, Boysenberries and Loganberries)	Anthrachnose ( <i>Elsinoe veneta</i> ) Botrytis ( <i>Botrytis cinerea</i> ) and Cladosporium ( <i>Cladosporium cladosporoides</i> )	80 mL	1 day	Begin applications at the onset of the disease. The applicable spray volume should be in the range of 500-1000 L/ha. Apply a maximum of 3 applications of Kenso Agcare Avior 250 SC Fungicide per season with a re-treatment interval of 14 days.
Tomatoes Except greenhouse	Early Blight (Target Spot) ( <i>Alternaria solani</i> )	400 mL/ha or 40 mL/100L	1 day	<b>Application</b> Apply in a sufficient volume of water to achieve thorough coverage of all foliage. The volume of water required to achieve this will depend on the stage of growth of the tomatoes and the method of trellising which influences canopy volume. In the case of dilute spraying (g/100 L) apply in the range of 400 to 500 L/ha after transplanting and increase to 800 to 1000 L/ha at full canopy. In the case of fully trellised tomatoes at full canopy, application volumes should be increased to 1500 L/ha to achieve these results with high volume spraying. For Late Blight and Sclerotinia controls use the higher rates when climatic conditions are humid and mild, which favours disease infection. <b>Spray Interval</b> Consecutive applications should be applied at 7 to 14 day intervals commencing soon after transplanting and continuing up to fruit maturity. Use the recommended shorter application interval in the following circumstances: 1. Under humid weather conditions which are favourable for disease infection. 2. When there is rapid vegetative growth during the early part of the crop cycle.
	Late Blight ( <i>Phytophthora infestans</i> ), Sclerotinia ( <i>Sclerotinia minor</i> )	500 to 600 mL/ha or 50 to 60 mL/100 L		

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

#### WITHHOLDING PERIODS

##### HARVEST

**Avocados:** DO NOT HARVEST FOR 7 DAYS AFTER APPLICATION.  
**Cucurbits, Passionfruit, Tomatoes:** DO NOT HARVEST FOR 1 DAY AFTER APPLICATION.  
**Grapes:** DO NOT HARVEST FOR 14 DAYS AFTER APPLICATION.  
**Mangoes:** DO NOT HARVEST FOR 3 DAYS AFTER APPLICATION.  
**Poppies:** DO NOT HARVEST FOR 6 WEEKS AFTER APPLICATION.  
**Potatoes, Pyrethrum:** NOT REQUIRED WHEN USED AS DIRECTED.  
**Olives:** DO NOT HARVEST FOR 21 DAYS AFTER APPLICATION.  
**Ribberries (*Syzygium* spp.):** DO NOT HARVEST FRUIT UNTIL 14 DAYS AFTER THE FINAL APPLICATION.  
**Anise myrtle and Lemon myrtle:** DO NOT HARVEST LEAVES UNTIL 4 MONTHS AFTER THE FINAL APPLICATION.  
**Carrot:** DO NOT HARVEST FOR 21 DAYS AFTER FINAL APPLICATION.  
**Rubus crops including, Blackberries, Raspberries, Loganberries and Boysenberries:** DO NOT HARVEST FOR 1 DAY AFTER APPLICATION.

##### GRAZING

**Pyrethrum:** DO NOT GRAZE OR CUT TREATED AREA FOR STOCK FOOD.  
**Anise Myrtle, Lemon Myrtle and Ribberries (*Syzygium* spp.):** DO NOT ALLOW LIVESTOCK TO GRAZE TREATED AREAS UNTIL 21 DAYS AFTER THE FINAL APPLICATION.

#### GENERAL INSTRUCTIONS

##### Application

DO NOT use concentration factors exceeding 4X when applying through low volume application equipment, except when applying Kenso Agcare Avior 250 SC Fungicide by air. In these cases adequate coverage of all plant surfaces is still required to achieve control of diseases.

#### TREE CROPS AND WINES

**Dilute spraying:** Use a sprayer designed to apply high volumes of water up to the point of run-off and matched to the crop being sprayed. Set up and operate the sprayer to achieve even coverage throughout the crop canopy. Apply sufficient water to cover the crop to the point of run-off. Avoid excessive run-off. The required water volume may be determined by applying different test volumes, using different settings on the sprayer, from industry guidelines or expert advice. Add the amount of product specified in the Directions of Use table for each 100L of water. Spray to the point of run-off. The required dilute spray volume will change and the sprayer set-up and operation may also need to be changed, as the crop grows.

**Concentrate spraying:** Use a sprayer designed and set-up for concentrate spraying (that is a sprayer which applies water volumes less than those required to reach the point of run off) and matched to the crop being sprayed. Set up and operate the sprayer to achieve even coverage throughout the crop canopy using your chosen water volume. Determine an appropriate dilute spray volume (see Dilute spraying above) for the crop canopy. This is needed to calculate the concentrate mixing rate. The mixing rate for concentrate spraying can then be calculated in the following way:

##### Example only

- Dilute spray volume as determined above: for example 1000 L/ha
- Your chosen concentrate spray volume: for example 500 L/ha
- The concentration factor in this example is: 2 x (ie 1000 L ÷ 500 L = 2)
- If the dilute label rate is 80 mL/100 L, then the concentrate rate becomes 2 x 80; that is 160 mL/100 L of concentrate spray.

The chosen spray volume, amount of product per 100 L of water, and the sprayer set up and operation may need to be changed as the crop grows. For further information on concentrate spraying, users are advised to consult relevant industry guidelines, undertake appropriate competency training and follow industry Best Practices.

#### Mixing

Half-fill the spray tank with clean water and start agitation. Shake the closed Kenso Agcare Avior 250 SC Fungicide container. Whilst filling the remainder of the spray tank add the required amount of Kenso Agcare Avior 250 SC Fungicide, adding any tank-mix products last. Maintain agitation until spraying is complete. DO NOT leave the spray mix in the sprayer overnight.

#### Export of Treated Produce

##### Grapes

While Maximum Residue Limits (MRLs) have been set in many major wine export destinations, some export destinations have not finalised MRL applications. For further information regarding export tolerances please contact your winery, Kenso Agcare Pty Ltd representative or the Australian Wine Research Institute.

##### Other Crops

While Maximum Residue Limits (MRLs) have been set in many major export destinations, it should be noted that MRLs or import tolerances may not be established in all export destinations. For further information regarding export tolerances please contact your export organisation or Kenso Agcare Pty Ltd representative.

### GROUP 11 FUNGICIDE

#### FUNGICIDE RESISTANCE WARNING

Kenso Agcare Avior 250 SC Fungicide is a member of the Quinone outside Inhibitors (QoIs) group of fungicides. For fungicide resistance management the product is a Group 11 fungicide. Some naturally occurring individual fungi resistant to the product and other Group 11 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by this product or other Group 11 fungicides, thus resulting in a reduction in efficacy and possible yield loss.

Since the occurrence of resistant fungi is difficult to detect prior to use, Kenso Agcare Pty Ltd accepts no liability for any losses that may result from the failure of this product to control resistant fungi.

#### RESISTANCE MANAGEMENT

Kenso Agcare Avior 250 SC Fungicide should be applied in a protective spray program containing fungicides from different chemical groups. DO NOT wait until disease levels have built up to make applications as this reduces the effectiveness of control and increases risk of resistance development. Disease control may be reduced if strains of pathogens less sensitive to Kenso Agcare Avior 250 SC Fungicide develop.

Kenso Agcare Avior 250 SC Fungicide should be applied as specified in the Directions for Use in association with the following CropLife Fungicide Resistance Management Strategies:

- DO NOT apply more than 1/3 of the total fungicide sprays per crop as Kenso Agcare Avior 250 SC Fungicide.
- A maximum of 2 consecutive applications of Kenso Agcare Avior 250 SC Fungicide are to be applied. They must be followed by at least the same number of applications of fungicide(s) from a different fungicide group(s), before Kenso Agcare Avior 250 SC Fungicide is used again in that crop.
- Where crops are grown successively alternation should continue between crops.

#### RE-ENTRY PERIODS

Do not enter treated areas until the spray has dried, unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing). Clothing must be laundered after each day's use.