

**CAUTION**  
KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

*KENSO AGCARE*  
**EPOXY**  
**125 SC**  
**FUNGICIDE**

ACTIVE CONSTITUENT: 125 g/L EPOXICONAZOLE

**GROUP 3 FUNGICIDE**

For the control of certain fungal diseases including stripe rust of wheat and specified fungal diseases of barley as per DIRECTIONS FOR USE table.



Kenso Corporation (M) Sdn Bhd  
3C / 59 Oxford Street  
Bulimba QLD 4171  
Phone (07) 3217 9788  
[www.kenso.com.au](http://www.kenso.com.au)

**IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USE**

**CONTENTS: 20 Litres APVMA Approval No.: 80799/ 101675**

# KENSO AGCARE EPOXY 125 SC FUNGICIDE

## STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, triple rinse, break, crush or puncture and deliver empty packaging or appropriate disposal at an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. Empty containers and product must not be burnt.

## SAFETY DIRECTIONS

Harmful if inhaled or swallowed. Will irritate the eyes, nose and throat and skin. Avoid contact with eyes and skin. DO NOT inhale vapours. If product on skin, immediately wash area with soap and water. Wash hands after use. When opening the container and preparing spray, wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow length PVC gloves and halfpiece respirator. When using the prepared spray, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and a washable hat. After each day's use, wash gloves, respirator and if rubber, wash with warm water. Wash contaminated clothing.

## FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26

## SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet which can be obtained from the supplier.

## CONDITIONS OF SALE

Kenso Corporation (M) Sdn. Bhd. will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Kenso Corporation (M) Sdn. Bhd. is limited to the replacement of the goods or (at the option of Kenso Corporation (M) Sdn. Bhd.) the refund of the price paid and where possible sufficient part of the goods to enable proper examination being returned to Kenso Corporation (M) Sdn. Bhd. within thirty days of delivery.

Product of Australia

In a Transport Emergency  
Dial **000**  
Police or Fire Brigade



9 328666 002932



**CAUTION**  
KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

**KENSO**  
*agcare*

**EPOXY  
125 SC**

Fungicide

ACTIVE CONSTITUENT: 125 g/L  
EPOXICONAZOLE

GROUP **3** FUNGICIDE

For the control of certain fungal diseases including stripe rust of wheat and specified fungal diseases of barley as per Directions for Use table.

READ THIS LEAFLET BEFORE USE

APVMA Approval No.: 80799/101675



**Kenso Corporation (M) Sdn. Bhd.**  
3C / 59 Oxford Street  
Bulimba QLD 4171  
Phone (07) 32179788

#### GENERAL INSTRUCTIONS

Kenso Agcare Epoxy 125 SC Fungicide is a broad spectrum contact and systemic fungicide. It is taken up via the stem and foliage and is translocated upwards and outwards to provide protection to new growth. Thorough coverage of the crop is necessary for best results. Check the crop regularly for any symptoms of disease infection. Apply as soon as disease is present as indicated in the Directions for Use.

#### MIXING

Kenso Agcare Epoxy 125 SC Fungicide is a suspension concentrate formulation. The addition of a non-ionic surfactant is generally not required, although under certain environmental conditions and lower total application volumes, the addition of a recommended 1000g/L surfactant may assist in providing better coverage of sprayed surfaces. Add the product to the half filled spray tank while agitating. Continue to agitate while topping up the tanks and during spraying.

#### APPLICATION

##### Ground Application

Apply in a water volume of between 50 and 100 L/ha, using flat fan nozzles operating at around 50cm above the top of the crop. Use the higher water volume in crops with heavier canopies.

##### Aerial Application

Apply with suitable aircraft, set up and operated to apply fungicides to cereal crops in a minimum water volume of 20L/ha.

#### COMPATIBILITY

For information on compatibility, please contact your local reseller or Kenso Agcare Pty Ltd representative.

#### STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, triple rinse, break, crush or puncture and deliver empty packaging or appropriate disposal at an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. Empty containers and product must not be burnt.

#### SAFETY DIRECTIONS

Harmful if inhaled or swallowed. Will irritate the eyes, nose and throat and skin. Avoid contact with eyes and skin. DO NOT inhale vapours. If product on skin, immediately wash area with soap and water. Wash hands after use. When opening the container and preparing spray, wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow length PVC gloves and halfpiece respirator. When using the prepared spray, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and a washable hat. After each day's use, wash gloves, respirator and if rubber, wash with warm water. Wash contaminated clothing.

#### FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

#### FUNGICIDE RESISTANCE WARNING

GROUP **3** FUNGICIDE

Kenso Agcare Epoxy 125 SC Fungicide is a member of the DMI group of fungicides. For fungicide resistance management the product is a Group 3 fungicide. Some naturally occurring individual fungi resistant to the product and other Group 3 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by this product or other Group 3 fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, Kenso Agcare Pty Ltd accepts not liability for any losses that may result from the failure of this product to control resistant fungi.

#### PRECAUTIONS

**Re-entry period:** DO NOT allow entry into treated areas until the spray has dried.

#### CAUTION

##### Export of treated cereals

Growers should note that Maximum Residue Limits (MRL's) or import tolerances do not exist in all markets for cereals treated with Kenso Agcare Epoxy 125 SC Fungicide. Additionally, some export markets have established MRLs different to those in Australia. If you are growing cereals for export, please check with Kenso Agcare Pty Ltd or your grain exporter for the latest information on MRLs and import tolerances BEFORE using Kenso Agcare Epoxy 125 SC Fungicide.

#### SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet which can be obtained from the supplier.

#### CONDITIONS OF SALE

Kenso Corporation (M) Sdn. Bhd. will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Kenso Corporation (M) Sdn. Bhd. is limited to the replacement of the goods or (at the option of Kenso Corporation (M) Sdn. Bhd.) the refund of the price paid and where possible sufficient part of the goods to enable proper examination being returned to Kenso Corporation (M) Sdn. Bhd. within thirty days of delivery.

In a Transport Emergency  
Dial **000**  
Police or Fire Brigade



**Kenso Corporation (M) Sdn. Bhd.**  
3C / 59 Oxford Street  
Bulimba QLD 4171  
Phone (07) 32179788

APVMA Approval No.: 80799/101675

#### LIVESTOCK DESTINED FOR EXPORT MARKETS

The grazing withholding period only applies to stock slaughtered for the domestic market. Some export markets apply different standards. To meet these standards, ensure that in addition to complying with the grazing withholding period, the Export Slaughter Interval is observed before stock are sold or slaughtered.

#### EXPORT SLAUGHTER INTERVAL (ESI)

AFTER OBSERVING THE WITHHOLDING PERIOD FOR GRAZING OR CUTTING FOR STOCKFOOD, LIVESTOCK THAT HAVE BEEN GRAZED OR FED TREATED CROPS SHOULD BE PLACED ON CLEAN FEED FOR 3 DAYS PRIOR TO SLAUGHTER.

#### PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET CROPS

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

#### PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Highly toxic to aquatic organisms. DO NOT apply Kenso Agcare Epoxy 125 SC Fungicide aerially if waterbodies, watercourses or wetlands are within 30 metres downwind of the application area. DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

## DIRECTIONS FOR USE

CROP	DISEASE	RATE	CRITICAL COMMENTS
Wheat	Leaf rust ( <i>Puccinia recondita</i> )	500 mL/ha	Apply when conditions favour disease development and prior to development of disease in the crop. Aim to apply between stem elongation and ear emergence complete (Z32-59) if disease is present. DO NOT apply later than Z59.
	Stripe rust ( <i>Puccinia striiformis</i> ) Septoria nodorum blotch ( <i>Phaeosphaeria nodorum</i> )	250 – 500 mL/ha	Apply when conditions favour disease development and prior to development of disease in the crop. Aim to apply between stem elongation and ear emergence complete (Z32-59) if disease is present. DO NOT apply later than Z59. Apply the higher rate when disease is present on the top leaf or conditions are favourable for disease development at these stages of crop development. Repeat spraying may be required, particularly if infection occurs early. Regularly monitor the crop from 3-4 weeks after the first application for signs of re-infection.
	Powdery mildew ( <i>Blumeria graminis f.sp. tritici</i> )	250 mL/ha	Apply when conditions favour disease development and prior to development of disease in the crop. Aim to apply during early tillering (Z21-22) stage of the crop if disease is present. DO NOT apply later than Z59. Repeat spraying may be required, particularly if infection pressure persists. Regularly monitor the crop from 3-4 weeks after the first application for signs of re-infection.
Barley	<b>DO NOT exceed a total rate of 500 mL/ha of Kenso Agcare Epoxy 125 SC Fungicide in a growing season</b>		
	Leaf rust ( <i>Puccinia hordei</i> )	250 – 500 mL/ha	Apply when conditions favour disease development and prior to development of disease in the crop. Aim to apply between stem elongation and ear emergence complete (Z32-59) if disease is present. DO NOT apply later than Z59. Apply the higher rate when disease is present on the top leaf or conditions are favourable for disease development at these stages of crop development.
	Net form of net blotch ( <i>Pyrenophora teres f. Sp. hordei</i> )	250 – 500 mL/ha	Apply when conditions favour disease development and prior to development of disease in the crop. Aim to apply between stem elongation and ear emergence complete (Z32-59) if disease is present. DO NOT apply later than Z59. Two applications of Kenso Agcare Epoxy 125 SC Fungicide at the lower rate may provide better control than a single application under certain conditions. If applying a two spray program, only apply 250 mL/ha per application, with the first application at Z31-32 if the disease is present. Aim to apply the second application around Z37-39, but no later than Z59. if infection pressure persists. DO NOT exceed a total rate of 500 mL/ha of Kenso Agcare Epoxy 125 SC Fungicide in barley per growing season.
	Leaf scald ( <i>Rynchosporium secalis</i> ) Powdery mildew ( <i>Blumeria graminis f. Sp. hordei</i> )	250 mL/ha	Apply when conditions favour disease development and prior to development of disease in the crop. Aim to apply between early tillering and flag leaf emergence (Z21-39) if disease is present. DO NOT apply later than Z59. Repeat spraying may be required, particularly if infection pressure persists. Regularly monitor the crop from 3-4 weeks after the first application for signs of re-infection. DO NOT exceed a total rate of 500 mL/ha of Kenso Agcare Epoxy 125 SC Fungicide in barley per growing season.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

### WITHHOLDING PERIODS

**DO NOT HARVEST FOR 6 WEEKS AFTER APPLICATION.**

**DO NOT GRAZE OR CUT FOR STOCKFOOD FOR 6 WEEKS AFTER APPLICATION.**