

CAUTION

KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

KENSO AGCARE

GUNDY 700 WG HERBICIDE

ACTIVE CONSTITUENT: 700 g/kg IMAZAPIC

GROUP B HERBICIDE

For the pre-emergence control of certain annual grass and broadleaf weeds in fallow situations and sugarcane as specified in DIRECTIONS FOR USE Table.

Kenso Corporation (M) Sdn Bhd
3C / 59 Oxford Street
Bulimba QLD 4171
Phone (07) 3217 9788
www.kenso.com.au

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USE



CONTENTS: 5 Kilograms APVMA Approval No.: 69339/ 60532

KENSO AGCARE GUNDY 700 WG HERBICIDE

STORAGE AND DISPOSAL

KEEP OUT OF REACH OF CHILDREN.

Store in the closed, original container, in a cool, dry, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Shake and empty contents into spray tank. DO NOT dispose of undiluted chemicals on site. Break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

SAFETY DIRECTIONS

Will irritate the eyes and skin. Avoid contact with eyes and skin. When opening the container and preparing spray, wear elbow-length PVC gloves and face-shield or goggles. If product in eyes, wash it out immediately with water. Wash hands after use. After each days use, wash gloves and face shield or goggles.

FIRST AID

If poisoning occurs, contact a Doctor or Poisons Information Centre (Phone Australia 13 11 26).

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet which can be obtained from the supplier.

CONDITIONS OF SALE

Kenso Corporation (M) Sdn. Bhd. will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Kenso Corporation (M) Sdn. Bhd. is limited to the replacement of the goods or (at the option of Kenso Corporation (M) Sdn. Bhd.) the refund of the price paid and where possible sufficient part of the goods to enable proper examination being returned to Kenso Corporation (M) Sdn. Bhd. within thirty days of delivery.

In a Transport Emergency
Dial **000**
Police or Fire Brigade

Product of China



9 528666 0027 89



Batch No.:

Date of Manufacture:

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

KENSO
 agcare

GUNDY 700 WG

Herbicide

ACTIVE CONSTITUENT: 700 g/kg IMAZAPIC

GROUP B HERBICIDE

For the pre-emergence control of certain annual grass and broadleaf weeds in fallow situations and sugarcane as specified in DIRECTIONS FOR USE Table.

READ THIS LEAFLET BEFORE USE

APVMA Approval No.: 69339/60532



Kenso Corporation (M) Sdn. Bhd.

3C / 59 Oxford Street

Bulimba QLD 4171

Phone (07) 32179788

SAFETY DIRECTIONS

Will irritate the eyes and skin. Avoid contact with eyes and skin. When opening the container and preparing spray, wear elbow-length PVC gloves and face-shield or goggles. If product in eyes, wash it out immediately with water. Wash hands after use. After each days use, wash gloves and face shield or goggles.

FIRST AID

If poisoning occurs, contact a Doctor or Poisons Information Centre (Phone Australia 13 11 26).

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet which can be obtained from the supplier.

CONDITIONS OF SALE

Kenso Corporation (M) Sdn. Bhd. will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Kenso Corporation (M) Sdn. Bhd. is limited to the replacement of the goods or (at the option of Kenso Corporation (M) Sdn. Bhd.) the refund of the price paid and where possible sufficient part of the goods to enable proper examination being returned to Kenso Corporation (M) Sdn. Bhd. within thirty days of delivery.

In a Transport Emergency
 Dial **000**
 Police or Fire Brigade



Kenso Corporation (M) Sdn Bhd

3C / 59 Oxford Street

Bulimba QLD 4171

Phone (07) 3217 9788

APVMA Approval No: 69339/60532

DIRECTIONS FOR USE

RESTRAINTS:

DO NOT apply to soils saturated with water.

DO NOT apply by aircraft.

SITUATION	WEEDS CONTROLLED	RATE g/ha	CRITICAL COMMENTS
Fallow prior to planting wheat, barley or chickpea (Qld and Nthn NSW only)	Barnyard grass (<i>Echinochloa colona</i>), liverseed grass (<i>Urochloa panicoides</i>), stink grass (<i>Eragrostis ciliaris</i>), blowaway grass (<i>Panicum decompositum</i>), button grass (<i>Dactyloctenium radulans</i>), Pigweed (<i>Portulaca oleracea</i>), yellow vine (<i>Tribulus terrestris</i>), mintweed (<i>Salvia reflexa</i>), Boggabri weed (<i>Amaranthus mitchellii</i>), dwarf amaranth (<i>Amaranthus macrocarpus</i>), peachvine/cowvine (<i>Ipomea lonchophylla</i>)	50 to 70	Treatments may be applied from the end of the previous crop harvest but not less than 4 months prior to planting of winter crop. Best results are obtained where treatments are applied to dry, weed-free soil prior to weed germination. Treatments can be applied to standing crop residues. Sufficient rainfall is required after application and prior to weed emergence to wet soil to a depth of 5 cm. Use the higher rates for increased residual control of weeds or anticipated high weed pressure. Emerged weeds must be controlled by prior cultivation, or by application of a knockdown herbicide. Where infestation of emerged weeds is light, tank mix treatment with Ken-Up 450 CT. Cultivation following application of treatments may reduce pre-emergence weed control provided by Kenso Agcare Gundy 700 WG Herbicide. Prolonged wet soil conditions following treatment will reduce the effective life of pre-emergence treatments. Weed escapes may require follow-up application of knockdown herbicides. DO NOT apply more than 70 g/ha per season on cropland. DO NOT apply in more than two successive seasons on cropland. DO NOT overspray headlands.
Sugarcane (Pre-emergence to variety Q124 only) (Qld, NSW and WA only)	Cobbler's peg (<i>Bidens pilosa</i>), common sida (<i>Sida rhombifolia</i>), glossy nightshade (<i>Solanum americanum</i>), green amaranth (<i>Amaranthus viridis</i>), green summer grass (<i>Brachiaria subquadriflora</i>),	140	DO NOT apply to emerged sugarcane as damage may result. PLANT CANE: Apply after planting and before emergence of the sugarcane. RATOON: Apply after harvest and before new cane growth appears. Heavy trash layers may prevent the product from reaching the soil. Apply to weed-free soil prior to weed germination. Sufficient rainfall, or overhead irrigation, is required after application and prior to weed emergence to wet soil to a depth of 5 cm. Emerged weeds must be controlled by prior cultivation, or application of an appropriate herbicide. Cultivation following application of treatments may reduce pre-emergence weed control provided by Kenso Agcare Gundy 700 WG Herbicide. PLEASE TURN OVER

SITUATION	WEEDS CONTROLLED	RATE g/ha	CRITICAL COMMENTS
	<i>Ipomoea</i> spp., milkweed (Mexican fire plant) (<i>Euphorbia heterophylla</i>), summer grass (<i>Digitaria ciliaris</i>)		Heavy rain immediately after application may concentrate Kenso Agcare Gundy 700 WG Herbicide in the furrow and cause temporary yellowing of cane leaves. Prolonged wet soil conditions following treatment will reduce the effective period of weed control. Weed escapes may require follow-up application of an appropriate herbicide.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD:

WHEAT, BARLEY, CHICKPEA STUBBLE:

DO NOT GRAZE TREATED STUBBLE OR WEEDS WITHIN 4 WEEKS AFTER TREATMENT.

SUGARCANE:

NOT REQUIRED WHEN USED AS DIRECTED.

GENERAL INSTRUCTIONS

Fallow application: Kenso Agcare Gundy 700 WG Herbicide is for use in fallow prior to the planting of wheat, barley or chickpea and is a pre-emergence herbicide which is recommended to provide short term (6-12 weeks) pre-emergence control of summer germinating grasses and broadleaf weeds in cultivated or uncultivated fallow.

Sugarcane application: Kenso Agcare Gundy 700 WG Herbicide should be applied to plant or ratoon cane before the cane has emerged. If emerged weeds are present, Kenso Agcare Gundy 700 WG Herbicide should be mixed with an appropriate knockdown herbicide.

MIXING

Kenso Agcare Gundy 700 WG Herbicide is a Water Dispersible granule formulation. Part fill the spray tank with water, then with agitator running, add the required amount of product, then fill the tank with water.

When tank mixing this product with other recommended compatible products, first add the other product to the tank and mix thoroughly before adding this product. No adjuvants or surfactants are required when Kenso Agcare Gundy 700 WG Herbicide is used alone for pre-emergence weed control.

APPLICATION

DO NOT apply by aircraft.

Fallow application: For ground application only: Apply with boom equipment in not less than 50 L/ha water using flat fan nozzles. Avoid overlap and do not overspray headlands.

Sugarcane: Ensure thorough coverage of soil with well-calibrated spray equipment. Apply in 50 to 200 L water/ha.

EQUIPMENT CLEAN-UP

Thoroughly flush all spray equipment with water following the use of Kenso Agcare Gundy 700 WG Herbicide and before use with other products.

COMPATIBILITY

Kenso Agcare Gundy 700 WG Herbicide is compatible with glyphosate, paraquat, Ken-Met 600 and Triclopyr 600. It is NOT compatible with dicamba or Ken-Trel 300.

RESISTANT WEEDS WARNING

GROUP B HERBICIDE

Kenso Agcare Gundy 700 WG Herbicide is a member of the imidazolinone group of herbicides. The product has the inhibitors-of-acetolactate-synthase (ALS) mode of action. For weed resistance management, the product is a Group B herbicide. Some naturally-occurring weed biotypes resistant to this product and other inhibitors-of-acetolactate-synthase (ALS) herbicides, may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population, if these herbicides are used repeatedly. These resistant weeds will not be controlled by this product or other inhibitors-of-acetolactate-synthase (ALS) herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Kenso Agcare Australia Ltd accepts no liability for any losses that may result from the failure of this product to control resistant weeds.

FOLLOW CROPS

Under conditions which do not favour breakdown of this product, carry-over soil residues can affect susceptible follow crops.

As environmental and agronomic factors make it impossible to eliminate all risks associated with this product, rotational crop injury is always possible.

The following minimum re-cropping intervals (months after application) should be observed.

Following use in summer fallow:-

MONTHS AFTER APPLICATION			
4	12	24	36
wheat*, barley*, chickpea, peanuts	mungbean, lucerne	sorghum, cotton, sunflower	any other crops

* The following additional requirements apply if it is intended to plant wheat or barley during the next winter season.

Do NOT apply Kenso Agcare Gundy 700 WG Herbicide in fallow later than the end of December.

Do NOT apply Kenso Agcare Gundy 700 WG Herbicide in areas where rainfall from spraying to sowing of cereals is expected to be below 200 mm.

Following use in sugarcane: Observe the minimum re-cropping intervals listed above under summer fallow.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

Do NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

Do NOT spray within 50 m of wetlands or waterways.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Do NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

KEEP OUT OF REACH OF CHILDREN.

Store in the closed, original container, in a cool, dry, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Shake and empty contents into spray tank.

DO NOT dispose of undiluted chemicals on site. Break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.