

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

KENSO AGCARE

KEN-UP 450 CT

NON-SELECTIVE, TRANSLOCATED HERBICIDE

ACTIVE CONSTITUENT: 450 g/L GLYPHOSATE present as the ISOPROPYLAMINE SALT

Also contains: 144 g/L polyethanoxy (15) tallow amine

GROUP M HERBICIDE

For the control of many annual and perennial grasses and broadleaf weeds prior to crop establishment and as an aid to pasture management through manipulation and seed set control as per directions for use table.

Kenso Corporation (M) Sdn Bhd
Level 1, 98 Commercial Road,
Teneriffe QLD 4005
Phone (07) 3216 1188
www.kenso.com.au



IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USE

CONTENTS: 20 Litres APVMA Approval No.: 55080/ 115598

KENSO AGCARE KEN-UP 450 CT NON-SELECTIVE, TRANSLOCATED HERBICIDE

STORAGE AND DISPOSAL

Store in the closed original container in a well ventilated area as cool as possible. Do not store for prolonged periods in direct sunlight.

Triple-rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

SAFETY DIRECTIONS

Product will irritate the eyes and skin. Avoid contact with eyes and skin. When preparing product for use, wear elbow length PVC gloves and face shield or goggles. When using controlled droplet applicators wear protective waterproof clothing and impervious footwear. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each days use, wash gloves, face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131 126. If skin contact occurs, remove contaminated clothing and wash skin thoroughly. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet, which can be obtained from www.kenso.com.au.

CONDITIONS OF SALE

Kenso Corporation (M) Sdn. Bhd. will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Kenso Corporation (M) Sdn. Bhd. is limited to the replacement of the goods or (at the option of Kenso Corporation (M) Sdn. Bhd.) the refund of the price paid and where possible sufficient part of the goods to enable proper examination being returned to Kenso Corporation (M) Sdn. Bhd. within thirty days of delivery.

In a Transport Emergency
Dial **000**
Police or Fire Brigade



Batch No.:
Date of Manufacture:

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

KENSO
agcare

KEN-UP
450 CT

Non-Selective, Translocated Herbicide

ACTIVE CONSTITUENT: 450 g/L GLYPHOSATE
present as the ISOPROPYLAMINE SALT

Also contains: 144 g/L polyethanox (15) tallow amine

GROUP M HERBICIDE

For the control of many annual and perennial grasses and broadleaf weeds prior to crop establishment and as an aid to pasture management through manipulation and seed set control as per directions for use table.

READ THIS LEAFLET BEFORE USE

APVMA Approval No.: 55080/115598



Kenso Corporation (M) Sdn. Bhd.
Level 1, 98 Commercial Road,
Teneriffe QLD 4005
Phone (07) 3216 1188

Tank Mixtures — How to Choose

Use the following list of suggestions to determine the appropriate tank mixtures. Adhere to manufacturers' instructions in all cases.

- **To improve knockdown and give residual weed control** -Atrazine (flowable only) can be added. Use with ammonium sulphate to overcome antagonism. Do not spray by air. This mixture does not control barnyard grass.
- **-in fallow or in crop.** Ken-Chlor 750 can be added.
- **-prior to sowing lupins.** Simazine (flowable only) can be added. Use ammonium sulphate to overcome antagonism.
- **To improve and accelerate knockdown symptoms prior to planting wheat or barley.** Add 75 mL of Ox 240 per hectare.
- **To improve control of certain broadleaved weeds.** Add Ken-Ester LV 680. Observe any regional restrictions on use.
- **To improve control of Sorrel, Subterranean Clover, Medics, and White Clover.** Add Kenso Agcare Dicamba 500. Observe any regional restrictions on use.
- **To improve control of Annual Ryegrass, Silver grass, and perennial grasses.** Add 200 mL of Silken Penetrant per 100 L of spray mix. Use only if recommended.
- **To compensate for adverse growing conditions such as cool or cloudy weather.** Control of annual weeds by Ken-Up 450 CT Non-Selective, Translocated Herbicide alone may possibly be improved by adding 2L of ammonium sulphate per 100 L of spray mix.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

Avoid contact with foliage, green stems or fruit of crops, desirable plants and trees, as severe injury or destruction may result.

DO NOT apply under weather conditions, or from spraying equipment that may cause spray to drift onto nearby susceptible plants, crops, cropping lands or pastures.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Do not contaminate dams, rivers or streams with the product or used container. Do not apply to weeds growing in or over water. Do not spray across open bodies of water.

STORAGE AND DISPOSAL

Store in the closed original container in a well ventilated area as cool as possible. Do not store for prolonged periods in direct sunlight.

Triple-rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

SAFETY DIRECTIONS

Product will irritate the eyes and skin. Avoid contact with eyes and skin. When preparing product for use, wear elbow length PVC gloves and face shield or goggles. When using controlled droplet applicators wear protective waterproof clothing and impervious footwear. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each days use, wash gloves, face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131 126. If skin contact occurs, remove contaminated clothing and wash skin thoroughly. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet, which can be obtained from www.kenso.com.au.

CONDITIONS OF SALE

Kenso Corporation (M) Sdn. Bhd. will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Kenso Corporation (M) Sdn. Bhd. is limited to the replacement of the goods or (at the option of Kenso Corporation (M) Sdn. Bhd.) the refund of the price paid and where possible sufficient part of the goods to enable proper examination being returned to Kenso Corporation (M) Sdn. Bhd. within thirty days of delivery.

In a Transport Emergency
Dial **000**
Police of Fire Brigade

* Not a registered trademark of Kenso Corporation



Kenso Corporation (M) Sdn. Bhd.
Level 1, 98 Commercial Road,
Teneriffe QLD 4005
Phone (07) 3216 1188

APVMA Approval No.: 55080/115598

DIRECTIONS FOR USE:

Restraints

To ensure herbicide absorption, DO NOT disturb weeds by cultivation, sowing or grazing for 1 day after treatment of annual weeds and 7 days for perennial weeds, except where noted.

Crop/Situation	State	Weeds controlled	Rate Vol/ha	Critical comments
TREE AND VINE CROPS Avocado, Banana, Blueberries, Citrus fruits, Custard apples, Duboisa, Figs – dessert, Guava, Kiwifruit, Litchi, Mango, Monstera – fruit, Nuts (including Almond, Pecan, Macadamia, Pistachio and Walnut), Olives, Pawpaw, Persimmons, Pome fruit, Raspberries, Stone fruit, Tea, Vineyards	All States	Amaranth, Barley grass, Brome grass, Barnyard grass, Caltrop, Canary grass, Capeweed, Chickweed, Deadnettle, Doublegee, Liverseed Grass, Mintweed, Paterson's Curse, Pigweed, Ryegrass, Silvergrass, Spear thistle, Thornapple, Wild mustard, Wild oats, Wild turnip, Winter grass, Variegated thistle	Boom : 1.6 – 2.4 L/ha Handgun : 400 – 600 mL per 100 L Knapsack : 60 – 80 mL per 15 L	Apply as a directed or shielded spray or using wiper equipment. Do NOT apply as a spray near trees or vines less than 3 years old unless they are effectively shielded from spray and spray drift. Do NOT allow wiper surface to contact any part of the tree, vine or palm. Citrus fruit, Nuts, Olives, Pome Fruit & Vineyards Do NOT allow spray to drift to contact green bark or stems, canes, laterals, suckers, fresh wounds, foliage or fruit. Tea Apply a maximum of 2.4 L/ha by shielded boom or directed off-centre nozzle or 0.4 L/100 L by directed handgun or knapsack to avoid application to the crop. All other crops Do NOT allow spray or spray drift to contact any part of the plant including the trunk. CAUTION Where split bark on Kiwifruit and green stems on Pawpaw occur, extreme care is required. Annual weeds may be sprayed anytime they are actively growing. Use the lower rate on weeds, up to 15 cm tall.
SOUTHERN AUSTRALIA Prior to sowing a crop or pasture with full soil disturbance by cultivation or sowing with a tyned implement	WA, SA, VIC, NSW only	Barley Grass, Brome grass, Volunteer cereals, Wild oats Annual phalaris (Canary grass), Annual ryegrass, Silvergrass, Winter grass Calomba daisy, Capeweed, Doublegee (Spiny Emex) Amsinckia, Fumitory, Paterson's Curse, Saffron Thistle, Scotch Thistle, Spear thistle, Variegated thistle, Volunteer lupins, Wild turnip Dock (Seedling)	400 – 800 mL pre tillering 800 mL – 1.0L post tillering 800 mL – 1.0L pre tillering 1.0 L – 1.2 L post tillering 400 – 800 mL less than 8 cm diameter 800 mL – 1.2L greater than 12 cm diameter 800 mL – 1.0 L less than 12 cm diameter 1.0 L – 1.2L greater than 12 cm diameter 800 mL – 1.2L	Treat actively growing weeds not under stress from low moisture, frost, cold, disease or waterlogging. If heavy grazing has occurred allow regrowth to 6-8 cm before spraying and use the higher rate. RATE SELECTION: Increase to higher rates late in the season or when treating under cold/overcast conditions. FULL DISTURBANCE with cultivation or sowing with a tyned implement may start 1 day after treatment (7 days if Dock, Phalaris, Skeleton weed, Soursob or Sorrel are present) and should occur within 21 days after treatment. Where cultivation or sowing does not occur within 21 days, new weed growth may require further treatment. When treating light infestations of seedling annual grasses (pre-tillering) and annual broadleaved weeds (less than 8cm dia/height) cultivation or sowing may start 6 hours after treatment and should occur within 21 days. CROP ESTABLISHMENT: Sowing should not proceed until conditions allow the formation of a satisfactory seedbed. See Crop Establishment for directions ANNUAL RYEGRASS, SILVERGRASS AND PERENNIAL GRASSES: Addition of Silken Penetrant, 200mL/100L of spray solution may improve control. When treating dense infestations of Silvergrass, use of low volume nozzles (eg. SS 11001, Hardi No 10) and a spray volume of 70 L/Ha or more is recommended to improve plant spray coverage. TANK MIXTURES: For improved control of clover add Kenso Agcare Dicamba 500 Selective Herbicide. Read and follow all label directions, restraints, plant back periods, withholding periods, regional use restrictions and safety directions for the tank mix products. See Tank Mixtures for directions PERENNIAL WEEDS: For Perennial phalaris, Soursob, Skeleton weed and Sorrel, this product will provide knockdown, seasonal suppression and reduction in treated plant numbers.
	Tas only	All the above weeds	1.2 – 2.4 L	TASMANIA: Use 1.2L/ha on annual weeds. Increase to 2.4L/ha where perennial weeds are being treated. To control White clover and improve control of Sorrel and Dock, add 1L/ha of Kenso Agcare Dicamba 500 Selective Herbicide. Observe Kenso Agcare Dicamba 500 Selective Herbicide label directions and plant back periods.

Crop/Situation	State	Weeds controlled	Rate Vol/ha	Critical comments
SOUTHERN AUSTRALIA Prior to establishing a crop or pasture with an implement that gives minimal or no soil disturbance.	NSW, SA, VIC, WA, only	Barley grass, volunteer cereals, Wild oats Brome grass, Canary grass, Capeweed, Variegated thistle, Winter grass Annual ryegrass, Paterson's curse, Saffron thistle, Scotch thistle, Silvergrass, Spear thistle, Wild mustard, Wild radish, Wild turnip Erodium, Perennial phalaris, Plantain, Sorrel, Sub clover, Yorkshire fog	800 mL – 1.2 L 1.0 – 1.6 L 1.2 – 1.6 L 1.5 – 2.0 L	Treat actively growing weeds not under stress from low moisture, frost, cold, disease or waterlogging. If heavy grazing of mature plants has occurred allow regrowth to 6-8 cm before spraying and use the higher rate. RATE SELECTION: Use the lower rate on young weeds, increase to the higher rate where grasses reach full tillering or where broadleaf weeds reach stem elongation/budding. Increase to higher rates in spring or when treating under cold/overcast conditions. AERIAL APPLICATION: Use the higher rates. See also Aerial Application . ANNUAL RYEGRASS, SILVERGRASS AND PERENNIAL GRASSES: Add Silken Penetrant, 200mL/100L of spray solution. When treating dense infestations of Silvergrass, use of low volume nozzles (eg. SS 11001, Hardi No 10) and a spray volume of 70 L/Ha or more is recommended to improve plant spray coverage. TANK MIXTURES: For improved control of Dock, Sorrel and Sub clover add Kenso Agcare Dicamba 500 Selective Herbicide. Read and follow all label directions, restraints, plant back periods, withholding periods, regional use restrictions and safety directions for the tank mix products. See Tank Mixtures for directions. Addition of Ammonium Sulphate, 2kg/100L, may improve control when treating under adverse environmental conditions. PASTURE OR CROP ESTABLISHMENT: Do NOT sow into excessive trash. Trash may be removed by grazing after treatment. Grazing may commence one day after treatment of annual weeds (small) and 7 days for perennial weeds. Delay grazing for 3 days where annual weeds are large. Sowing may proceed when excessive trash is removed, but not sooner than one day after treatment of annual weeds and 7 days for perennial weeds. See also Crop Establishment
	Tas only	All the above weeds	1.2 – 2.4 L	AERIAL (OR SURFACE) SEEDING: Delay seeding until trash is completely removed by grazing and/or plant decay. When establishing pasture, ensure application of fertilizer and insecticides and follow-up management is undertaken as required. TASMANIA: Use 1.2L/ha on annual weeds. Increase to 2.4L/ha where perennial weeds are being treated. To control White clover and improve control of Sorrel and Dock, add 1L/ha of Kenso Agcare Dicamba 500 Selective Herbicide. Observe Kenso Agcare Dicamba 500 Selective Herbicide label directions and plant back periods.
SOUTHERN AUSTRALIA To commence a fallow	NSW, VIC, SA, WA only	Barley grass, volunteer cereals, Wild oats Annual ryegrass, Brome grass, Capeweed, Paterson's curse (rosette), Saffron thistle, Scotch thistle, Silvergrass, Spear thistle, Wild mustard, Wild radish, Wild turnip Hoary cress, Soursob Couch	800 mL – 1.2 L 1.2 - 1.6 L 1.2 L 1.2 – 2.4 L	Treat actively growing weeds not under stress from low moisture, frost, cold, disease or waterlogging. If heavy grazing has occurred allow regrowth to 6-8 cm before spraying and use the higher rate. RATE SELECTION: Use the lower rate on young weeds, or where cultivation is to follow within 21 days, increasing to the higher rate where grasses reach full tillering or where broadleaf weeds reach stem elongation/budding. ANNUAL RYEGRASS, SILVERGRASS AND PERENNIAL GRASSES: Add Silken Penetrant, 200mL/100L of spray solution. When treating dense infestations of Silvergrass, use of low volume nozzles (eg. SS 11001, Hardi No 10) and a spray volume of 70 L/Ha or more is recommended to improve plant spray coverage. HOARY CRESS: Treat from late rosette to early flowering SOURSOB: Treat at tuber exhaustion COUCH: Use the higher rate on dense infestations. Apply sequential treatments during summer and autumn, with autumn being most effective. Repeat applications will be required for full control. For improved control use in conjunction with cultivation. TANK MIXTURES: Read and follow all label directions, restraints, plant back periods, withholding periods, regional use restrictions and safety directions for the tank mix products. See Tank Mixtures for directions.
	TAS only	All the above weeds	1.2 – 2.4 L	TASMANIA: Use 1.2L/ha on annual weeds. Increase to 2.4L/ha where perennial weeds are being treated. To control White clover and improve control of Sorrel and Dock, add 1L/ha of Kenso Agcare Dicamba 500 Selective Herbicide. Observe Kenso Agcare Dicamba 500 Selective Herbicide label directions and plant back periods
PASTURE TOPPING For annual grass, Capeweed and Calomba daisy seed-set reduction	WA, SA, VIC, TAS, NSW only	Barley grass, Brome grass, Capeweed, Silvergrass Annual ryegrass, Calomba daisy	240 mL – 360 mL 360 mL	Remove stock prior to treatment to allow even regrowth. Apply to Capeweed and Annual ryegrass at FLOWERING. For other grasses apply from HEAD to MILKY DOUGH stage. Use the higher rate for dense infestations or where Annual ryegrass is present. Apply before signs of plants "haying off". Reduction in pasture legume population may occur as a result.. DO NOT apply to clover or medic crops intended for seed or hay.
SEED- HEAD SUPPRESSION OF PERENNIAL GRASSES	VIC, TAS, NSW, WA, SA, only	Bent grass	300 – 500 mL	TIMING : Treat from late October to late November. Apply before seedheads have emerged. Use the higher rate where growth is excessive and renovation is intended the following autumn. FOLLOW-UP MANAGEMENT: Graze hard after spraying
BENT GRASS INFESTED PASTURE For control/ suppression prior to establishing crops or improved pasture species	VIC, TAS only	Most annual weeds and Bent grass	2.0 L	TIMING: Apply to actively growing plants in late spring when they have some seed-head development, but before summer moisture stress. Remove stock to ensure there is full leaf growth. FOLLOW-UP MANAGEMENT: Full disturbance with a tyned implement should follow 10-21 days after spraying. Then follow with a summer crop, and/or reseeded pasture or crop the following autumn.

PASTURE MANIPULATION For suppression or control of pasture species prior to drilling, improved pasture, forage species, soybeans or Leucaena BAND SPRAYING: May also be applied as a band or strip spray	NSW, VIC, WA only	Carpet grass, Kikuyu, Paspalum	1.1 – 4.8 L	RATE SELECTION: For suppression apply the low rate. Where complete control is required apply up to the high rate BAND SPRAYING: Band spraying may be done immediately after the sowing operation. Mount the nozzles behind the coulters/tyne/press wheel assembly of the band seeder. Adjust to spray 0.5 to 1.0m strips. Ensure minimal disturbance of the pasture. Excessive dust created in the seeding operation may reduce herbicide activity. Pasture seed must be drilled at the appropriate depth and covered by soil. LEUCAENA (QLD ONLY): Apply 2L/ha through a single taper fan nozzle LF1-80 mounted at the rear of the single row planter providing a 1m swath. Planting rows to be 4m apart.
	QLD only	Carpet grass, Paspalum Kikuyu Barbed wire grass, Black speargrass, Love grasses, Red Natal grass, Wire grasses	1.1 – 4.8 L 500 mL – 4.8 L 2.0 L	
POA TUSsock INFESTED PASTURE For reduction of ground cover allowing pasture renovation	NSW, Tas, VIC, QLD only	Most annual weeds and suppression of Poa tussock	2.4 – 3.2 L	TIMING: Graze heavily, then remove stock at least 14 days before spraying to allow fresh regrowth. Apply to actively growing plants after the autumn break but before heavy frosts (March – May) APPLICATION: Increasing to the higher rate may give more effective reductions. If aerial spraying see Aerial Equipment. FOLLOW-UP MANAGEMENT: Sowing may start from 14 days after spraying. It is essential that correct follow-up pasture establishment and management occurs after treatment. Spot treatment will limit re-infestations.

Crop/Situation	State	Weeds controlled	Rate Vol/ha	Critical comments
COTTON PRE-HARVEST Do not use on crops intended for seed production	QLD, NSW only	Bathurst burr, Noogoora burr, Winter annual weeds including Sowthistle/ Milkthistle	1.0 – 2.0 L	Use the lower rate on light infestations of small weeds, where the crop canopy allows adequate spray coverage of the weeds. Increase to the higher rate when the crop canopy may limit spray coverage, when treating dense infestations, or when treating larger weeds. Apply alone or in tank mixtures with Kenso Agcare Thiazuron 500 SC Cotton Defoliant. Apply when at least 60% of bolls are open and immature bolls cannot be easily cut with a sharp knife. Where a leafy canopy limits spray coverage, reduced weed control can be expected. For better results under these conditions, delay application until canopy re-opens following initial conditioning treatment. Where control of Nutgrass or Noogoora burr is required, treatment should be applied prior to the onset of frosts. When tank mixed with defoliant, a slightly higher proportion of cotton leaf may be retained, particularly where the higher rate is used. Read and follow all label directions for tank mix products.
		Nutgrass (seasonal suppression only)	2.0 L	
COTTON Shielded Sprayers	QLD, NSW only	Refer to weeds controlled section Northern Australia in fallows or prior to sowing a crop		Apply this product to weeds growing between crop rows using a shielded sprayer. Do not apply in crops less than 20 cm high. Do not allow spray or spray drift to contact any part of the cotton plants as severe injury or destruction may result.

KNAPSACK RATE:

Annual Weeds including :	APPLICATION METHOD	WEEDS CONTROLLED	RATE	CRITICAL COMMENTS
Barley Grass, Barnyard grass, Capeweed, Doublegee, Fumitory, Paterson's curse, Potato weeds, Ryegrass, Sub. clover, Sowthistle, Milkthistle, Spear thistle, Wild oats, Winter grass	Leaf spray	General mixed weeds (Annual and perennial weeds)	8mL per 1L water	Timing: Apply when weeds are actively growing. For best results on perennial weeds treat after flowering. Treat woody weeds from flowering to leaf fall (Jan-May). DO NOT spray plants bearing edible berries. Application: Uniform and complete spray to wet all leaves. Run off should not occur. 1 litre of spray will cover 10m ² . The addition of a Silken Penetrant is recommended at a rate of 1mL of a 600g/L product (or equivalent) per 1L spray solution.
Perennial Weeds including : Couch grass, Johnson grass, Lovergrass African, Paspalum. Dock, Kikuyu, Nutgrass, Soursob.		Woody weeds	12mL per 1L water	
Woody Weeds including : Blackberry, Groundsel Bush, Lantana, Boxthorn-African, Hawthorn, Sweet briar.		Annual weeds (small)	4mL per 1L water	

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHOLDING PERIOD: not required when used as directed.

GENERAL INSTRUCTIONS

Ken-Up 450 CT Non-Selective, Translocated Herbicide is a non-selective herbicide that will kill most emerged weeds and plants. It provides no residual weed control and may therefore be used before sowing any crop, but not prior to transplanting tomato seedlings. If required, Ken-Up 450 CT Non-Selective, Translocated Herbicide can be mixed with certain other herbicides to achieve both knockdown and residual weed control (see Tank Mixtures).

Ken-Up 450 CT Non-Selective, Translocated Herbicide is absorbed by foliage and green stems and moves into the root system. Weeds should be actively growing when treated. Do not treat weeds under poor growing or dormant conditions (such as occur in drought, waterlogging, disease, insect damage or following frosts), as reduced weed control may result. Reduced results may also occur when treating weeds heavily covered with dust or silt. Prior herbicide application may also induce stress in weeds. For annual weeds, it may be 3 to 7 days before the effects of the treatment become apparent; for perennials, it may be as much as 2 to 3 weeks or more if cool and cloudy. Ken-Up 450 CT Non-Selective, Translocated Herbicide is a non-volatile liquid that mixes readily with water. Just mix with the required volume of water, add Silken Penetrant, and spray.

Rain occurring up to 6 hours after application may reduce effectiveness. Heavy rainfall within 2 hours of application may wash the chemical off the foliage and a repeat treatment may be required. Delay treatment of plants wet with dew or rain, if water droplets runoff when plants are disturbed. Avoid contact with foliage, green stems or fruit of crops, desirable plants and trees, as severe injury or destruction may result. Ken-Up 450 CT Non-Selective, Translocated Herbicide is compatible with certain herbicides, insecticides and additives (see Compatibility).

The active constituent of Ken-Up 450 CT Non-Selective, Translocated Herbicide is 450 g/L glyphosate present as the isopropylamine salt. A withholding period for stock is not required, however, it is recommended that grazing of treated plants be delayed for one day after treatment of annual weeds, or 7 days if perennial weeds are present to ensure absorption of Ken-Up 450 CT Non-Selective, Translocated Herbicide. Certain plants (eg. Soursob, variegated thistle) may be naturally toxic to stock. When known toxic plants are present, do not allow stock to graze until complete browning of treated plants has occurred.

Resistant Weeds Warning

GROUP M HERBICIDE

Ken-Up 450 CT Non-Selective, Translocated Herbicide is a member of the Glycines group of herbicides. Ken-Up 450 CT Non-Selective, Translocated Herbicide has the inhibition of EPSP Syntheses mode of action. For weed resistance management KEN-UP 450 CT Non Selective, Translocated Herbicide is a group M Herbicide. Some naturally-occurring weed biotypes resistant to Ken-Up 450 CT Non-Selective, Translocated Herbicide and other Group M herbicides may exist through normal genetic variability in any weed population. These resistant weeds will not be controlled by Ken-Up 450 CT Non-Selective, Translocated Herbicide or other Group M herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Kenso Corporation (M) Sdn. Bhd. accepts no liability for any losses that may result from the failure of Ken-Up 450 CT Non-Selective, Translocated Herbicide to control resistant weeds.

Crop Establishment

This product is recommended for control of emerged weeds prior to crop establishment. Suitable cultivation and/or sowing operations are required to produce the most satisfactory seedbed for crop germination and development, proceed as follows:

- Spray Ken-Up 450 CT Non-Selective, Translocated Herbicide early when the weeds are young.
- If weeds are few and the soil is friable, you can start seeding as early as one day after spraying.
- If the weed density is high, you should delay sowing until the dead weeds are sufficiently decayed, to ensure that crop emergence is not delayed as a result of dead green or decaying weeds being incorporated by cultivation or sowing. This is particularly important in cold or wet weather. Grazing will help to reduce the weed density, and you can assist weed decay by cultivation to leave trash on the surface.
- If seedbed conditions are marginal, be careful to seed at the correct depth. Do not use pre-emergent herbicides if their labels indicate that they may delay crop emergence.

Mixing

NOTE: Reduced results may occur if water containing soil is used, eg. water from ponds and unlined ditches; or if hard water containing calcium salts is used.

Ken-Up 450 CT Non-Selective, Translocated Herbicide is a non-volatile liquid that mixes readily with water. DO NOT mix or store or apply this product in galvanised steel, unlined steel containers or spray tanks, since a highly flammable gas mixture may be formed. So use only spray tanks and equipment made of plastic or plastic lined, fibreglass, rubber, aluminium, brass, copper, or stainless steel.

- Make sure the spray tank is completely free of previous spray residues.
- Half fill the spray tank with clean water. Where possible avoid using turbid water, or hard water containing calcium salts, as this may reduce weed control.
- If Ken-Up 450 CT Non-Selective, Translocated Herbicide is being used alone, go to step 5.
- If Ken-Up 450 CT Non-Selective, Translocated Herbicide is being used with insecticides, other herbicides or additives (see Tank Mixtures), add these products now according to their label directions. If ammonium sulphate (500g/L) is required, add it first using 2 L/100 L of spray volume.
- Add the required volume of Ken-Up 450 CT Non-Selective, Translocated Herbicide, and mix well. Mechanical agitators may cause excessive foaming and should not be used unless required by any tank mix partners.
- Add the rest of the water. Near the end, add Silken Penetrant, see below. Make sure the filling hose is submerged to avoid excessive foaming, and on completion remove it immediately to avoid back siphoning into the water supply. Use the tank mix promptly. And certainly within 5 days since a gradual loss of activity will occur.

Spray tanks, pumps, lines and nozzles should be thoroughly rinsed with clean water following application to prevent corrosion.

Surfactant Addition

The use of a Silken Penetrant may improve weed control where water rates are high or product rates low. Use 200mL of 1000g/L Silken Penetrant per 100L spray solution. Do not add any other agricultural chemicals, spraying oils or other materials except as directed on the label.

Application

Ken-Up 450 CT Non-Selective, Translocated Herbicide is a non-selective translocated herbicide. Direct spray contact or even slight drift may cause severe injury or destruction of any growing crop or other desirable plants including trees. Thoroughly flush your mixing and spray equipment with clean water after use.

Ground Application

- Apply 25-100 L of spray mix per hectare.
- Where possible use fan nozzle equipment with pressures of 240-280 kPa.
- Set the boom high enough to ensure double overlap of nozzle patterns at the top of the weed canopy.

Aerial Application

- Use aerial spraying only in pasture or fallow before establishing new pasture, field crops or fodder crops, or for pre-harvest application to Sorghum. **Do not apply Ken-Up 450 CT Non-Selective, Translocated Herbicide by air within intensive cropping areas as the consequences of accidental drift damage are too extreme.**
- Do not exceed 3.2 L of Ken-Up 450 CT Non-Selective, Translocated Herbicide per hectare.
- Use at least 15 L/ha with Micronair or boom equipment.
- On cool days apply at least 15 L of spray mix per hectare using an average droplet size (or VMD) of 250-350 microns and a swath 15-17 metres wide.
- On hilly terrain, increase the water volume to 30-80 L/ha and the minimum average droplet size to 300 micron.
- At 25°C increase the water volume to at least 30 L/ha and the minimum average droplet size to 300 micron to compensate for evaporation. Do not spray by air at temperatures above 35°C as excessive evaporation may occur that will reduce weed control.
- Drift is likely when droplet size is 150 microns or less, when wind speed is near zero or over 8 km/hr, the air is hot and dry, or there is a temperature inversion. Do not spray under these conditions.
- After each day of spraying, thoroughly wash the aircraft and landing gear with clean water to remove herbicide residues.

Drift Warning

DO NOT use when breeze is blowing towards nearby desirable plants. DO NOT apply under weather conditions, or from spraying equipment that may cause spray to drift onto nearby susceptible plants, crops, cropping lands or pastures.

Tank Mixtures — Compatibility

Ken-Up 450 CT Non-Selective, Translocated Herbicide may be tank-mixed with the following. Read and follow all label directions, restraints, plant-back periods, withholding periods, regional use restrictions and safety directions for the tank mix product:

Herbicides: Ken-Ester LV 680, Kenso Agcare Tri-Allate Selective Herbicide, Kenso Agcare Dicamba 500 Selective Herbicide, Ken-Chlor 750 WG, Ken-Met 600 WG, Ox 240, Kenso Agcare Atrazine 900 WG Herbicide (Do not apply the tank mix for control of Barnyard grass or Liverseed grass) plus ammonium sulphate, Kenso Agcare Simazine 900 WG Herbicide plus ammonium sulphate.

Insecticides: Dimethoate, Imidan*, Le-Mat* Kensban 500, Perfekthion* EC 400.

Additives: Ammonium Sulphate, Silken Penetrant as specified.

Crop /Situation	State	Weeds controlled	Rate Vol/ha	Critical comments
NORTHERN AUSTRALIA In fallows or prior to sowing a crop	QLD, NSW only	Annual phalaris (Canary grass), Barley grass, volunteer cereals, Wild oats	400 - 800 mL	Treat actively growing weeds not under stress from low moisture, frost, cold, disease or waterlogging. If heavy grazing has occurred allow regrowth to 6-8 cm. NOTE that under summer (hot) conditions, dense infestations of Barnyard grass and Liverseed grass may require follow-up treatment for complete control. In winter (cold) conditions, symptoms on Deadnettle may be slow to develop.
		Barnyard grass, Button grass, Columbus grass (seedling), Liverseed grass, Native Millet, Stinkgrass (Lovegrass), Volunteer Sorghum	800 mL –1.6 L	RATE SELECTION: Use the lower rate on young weeds, increase to the higher rate where grasses reach full tillering or where broadleaf weeds reach stem elongation/budding. At more advanced stages of growth certain broadleaf weeds require a higher rate range or the addition of Ken-Ester LV 680.
		Australian bluebell (QLD only), Cudweed, Fumitory, Mexican poppy, New Zealand spinach, Saffron thistle, Spear thistle, Spurge, Stinking goosefoot	800 mL - 1.2 L	CROP ESTABLISHMENT: Sowing should not proceed until conditions allow for a formation of a satisfactory seedbed. See Crop Establishment for directions.
		Black (giant) pigweed, Boggabri weed, Calltrop (Yellowvine), Indian hedge mustard, Mintweed, Summer grass	400 – 800 mL up to 5 true leaves or 3cm diameter/ height 800mL – 1.2 L greater than 3cm diameter/height	TANK MIXTURES: Read and follow all label directions, restraints, plant back periods, withholding periods, regional use restrictions and safety directions for the tank mix products. DO NOT tank mix with atrazine when spraying Barnyard grass or Liverseed grass.
		African turnip weed, Deadnettle, Sweet summer grass, Variegated thistle, Volunteer Sunflower	600 – 800 mL up to 5 true leaves or 3cm diameter/ height 800mL – 1.6 L greater than 3cm diameter/height	AERIAL APPLICATION: For instructions on Aerial application under hot conditions, see Aerial Application. DO NOT apply by aircraft when temperature is above 30°C.
		Annual ground cherry (Gooseberry), Bladder Ketmia, Camel melon, False castor oil plant/ Thornapple, Noogoora burr, Turnip weed, Wild lettuce, Wild turnip, Wireweed	800 mL – 1.2 L prior to stem elongation/ budding. After that use 400mL – 1.2 L plus 500 – 700 mL 2,4-D Ester (800g/L) OR 1.2 – 1.6 L of this product alone	
		Pigweed	800 mL - 1.6 L up to 20cm diameter	Use a higher rate on larger weeds. Control of Pigweed over a wide range of growth stages can be obtained with the addition of Ken-Met 600 WG. Observe recropping intervals.
		Sowthistle/ Milkthistle	600 – 800 mL rosette up to 3cm diameter 800mL – 1.6 L greater than 3cm diameter	Previously grazed plants may be difficult to control without allowing full recovery
		Couch	1.2 – 2.4 L	Use the higher rate for dense infestations. Apply sequential treatments during summer and autumn, with autumn being most effective. Repeat applications may be required for full control. For improved control use in conjunction with cultivation
		Johnson grass	1.5 – 2.4 L	Use the higher rates on plants approaching seedhead stage. Apply to plants with a minimum of 30 cm new growth. Sequential treatments will be required for long term control.
Nutgrass	2.4 + 2.4 L	Make first application to actively growing plants when at least 20% have reached the head stage (normally about February). After allowing maximum re-emergence to occur (normally in 6-8 weeks), it is essential to make a second application. NOTE: Follow-up treatments should be made as part of a Nutgrass control program.		

Crop/Situation	State	Weeds controlled	Rate Vol/ha	Critical comments
SORGHUM CONTROL Pre-harvest	QLD, NSW only	Sorghum, Grain Sorghum DO NOT apply to varieties intended for seed production or varieties prone to lodging	1.2 or 1.6 L	DO NOT apply if crop is under stress from low moisture, frost, cold or waterlogging. RATE SELECTION: Use the lower rate for control of crop and late tillers and suppression of ratoon regrowth. Use the higher rate for improved suppression of ratoon regrowth. TIMING: Apply when grain moisture is less than 25%. Application can be made when moderate browning has occurred. CAUTION: Treatment may increase potential for CROP LODGING, particularly if prior moisture stress has occurred. Harvest as soon as sufficient dry down has occurred to avoid possible lodging CAUTION: Sorghum may be naturally toxic to stock.
SORGHUM CONTROL Post-harvest	QLD, NSW only	Sorghum stubble, Grain sorghum	800 mL – 1.2 L for fresh regrowth from slashed stubble. 1.2 – 1.6 L for standing stubble if sufficiently green and for fresh spring regrowth	APPLY UNDER GOOD GROWING CONDITIONS ONLY. DO NOT apply if plants are under stress from low moisture, frost, cold or waterlogging. SLASHED STUBBLE AND SPRING REGROWTH: Apply when fresh regrowth is at least 20 cm high. STANDING STUBBLE: Apply only if sufficient green leaf is present. If grazing has occurred allow regrowth to 20 cm high before treatment. RATE SELECTION: Use the lower rate for knockdown and regrowth suppression where cultivation is to follow. Increase to the higher rate for improved regrowth control. NOTE: Variable results occur where the crop has been subject to stress or the growing conditions are marginal. CAUTION: Sorghum may be naturally toxic to stock
SUGAR CANE Ratoon spray out	QLD, NSW only	Sugar Cane ratoon regrowth	3.2 - 7.2 L	APPLY UNDER GOOD GROWING CONDITIONS ONLY to actively growing ratoons 60 – 120 cm tall. DO NOT apply if plants are under stress from low moisture, frost, cold or waterlogging. Use the lower rate for suppression or where cultivation is to follow. Use the higher rate for control.
RICE Direct drilling	NSW only	Annual phalaris (Canary grass), Annual ryegrass, Barley grass, Burr medic, Sub-clover, Winter grass	800 mL – 1.0 L	This product is less effective on drought stressed plants. In drought conditions a pre-watering prior to spraying is recommended. In grazed situations, if heavy grazing has occurred allow regrowth to 6-8 cm before spraying. ANNUAL RYEGRASS: Add Silken Penetrant at 200mL/100L spray solution and when dominant use the higher rate. SOWING: Direct drilling may take place 1 – 14 days after spraying. This product does not provide residual weed control. Permanent water and approved selective herbicides should be used to provide continual control of weeds.