

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

KENSO AGCARE

PAVO 250 EC FUNGICIDE

ACTIVE CONSTITUENT: 250 g/L PYRACLOSTROBIN
SOLVENT: 705 g/L LIQUID HYDROCARBONS

GROUP **11** FUNGICIDE

For the control of leaf speckle and leaf spot in bananas and downy and powdery mildew in grapevines, husk spot in macadamia and rust in almond as specified in the Directions for Use Table.

Kenso Corporation (M) Sdn Bhd
Level 1, 98 Commercial Road,
Teneriffe QLD 4005
Phone (07) 3216 1188
www.kenso.com.au



IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USE

CONTENTS: 5 Litres APVMA Approval No.: 83897/ 109210

KENSO AGCARE PAVO 250 EC FUNGICIDE

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight.

Triple-rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. When opening the container and preparing spray wear cotton overalls buttoned to the neck and wrist [or equivalent clothing], a washable hat, elbow-length PVC gloves and face shield or goggles. When using the prepared spray by ground application wear cotton overalls buttoned to the neck and wrist [or equivalent clothing], a washable hat and elbow-length PVC gloves. If product on skin, immediately wash area with soap and water. If product in eyes, wash it out immediately with water. Wash hands after use. After each days use, wash gloves, face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (TEL: 131126). If swallowed, DO NOT induce vomiting. Give a glass of water.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet, which can be obtained from www.kenso.com.au.

CONDITIONS OF SALE

Kenso Corporation (M) Sdn. Bhd. will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Kenso Corporation (M) Sdn. Bhd. is limited to the replacement of the goods or (at the option of Kenso Corporation (M) Sdn. Bhd.) the refund of the price paid and where possible sufficient part of the goods to enable proper examination being returned to Kenso Corporation (M) Sdn. Bhd. within thirty days of delivery.

In a Transport Emergency
Dial **000**
Police or Fire Brigade



Batch No.:

Date of Manufacture:

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

KENSO
agcare

PAVO 250 EC

Fungicide

ACTIVE CONSTITUENT: 250 g/L PYRACLOSTROBIN
SOLVENT: 705 g/L LIQUID HYDROCARBONS

GROUP 11 FUNGICIDE

For the control of leaf speckle and leaf spot in bananas and downy and powdery mildew in grapevines, husk spot in macadamia and rust in almond as specified in the Directions for Use Table.

READ THIS LEAFLET BEFORE USE

APVMA Approval No.: 83897/109210



Kenso Corporation (M) Sdn. Bhd.
Level 1, Commercial Road,
Teneriffe QLD 4005
Phone (07) 3216 1188

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet, which can be obtained from www.kenso.com.au.

CONDITIONS OF SALE

Kenso Corporation (M) Sdn. Bhd. will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Kenso Corporation (M) Sdn. Bhd. is limited to the replacement of the goods or (at the option of Kenso Corporation (M) Sdn. Bhd.) the refund of the price paid and where possible sufficient part of the goods to enable proper examination being returned to Kenso Corporation (M) Sdn. Bhd. within thirty days of delivery.

In a Transport Emergency
Dial **000**
Police of Fire Brigade



Kenso Corporation (M) Sdn. Bhd.
Level 1, 98 Commercial Road,
Teneriffe QLD 4005
Phone (07) 3216 1188

APVMA Approval No.: 83897/109210

DIRECTIONS FOR USE: RESTRAINTS

DO NOT use by aerial application in grapevines, almond and macadamia crops.
DO NOT apply within 48 hours of heavy rainfall. DO NOT irrigate to the point of run-off.
DO NOT apply to water-logged soil.

MANDATORY NO-SPRAY ZONES

DO NOT apply if there are aquatic and wetland areas including aquacultural ponds, surface streams and rivers downwind from the application area and within the mandatory no-spray zones shown in Table 1 below:

Table 1 – No-Spray Zones for Protection of the Aquatic	
FOR AERIAL APPLICATION	
Crop	Downwind Mandatory No-
Bananas	30 metres
FOR GROUND APPLICATION	
Crop	Downwind Mandatory No-
Bananas	10 metres
Grapes	10 metres
Macadamias	120 metres
Almonds	160 metres

CROP	DISEASE	RATE	WHP (Days)	CRITICAL COMMENTS
Bananas (bagged)	Leaf speckle (<i>Mycosphaerella musae</i>) Leaf spot (<i>Mycosphaerella musicola</i>) Black Sigatoka (<i>Mycosphaerella fijiensis</i>)	300 to 400 mL plus 3 to 5 L of water miscible oil/ha.	-	Maintain good crop hygiene by regularly removing old and diseased leaves to reduce the source of disease inoculum. DO NOT use on bananas unless bunch covers are in place. Ground application and aerial application: Apply by airblast sprayer or misting machine. Use the higher rates under conditions favouring disease infection. Apply up to four sprays per season, as part of a preventive disease control programme, commencing early in the season and using alternative mode-of-action products in between PAVO 250 EC Fungicide applications. Do NOT apply consecutive sprays of PAVO 250 EC Fungicide or other Group 11 fungicides. This use is subject to a Crop Life Australia resistance management strategy.
Grapevines	Downy mildew (<i>Plasmopara viticola</i>) Powdery mildew (<i>Uncinula necator</i>)	Dilute spray 40mL/100 L water Concentrate spray Refer to the "Application" section.	21	Also see 'CAUTION' section re export commodities. Apply up to three sprays per season as part of a complete disease control programme. Ideally, apply in a block of three sprays at 10 to 14 day intervals, commencing at flowering. Use the shorter intervals under conditions favouring disease infection. Apply by dilute or concentrate spraying equipment. Apply the same total amount of product to the target crop whether applying this product by dilute or concentrate spraying methods. This use is subject to a Crop Life Australia resistance management strategy.
Macadamia	Husk spot (<i>Pseudocercospora macadamiae</i>)	40mL/ 100L in maximum spray volume of 2000L/ha	-	Apply up to two sprays per season as part of a complete disease control program. Commence PAVO 250 EC Fungicide application at match head timing stage and repeat application at 14 to 28 days later. Ensure that fungicides from an alternative chemical group are included in the spray program each season.
Almond	Rust (<i>Tranzschelia discolor</i>)	40mL/ 100L	-	Apply up to two sprays per season as part of a complete disease control program. Commence application at flowering and repeat application at 10 to 14 days later. Ensure that fungicides from an alternative chemical group are included in the spray program each season.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS: HARVEST:

ALMOND, BANANAS (BAGGED), MACADAMIA - NOT REQUIRED WHEN USED AS DIRECTED.
GRAPES - DO NOT HARVEST FOR 21 DAYS AFTER APPLICATION

GENERAL INSTRUCTIONS

MIXING

To ensure even mixing, half-fill the spray tank with clean water and add the required amount of product. If required, add compatible products and agitate thoroughly, then add the remainder of the water. Agitate again before spraying commences.

APPLICATION

Bananas:

Ground application: Apply by airblast sprayer or misting machine to ensure even coverage. Aerial application: Apply in a minimum volume of 20 L/ha.

Grapevines/ Macadamia/ Almond:

Dilute Spraying

- Use a sprayer designed to apply high spray volumes, up to the point of run-off and matched to the crop being sprayed.
- Set up and operate the sprayer to achieve even coverage throughout the crop canopy. Apply sufficient spray solution to cover the crop to the point of run-off. Avoid excessive run-off.
- The required spray volume to achieve point of run off may be determined by applying different test volumes, using different settings on the sprayer, or from industry guidelines or other expert advice.
- Add the amount of product specified in the Directions for Use Table for each 100 L of water. Spray to the point of run-off.
- The required dilute spray volume to achieve point of run off will change and the sprayer set up and operation may also need to be changed, as the crop grows.

Concentrate Spraying

- Use a sprayer designed and set up for concentrate spraying (that is a sprayer which applies spray volumes less than those required to reach the point of run-off) and matched to the crop being sprayed.
- Set up and operate the sprayer to achieve even coverage throughout the crop canopy using your chosen spray volume.
- Determine an appropriate dilute spray volume (See Dilute Spraying above) for the crop canopy. This is needed to calculate the concentrate mixing rate.
- The mixing rate for concentrate spraying can then be calculated in the following way:

EXAMPLE ONLY

1. Dilute spray volume as determined above: For example 1500 L/ha
 2. Your chosen concentrate spray volume: For example 500 L/ha
 3. The concentration factor in this example is: $3 \times$ (i.e. $1500 \text{ L} \div 500 \text{ L} = 3$)
 4. If the dilute label rate is 20 mL/100 L, then the concentrate rate becomes 3×20 (that is 60 mL of product per 100L water for concentrate spraying).
- The chosen spray volume, amount of product per 100 L of water, and the sprayer set up and operation may need to be changed as the crop grows.
 - For further information on concentrate spraying, users are advised to consult relevant industry guidelines, undertake appropriate competency training and follow industry Best Practices.

COMPATIBILITY

For information on compatibility please contact your local Kenso Agcare representative.

GROUP 11 FUNGICIDE

FUNGICIDE RESISTANCE WARNING

PAVO 250 EC Fungicide is a member of the strobilurin group of fungicides. For fungicide resistance management, PAVO 250 EC Fungicide is a Group 11 fungicide. Some naturally-occurring individual fungi resistant to PAVO 250 EC Fungicide and other Group 11 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by PAVO 250 EC Fungicide or other Group 11 fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, Kenso Agcare Pty Ltd accepts no liability for any losses that may result from the failure of PAVO 250 EC Fungicide to control resistant fungi.

PRECAUTIONS

DO NOT use human flaggers/markers unless they are protected by engineering controls such as enclosed cabs.

RE-ENTRY PERIOD

Do NOT allow entry into treated areas until the spray has dried. When prior entry is necessary, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

CAUTION

EXPORT OF TREATED FRUIT OR WINE

Growers should note that Maximum Residue Limits (MRLs) or import tolerances do not exist in all markets for fruit treated with PAVO 250 EC Fungicide. Additionally, some export markets have established MRLs different to those in Australia. If you are growing fruit for export (either fresh, dried or for wine production), please check with Kenso Agcare Pty Ltd or the Australian Wine Research Institute <http://www.awri.com.au> for the latest information on MRLs and import tolerances BEFORE using PAVO 250 EC Fungicide.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

HIGHLY TOXIC TO AQUATIC ORGANISMS. Do NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple-rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. When opening the container and preparing spray wear cotton overalls buttoned to the neck and wrist [or equivalent clothing], a washable hat, elbow-length PVC gloves and face shield or goggles. When using the prepared spray by ground application wear cotton overalls buttoned to the neck and wrist [or equivalent clothing], a washable hat and elbow-length PVC gloves. If product on skin, immediately wash area with soap and water. If product in eyes, wash it out immediately with water. Wash hands after use. After each days use, wash gloves, face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126. If swallowed, do NOT induce vomiting. Give a glass of water.