

POISON

KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

KENSO AGCARE

PIRI-KEN 500 WG APHICIDE

ACTIVE CONSTITUENT: 500 g/kg PIRIMICARB (an anti-cholinesterase compound)

GROUP 1A INSECTICIDE

For control of certain Aphids on crops and pastures as per directions for use table.

Kenso Corporation (M) Sdn Bhd
Level 1, 98 Commercial Road,
Teneriffe QLD 4005
Phone (07) 3216 1188
www.kenso.com.au



IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USE

CONTENTS: 5 kilograms APVMA Approval No.: 66473/ 115545

KENSO AGCARE PIRI-KEN 500 WG APHICIDE

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Do not store near food, feedstuffs, fertilisers or seeds. Do not dispose of undiluted chemicals on-site. When the container is empty, shake any residual material into the spray tank. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver to an approved waste management facility. If an approved waste management facility is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots in compliance with relevant State, Territory or Local government regulations. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Poisonous if absorbed by skin contact or inhaled or swallowed. May irritate the nose and throat and will irritate the eyes. Avoid contact with eyes and skin. Do not inhale dust or spray mist. When opening the container and preparing product for use wear cotton overall, over normal clothing, buttoned to the neck and wrist and a washable hat, elbow length chemical resistant gloves, goggles and half face piece respirator with dust cartridge or canister. If applying by boomspray equipment wear cotton overalls buttoned to neck and wrist (or equivalent clothing) and elbow length chemical resistant gloves. If applying by low pressure hand wand or high pressure hand wand, wear cotton overalls, over normal clothing, buttoned to neck and wrist and a washable hat, elbow length chemical resistant gloves, goggles and a half face piece respirator with dust cartridge or canister. If clothing becomes contaminated with product, or wet with spray remove clothing immediately. If product or spray on skin, immediately wash area with soap and water. If product or spray in eyes wash it out immediately with water. Wash hands after use. After each day's use, wash gloves, goggles, respirator (if rubber wash with detergent and warm water) and contaminated clothing.

FIRST AID

If swallowed, splashed on skin or in eyes, or inhaled, contact the Poisons Information Centre (Phone Australia 13 11 26) or a doctor at once. Remove any contaminated clothing and wash skin thoroughly. If swallowed, activated charcoal may be advised. Give atropine if instructed.

SAFETY DATA SHEET

For further Information refer to the Safety Data Sheet (SDS) which can be obtained from www.kenso.com.au.

CONDITIONS OF SALE

Kenso Corporation (M) Sdn. Bhd. will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply of these goods other

than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Kenso Corporation (M) Sdn. Bhd. is limited to the replacement of the goods or (at the option of Kenso Corporation (M) Sdn. Bhd.) the refund of the price paid and where possible sufficient part of the goods to enable proper examination being returned to Kenso Corporation (M) Sdn. Bhd. within thirty days of delivery.

In a Transport Emergency
Dial **000** Police or Fire Brigade



UN NO. 2757
CARBAMATE
PESTICIDE,
SOLID, TOXIC

PG III HAZCHEM 2X



Additional statements required by Globally Harmonised System of Classification and Labelling of Chemicals (GHS) and Safe Work Australia: **TOXIC IF SWALLOWED. HARMFUL IN CONTACT WITH SKIN. HARMFUL IF INHALED.** *Precautionary:* Avoid breathing dust/fume/gas/mist/vapours/spray. Wash contacted areas thoroughly after handling. Do not eat, drink or smoke when using this product. Use only outdoors or in a well ventilated area. Wear protective gloves, protective clothing and eye or face protection. IF SWALLOWED: immediately call a POISON CENTER or doctor. IF ON SKIN: Wash with plenty of soap and water. IF INHALED: Remove victim to fresh air and keep at rest in a position comfortable for breathing. Call a POISON CENTER or doctor/physician if you feel unwell. Specific treatment (see FIRST AID on this label). Rinse mouth. Wash contaminated clothing before reuse. Store locked up.

Batch No.:

Date of Manufacture:

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

KENSO
agcare

PIRI-KEN
500 WG

Aphicide

ACTIVE CONSTITUENT: 500 g/kg PIRIMICARB (an anti-cholinesterase compound)

GROUP 1A INSECTICIDE

For control of certain Aphids on crops and pastures as per directions for use table.

IMPORTANT: READ THE LEAFLET BEFORE USE

APVMA Approval No.: 66473/115545



Kenso Corporation (M) Sdn. Bhd.
Level 1, 98 Commercial Road,
Teneriffe QLD 4005
Phone (07) 3216 1188

SAFETY DIRECTIONS

Poisonous if absorbed by skin contact or inhaled or swallowed. May irritate the nose and throat and will irritate the eyes. Avoid contact with eyes and skin. Do not inhale dust or spray mist. When opening the container and preparing product for use wear cotton overall, over normal clothing, buttoned to the neck and wrist and a washable hat, elbow length chemical resistant gloves, goggles and half facepiece respirator with dust cartridge or canister. If applying by boomspray equipment wear cotton overalls buttoned to neck and wrist (or equivalent clothing) and elbow length chemical resistant gloves. If applying by low pressure hand wand or high pressure hand wand, wear cotton overalls, over normal clothing, buttoned to the neck and wrist and a washable hat, elbow length chemical resistant gloves, goggles and a half piece respirator with dust cartridge or canister. If clothing becomes contaminated with product, or wet with spray remove clothing immediately. If product or spray on skin, immediately wash area with soap and water. If product or spray in eyes wash it out immediately with water. Wash hands after use. After each day's use, wash gloves, goggles, respirator (if rubber wash with detergent and warm water) and contaminated clothing.

FIRST AID

If swallowed, splashed on skin or in eyes, or inhaled, contact the Poisons Information Centre (Phone Australia 13 11 26) or a doctor at once. Remove any contaminated clothing and wash skin thoroughly. If swallowed, activated charcoal may be advised. Give atropine if instructed.

SAFETY DATA SHEET

For further Information refer to the Safety Data Sheet (SDS) which can be obtained from www.kenso.com.au.

CONDITIONS OF SALE

Kenso Corporation (M) Sdn. Bhd. will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Kenso Corporation (M) Sdn. Bhd. is limited to the replacement of the goods or (at the option of Kenso Corporation (M) Sdn. Bhd.) the refund of the price paid and where possible sufficient part of the goods to enable proper examination being returned to Kenso Corporation (M) Sdn. Bhd. within thirty days of delivery.

* Not a registered trademark of Kenso Corporation (M) Sdn. Bhd.

In a Transport Emergency
Dial **000**
Police of Fire Brigade



Kenso Corporation (M) Sdn. Bhd.
Level 1, 98 Commercial Road,
Teneriffe QLD 4005
Phone (07) 3216 1188

APVMA Approval No.: 66473/115545

DIRECTIONS FOR USE

Restrains:

DO NOT apply by equipment carried on the back of the user.
DO NOT apply from the air when the crosswind is less than 3km/hr.

1. FRUIT CROPS - TREE AND VINE CROPS

Rate	Critical Comments
In the following table: "Fruit Crops - Tree and Vine Crops" all rates given are for dilute spraying. For concentrate spraying, refer to the mixing/Application Section.	For all uses in the table "Fruit Crops - Tree and Vine Crops": Apply by dilute or concentrate spraying equipment. Apply the same total amount of product to the target crop whether applying this product by dilute or concentrate spraying methods.

CROPS	PEST	STATE	RATE		WHP	CRITICAL COMMENTS
			PER 100L	PER HA		
Apples	Woolly Aphid	NSW, ACT, VIC, TAS, SA, WA only	50g	1.1-1.7kg	2 days	Use the higher rate when less than 2200L of spray per ha. Add 18mL Wetspray 600 per 100L.
Citrus	Citrus Aphid	NSW, ACT, VIC, TAS, SA, WA only	50g	1.1kg	2 days	Apply when aphids appear. Use at least 2200L spray per ha.
	Aphids	Qld, ACT, WA only	2.5g	-		Apply only where pest management is being implemented and only when it is apparent that parasite predators will not control the infestation within a reasonable time.
Stone fruit	Green Peach Aphid, Black Peach Aphid, Cherry Aphid	ALL STATES	50g	1.1kg		Apply at pink bud stage or when aphids appear. Use at least 1100L spray/ha.

2. FRUIT CROPS - NON TREE AND VINE CROPS

CROPS	PEST	STATE	RATE		WHP	CRITICAL COMMENTS
			PER 100L	PER HA		
Blueberries	Aphids	QLD, WA only	50g	-	2 days	Use the higher rate when less than 2200L of spray per ha. Add 18mL Wetspray 600 per 100L.
Pepinos		QLD, NSW, WA only	35g			Thorough coverage is necessary. If used in conjunction with predatory mites, apply before release or only after mites have fully established within the crop.
Strawberries						

3. VEGETABLES

CROPS	PEST	STATE	RATE		WHP	CRITICAL COMMENTS
			PER 100L	PER HA		
Artichokes (globe)	Aphids	ALL STATES	50g	-	2 days	-
Asparagus						
Beetroot	Green Peach Aphid		-	1kg		Use at least 500L spray/ha plus 18mL/100L Wetspray 600.
Beans	Cowpea Aphid (<i>Aphis craccivora</i>)	Vic, Tas, WA only		500g		Apply as beans emerge and thereafter whilst aphid attack continues. Spray thoroughly.
Broad, Beans, Lima Beans	Aphids	ALL STATES	50g	-		
Broccoli, Brussels Sprouts, Cabbages, Cauliflowers	Cabbage Aphid	ALL STATES	100 to 200g	500g or 1kg		Apply every 10 to 14 days commencing 1week after planting. Use 500L spray per ha. Add 18mL Wetspray 600 per 100L. If lower spray volumes per ha are used, increase the rate per 100L to maintain the equivalent rate of product per ha. Use the higher rate if temperature falls below 20°C.
Brussels Sprouts, Cabbages, Cauliflowers	Green Peach Aphid	ALL STATES	100 to 200g	500g or 1kg		
Chinese Cabbages	Cabbage Aphid, Green Peach Aphid	ALL STATES				
Capsicums	Aphids	ALL STATES				Spray when aphids are detected and repeat if necessary. Use the higher rates if applied in conditions below 20°C.
Cucurbits						Apply every 5 to 10 days during periods of rapid plant growth to delay virus buildup. Use the higher rates if applied in conditions below 20°C.

CROPS	PEST	STATE	RATE		WHP	CRITICAL COMMENTS
			PER 100L	PER HA		
Endive, Garden Cress	Aphids	ALL STATES	50g	-	2 days	-
Kale	Cabbage Aphid, Green Peach Aphid	ALL STATES	100g to 200g	500g or 1kg		Apply every 10 to 14 days commencing 1 week after planting. Use 500 litres spray per ha. Add 18mL Wetspray 600 per 100L. If lower spray volumes per ha are used, increase the rate per 100L to maintain the equivalent rate of product per ha. Use the higher rate if temperature falls below 20°C.
Kiwano	Aphids	ALL STATES	50g	-		Apply every 5 to 10 days during period of rapid plant growth to delay virus buildup.
Leeks			50g	-		-
Lettuce		ALL STATES	100 to 200g	500g or 1kg		Spray when aphids are detected and repeat if necessary. Use the higher rates if applied in conditions below 20°C.
Okra		ALL STATES	50g	-		For use in IPM programs where Helicoverpa resistance strategy is being practiced.
Peas		ALL STATES	100 to 200g	500g or 1kg		-
		NSW only				Spray when aphids are detected and repeat if necessary. Use the higher rates if applied in conditions below 20°C.
Potatoes	Potato Aphid	ALL STATES	250 to 500g	500g or 1kg		Apply in 200L spray per ha when aphids first appear. If higher spray volumes per ha are used, decrease the rate per 100L to maintain the equivalent rate of product per ha. Use the higher rate if temperature falls below 20°C.
Radishes	Cabbage Aphid, Green Peach Aphid	ALL STATES	100 to 200g	500g or 1kg		Apply every 10 to 14 days commencing 1week after planting. Use 500L spray per ha. Add 18mL Wetspray 600 per 100L. If lower spray volumes per ha are used, increase the rate per 100L to maintain the equivalent rate of product per ha. Use the higher rate if temperature falls below 20°C.
			50g	-		-
Shallots, Spinach	Aphids		50g	-		-
Swedes, Turnips	Cabbage Aphid, Green Peach Aphid	ALL STATES	100 to 200g	500g or 1kg		Apply every 10 to 14 days commencing 1 week after planting. Use 500L spray per ha. Add 18mL Wetspray 600 per 100L. If lower spray volumes per ha are used, increase the rate per 100L to maintain the equivalent rate of product per ha. Use the higher rate if temperature falls below 20°C.
Tomatoes	Aphids	ALL STATES				Spray when aphids are detected and repeat if necessary. Use the higher rates if applied in conditions below 20°C.
Watercress	Aphids	ALL STATES	50g	-		-

4. ORNAMENTALS

CROPS	INSECT	STATE	RATE		WHP	CRITICAL COMMENTS
			PER 100L	PER HA		
Ornamen-tals	Aphids	NSW, WA only	50g	-	-	Apply when aphids appear. Spray to give complete coverage of plants.
	Aphids	QLD, WA only	35g			If applied in conjunction with predatory mites (<i>Phytoseiulus persimilis</i>) apply before release or only after mites have become fully established. DO NOT apply more than 2 sprays in quick succession where IPM is practiced. Where roses are grown in dense hedges and IPM is used, spray only the upper half of the hedge where the aphids are attacking the buds. Allow the lower half of the hedge to serve as a refuge for <i>P. persimilis</i> . In such situations Kenso Agcare Piri-Ken 500 WG Aphicide should NOT be mixed with a fungicide which requires very careful coverage. Note: Because of the wide range of crops and pests encompassed under Ornamentals, the phytotoxicity and efficacy of this compound cannot be guaranteed under all uses. Test the chemical for phytotoxicity on a small area of crop before widespread usage.

CROPS	INSECT	STATE	RATE		WHP	CRITICAL COMMENTS
			PER 100L	PER HA		
Roses, Chrysanthemum	Rose Aphid, Chrysanthemum Aphid	ALL STATES	50g	-	-	Apply when aphids appear. Spray to give complete coverage of plants.

5. BROADACRE CROPS

CROPS	INSECT	STATE	RATE		WHP	CRITICAL COMMENTS	
			PER 100L	PER HA			
Canola	Cabbage Aphid (<i>Brevicoryne brassicae</i>), Green Peach Aphid (<i>Myzus persicae</i>)	ALL STATES	-	500g or 1kg	14 days harvest/ grazing	Apply when aphids first appear using 30 to 100L water/ha (ground rig) or 20 to 50L/ha (aerial). Use higher rate if temperature falls below 20°C. Add 18mL Wetspray 600 per 100L spray.	
Cotton	Cotton Aphid (<i>Aphis gossypii</i>), Green Peach Aphid (<i>Myzus persicae</i>)	QLD, NSW, WA ONLY	-	500g or 750g Aerial Application Spray volume of 30 to 50L/ha Ground Application Spray volume of 100L/ha minimum	21 days harvest/ grazing	Thorough spray coverage is essential to produce best results. Stage I - apply when aphids exceed 90% plants infested. Stage II - apply when aphids exceed 95% plants infested. Stage III- apply when aphids exceed 50% plants infested, or - if 1 % bolls are open and honeydew is present apply when aphids exceed 10% of plants infested. If a spray is required for another pest, the aphid threshold for each stage should be reduced by half to determine the need to spray for aphids; eg. 50% becomes 25% plants infested. Use Kenso Agcare Piri-Ken 500 WG Aphicide no more than twice per season. Adhere to the resistance management strategy and rotate chemistry. DO NOT use Kenso Agcare Piri-Ken 500 WG Aphicide as consecutive sprays.	
Duboisia	Aphids	QLD, WA only	50g	1.1kg	-	Apply when populations warrant.	
Lucerne, Medic pastures	Spotted Alfalfa Aphid, Blue Green Aphid	ALL STATES	-	100 to 150g in 200 to 300L water	3 days grazing	Application: High rate to be used when there is a high aphid population density or with dense crop growth. Complete coverage and penetration of the crop is essential. Timing: QLD only – On seedling lucerne, treatment is necessary when one or two aphids are present per plant. Established lucerne and medics (Spring and Summer) – spray when there are 20 to 40 aphids per stem. Spotted Alfalfa Aphid (NSW, SA, TAS, WA only): Apply when aphids begin to build up on stems and repeat as necessary. Spotted Alfalfa Aphid and Pea Aphid (VIC only). Seedling lucerne and medics – treat when 1 to 2 aphids per plant are present. Established lucerne and medics (Spring and Summer) – spray when there are 20 to 40 aphids per stem. TAS only – A cold weather adapted species which may be found at most times of the year. Spraying should be carried out when aphids begin to increase and be repeated as necessary. For the control of both the above lucerne and medic pasture pests Kenso Agcare Piri-Ken 500 WG Aphicide at the recommended rate should be applied in 200 to 300L of water per ha. Complete coverage and penetration of the crop is essential. Blue Green Aphid This aphid is cooler weather tolerant and will be found at all times of the year. Spraying should be carried out as aphids are observed. In some areas Spotted Alfalfa Aphid has developed tolerance to pirimicarb and in those areas Kenso Agcare Piri-Ken 500 WG Aphicide will not give satisfactory control. However, apparent tolerance should not be confused with poor application. Consult your local Department of Agriculture representative for local information.	
Lucerne	Pea Aphid	VIC, WA only	-	100 to 150g	3 days grazing	Apply when aphids begin to build up on stems. Repeat as necessary. Use 200 to 300L per ha. Use the higher rate when there is a high aphid population density or dense crop or when temperature falls below 20°C.	
Lupins	Green Peach Aphid	NSW, VIC, WA only	-	250g	6 weeks harvest/ grazing	Apply when aphids first appear using 20 to 100L/ha (ground rig) or 4 to 15 L/ha (aerial). Repeat as necessary. If low relative humidity conditions prevail add 1% or 10% Summer Spray Oil (DC-Tron or Caltex) to spray solution for ground and aerial Application respectively. Note: For best coverage and hence best results with aerial application; apply when a slight crosswind with a minimum speed of 3km/hr is present.	
	Cowpea Aphid	NSW, VIC, WA only				300g	Apply when aphids first appear. Good coverage of the infested area is essential. Apply in 30 to 100L/ha for aerial application. Repeat as necessary. If low relative humidity conditions prevail add Summer Spray Oil to the mix.
	Green Peach Aphid	WA only					
Winter Cereals	Aphids (<i>Rhopalosiphum maidis</i>)	ALL STATES	-	150g	6 weeks harvest/ grazing	Apply when 20 or more aphids are present per tiller and when the crop is between emergence of last leaf and flowering stages. Use 20 to 100L/ha (ground rig) or 4 to 15L/ha (aerial). If low relative humidity conditions prevail add Summer Spray Oil as for lupins. Where there is concern for infection by Barley Yellow Dwarf virus, spraying should be carried out as soon as aphids are located in the crop.	
	Aphids (<i>R. maidis</i> , <i>R. padi</i>)	WA only		250g to 300g			

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS:

FRUIT, VEGETABLES: DO NOT HARVEST FOR 2 DAYS AFTER APPLICATION.

CANOLA: DO NOT HARVEST, GRAZE OR CUT FOR STOCK FOOD FOR 14 DAYS AFTER APPLICATION.

COTTON: DO NOT HARVEST, GRAZE OR CUT FOR STOCK FOOD FOR 21 DAYS AFTER APPLICATION.

LUCERNE, MEDIC PASTURES: DO NOT GRAZE, CUT FOR STOCK FOOD OR FEED TO POULTRY FOR 3 DAYS AFTER APPLICATION.

LUPINS, WINTER CEREALS: DO NOT HARVEST, GRAZE OR CUT FOR STOCK FOOD OR FEED TO POULTRY FOR 6 WEEKS AFTER APPLICATION.

GENERAL INSTRUCTIONS

MIXING

Fill tank 50 to 75% full then start agitation. Slowly pour granules into tank and maintain agitation throughout spraying. DO NOT pre-mix Kenso Agcare Piri-Ken 500 WG Aphicide granules.

APPLICATION

If this product is applied when the temperature is less than 15°C it is essential to have thorough spray coverage for quick and total aphid control.

FRUIT CROPS - TREE AND VINE CROPS

Dilute Spraying

Use a sprayer designed to apply high volumes of water up to the point of runoff and matched to the crop being sprayed. Set up and operate the sprayer to achieve even coverage throughout the crop canopy. Apply sufficient water to cover the crop to the point of run-off. Avoid excessive run-off. The required water volume may be determined by applying different test volumes, using different settings on the sprayer, from industry guidelines or expert advice. Add the amount of product specified in the Directions for Use table for each 100L of water. Spray to the point of run-off. The required dilute spray volume will change and the sprayer set up and operation may also need to be changed, as the crop grows.

Concentrate Spraying

Use a sprayer designed and set up for concentrate spraying (that is a sprayer which applies water volumes less than those required to reach the point of runoff) and matched to the crop being sprayed. Set up and operate the sprayer to achieve even coverage throughout the crop canopy using your chosen water volume. Determine an appropriate dilute spray volume (See Dilute Spraying above) for the crop canopy. This is needed to calculate the concentrate mixing rate. The mixing rate for concentrate spraying can then be calculated in the following way:

Example Only:

1. Dilute spray volume as determined above: For example 1500L/ha.
2. Your chosen concentrate spray volume: For example 500L/ha.
3. The concentration factor in this example is: 3 x (ie 1500L ÷ 500L = 3)
4. If the dilute label rate is 10mL/100L, then the concentrate rate becomes 3 x 10, that is 30mL/100L of concentrate spray.

The chosen spray volume, amount of product per 100L of water, and the sprayer set up and operation may need to be changed as the crop grows.

For further information on concentrate spraying, users are advised to consult relevant industry guidelines, undertake appropriate competency training and following industry Best Practices.

COMPATIBILITY

Kenso Agcare Piri-Ken 500 WG Aphicide may be mixed in the spray vat with anyone of the following products: Wetspray* and Omite*. A mixture of Kenso Agcare Piri-Ken 500 WG Aphicide with more than one of these products or with any other product may be ineffective or may cause serious damage. The use of such a mixture is not recommended and would therefore be entirely at the user's risk.

GROUP 1A INSECTICIDE

INSECTICIDE RESISTANCE WARNING

For insecticide resistance management Kenso Agcare Piri-Ken 500 WG Aphicide is a Group 1A insecticide. Some naturally occurring insect biotypes resistant to Kenso Agcare Piri-Ken 500 WG Aphicide and other Group 1A insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if Kenso Agcare Piri-Ken 500 WG Aphicide or other Group 1A insecticides are used repeatedly. The effectiveness of Kenso Agcare Piri-Ken 500 WG Aphicide on resistant individuals could be significantly reduced.

Since occurrence of resistant individuals is difficult to detect prior to use, Kenso Corporation (M) Sdn Bhd accepts no liability for any losses that may result from the failure of Kenso Agcare Piri-Ken 500 WG Aphicide Aphicide to control insects.

Kenso Agcare Piri-Ken 500 WG Aphicide may be subject to specific resistance management strategies. For further information contact your local supplier, Kenso representative or local agricultural department agronomist.

Resistance Management for Green Peach Aphid control

Some Green Peach Aphids are resistant to all commonly used insecticides including Kenso Agcare Piri-Ken 500 WG Aphicide. If resistance is known or suspected, DO NOT use Kenso Agcare Piri-Ken 500 WG Aphicide and seek advice on resistance management. If there is no Kenso Agcare Piri-Ken 500 WG Aphicide resistance, use a maximum of 2 non-consecutive aphid sprays per season. If Kenso Agcare Piri-Ken 500 WG Aphicide was used as the last aphid spray for the season, do not use it as the first aphid spray next season.

PROTECTION OF LIVESTOCK

Animals and poultry should be kept away from orchards and other treated areas during the horticultural spraying season. Keep animals and poultry away from spray and treated areas for a minimum of 3 days. (refer to Withholding Periods)

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

DO NOT apply under meteorological conditions (eg. crosswinds less than 3km/hr) or from spraying equipment which could be expected to cause spray to drift onto adjacent areas, particularly wetlands, waterbodies or watercourses.

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Do not store near food, feedstuffs, fertilisers or seeds. Do not dispose of undiluted chemicals on-site. When the container is empty, shake any residual material into the spray tank. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver to an approved waste management facility. If an approved waste management facility is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots in compliance with relevant State, Territory or Local government regulations. Empty containers and product should not be burnt.