

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

KENSO AGCARE

REBELLION

800

HERBICIDE

ACTIVE CONSTITUENT: 800 g/L PROSULFOCARB

GROUP J HERBICIDE

For the control of Annual Ryegrass (*Lolium rigidum*) and other grass and broadleaf weeds in Barley and Wheat as per the directions for use.



Kenso Corporation (M) Sdn Bhd
Level 1, 98 Commercial Road,
Teneriffe QLD 4005
Phone (07) 3216 1188
www.kenso.com.au

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USE

CONTENTS: 20 Litres APVMA Approval No.: 82377/ 112737

KENSO AGCARE REBELLION 800 HERBICIDE

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight.

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Will irritate the eyes and skin. Repeated exposure may cause allergic disorders. Sensitive workers should use protective clothing. Avoid contact with eyes and skin. When opening the container, preparing the product for use and using the product, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), elbow-length chemical-resistant gloves and face shield or goggles. In addition, when opening the container and mixing and loading, wear a disposable mist face mask covering mouth and nose. If product on skin, immediately wash area with soap and water. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves and face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone 131 126.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet.

CONDITIONS OF SALE

Kenso Corporation (M) Sdn. Bhd. will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Kenso Corporation (M) Sdn. Bhd. is limited to the replacement of the goods or (at the option of Kenso Corporation (M) Sdn. Bhd.) the refund of the price paid and where possible sufficient part of the goods to enable proper examination being returned to Kenso Corporation (M) Sdn. Bhd. within thirty days of delivery.

In a Transport Emergency
Dial **000**
Police or Fire Brigade



Batch No.:

Date of Manufacture:

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



REBELION 800

Herbicide

**ACTIVE CONSTITUENT: 800 g/L
 PROSULFOCARB**

GROUP J HERBICIDE

For the control of Annual Ryegrass (*Lolium rigidum*) and other grass and broadleaf weeds in Barley and Wheat as per the directions for use.

READ THIS LEAFLET BEFORE USE

APVMA Approval No.: 82377/112737



Kenso Corporation (M) Sdn. Bhd.
 Level 1, 98 Commercial Road,
 Teneriffe QLD 4005
 Phone (07) 3216 1188

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

HIGHLY TOXIC TO AQUATIC ORGANISMS. DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

DO NOT apply under meteorological conditions or from spraying equipment which could be expected to cause spray to drift onto adjacent areas, particularly wetlands, water bodies or watercourses.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight.

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Will irritate the eyes and skin. Repeated exposure may cause allergic disorders. Sensitive workers should use protective clothing. Avoid contact with eyes and skin. When opening the container, preparing the product for use and using the product, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), elbow-length chemical-resistant gloves and face shield or goggles. In addition, when opening the container and mixing and loading, wear a disposable mist face mask covering mouth and nose. If product on skin, immediately wash area with soap and water. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves and face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone 131 126.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet.

CONDITIONS OF SALE

Kenso Corporation (M) Sdn. Bhd. will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Kenso Corporation (M) Sdn. Bhd. is limited to the replacement of the goods or (at the option of Kenso Corporation (M) Sdn. Bhd.) the refund of the price paid and where possible sufficient part of the goods to enable proper examination being returned to Kenso Corporation (M) Sdn. Bhd. within thirty days of delivery.

In a Transport Emergency
 Dial **000**
 Police of Fire Brigade



Kenso Corporation (M) Sdn. Bhd.
 Level 1, 98 Commercial Road,
 Teneriffe QLD 4005
 Phone (07) 3216 1188

APVMA Approval No.: 82377/112737

DIRECTIONS FOR USE

Restraints:

DO NOT apply to waterlogged soils.
 DO NOT apply if heavy rains or storms that are likely to cause runoff are forecast within 2 days of application.

Mandatory No-Spray Zones

DO NOT apply if there are aquatic and wetland areas including aquacultural ponds, within 20 metres downwind from the application area.
 DO NOT apply if there is non-target vegetation within 10 metres downwind from the application area.

Crop	Weeds	Rate/ha	Critical Comments
Barley Wheat	Annual Ryegrass (<i>Lolium rigidum</i>) including control of Group D resistant populations	2.5 L	Apply pre-emergent and incorporate mechanically by sowing operation (IBS). Application should be made to a moist seedbed up to 7 days prior to sowing and sufficient rain to thoroughly wet the top 3 to 4 cm of soil should occur within 10 days after spraying. Avoid soil throw into adjacent seeding rows or sites where furrow walls may collapse. Refer to crop tolerance, incorporation and tillage requirements under General Instructions.

Crop	Weeds	Rate/ha	Critical Comments
Barley Wheat	Annual Ryegrass (<i>Lolium rigidum</i>), Red and White Fumitory, Paradoxa Grass (Canary Grass), Phalaris spp, Sand Fescue, soil surface Wild Oats, Wireweed (Hogweed) Suppression of Brome Grass, Deadnettle, Rough Poppy, Yellow Burr Weed	1.5 to 2.5 L <i>plus</i> 0.8 to 1.5 L of Kenso Agcare Trifluralin 480 Selective Herbicide	DO NOT use less than 2.5 L/ha REBELION 800 HERBICIDE where Group D resistance is confirmed or suspected. Use of REBELION 800 HERBICIDE below 2.5 L/ha alone or in a tank mix WILL NOT be effective in the control of Group D resistant Annual Ryegrass nor to delay the onset of herbicide resistance development. Refer to Tank Mixes under General Instructions. Apply pre-emergent and incorporate mechanically by sowing operation (IBS). Application should be made to a moist seedbed up to 24 hours prior to sowing and sufficient rain to thoroughly wet the top 3 to 4 cm of soil should occur within 10 days after spraying. Avoid soil throw into adjacent seeding rows or sites where furrow walls may collapse. Use 1.5 L/ha of Kenso Agcare Trifluralin 480 Selective Herbicide in minimum tillage knife/blade point systems only as per trifluralin label. Attention to sowing speed and soil throw is important to ensure crop safety. This is especially critical at higher use rates. Risk of crop injury may increase where greater than 1 L/ha Kenso Agcare Trifluralin 480 Selective Herbicide is applied with 2.5 L/ha REBELION 800 HERBICIDE. Always refer to use recommendations on trifluralin label regarding soil type restrictions and incorporation timing and rate selection under different tillage systems. This mixture is recommended for control or suppression of additional weeds, on sandy soils where product movement into the seeding zone poses a crop safety risk or where good soil moisture cannot be assured. Refer to crop tolerance, incorporation and tillage requirements under General Instructions.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHOLDING PERIODS:

Barley, Wheat

Harvest: NOT REQUIRED WHEN USED AS DIRECTED

Grazing: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 10 WEEKS AFTER APPLICATION

GENERAL INSTRUCTIONS

REBELION 800 HERBICIDE is a short residual, soil applied, pre-emergent herbicide that is absorbed by the roots and shoots (coleoptile) of germinating seedlings with inhibition of growth in the meristematic region. Foliar uptake is possible but of lower effect. Upstream translocation in the plant occurs but movement in the phloem is very unlikely. Best activity can be expected from applications before or shortly after the germination of target weeds.

The product should be applied to a moist smooth seedbed free of soil clods and emerged weeds. Product efficacy may be reduced by improper incorporation, high levels of crop or trash residues, stones or foreign matter and in areas of extremely high weed seed density such as header rows. For optimal efficacy and crop safety avoid shallow seeding and ensure accurate seed placement that avoids placement of seed in the herbicide band. Avoid water repellent soils, soils subject to water logging or where furrow walls are likely to collapse. Duration and effectiveness of weed control will depend on use rate, soil type and rainfall or irrigation after application. Adequate rainfall should occur within 10 days of application, as this herbicide requires sufficient available soil moisture to ensure soil movement and uptake by emerging weed seeds.

Crop Tolerance

The selectivity of REBELION 800 HERBICIDE in cereals is a combination of positional and physiological selectivity. Positional selectivity must be maintained by sowing at adequate depth below the herbicide band. The physiological selectivity of REBELION 800 HERBICIDE is given by the ability of cereals to metabolise the herbicide to inactive compounds more rapidly than susceptible weed species. Environmental factors such as rainfall events following application and soil type will influence product movement into the seed zone. Crop damage may result where tank mixed with other herbicides, heavy rainfall follows sowing, in the event of crop stresses including frost or in sandy soils with high leaching potential. Risk of crop damage may be exacerbated where varieties with short coleoptile length are planted in conjunction with the use of some fungicide seed treatments.

Tillage System

IBS with knife or blade points: Working speed should ensure adequate incorporation of product but avoid soil throw into the adjacent seeding row. Use of press-wheels will minimise potential for herbicide to be dragged back into seeding rows. Weed control may be reduced in seeding rows as a result of concentration of herbicide in crop inter-rows. A knife or blade point is defined as being <12 mm in width, having no wings, inverted T or blade and is generally on a minimum 200 mm row spacing.

Full disturbance or conventional cultivation, including trailing harrows/prickle chain: Care should be taken to ensure seed placement below the herbicide band as crop damage may result where seed is sown too shallow. Consult trifluralin label for tank mix rate selection. PRODUCT MUST BE INCORPORATED INTO THE SOIL WITHIN 7 DAYS OF APPLICATION.

Mixing

REBELION 800 HERBICIDE is an emulsifiable concentrate that mixes readily with water. Fill the spray tank to one quarter full. Add REBELION 800 HERBICIDE and continue adding water to make up to the final spray volume. Agitate while mixing and spraying.

When tank mixing wettable powder or water dispersible granule formulations should be added to the tank first followed by suspension concentrates (flowables), water soluble salts then REBELION 800 HERBICIDE or other emulsifiable concentrate formulations. Maintain thorough agitation during mixing and application. Agitate tank mixes vigorously if allowed to stand. Note: Tank mix spray solutions should NOT be left standing in the vat overnight.

Sprayer Clean Up

After using REBELION 800 HERBICIDE, empty the tank completely and drain the whole system. Thoroughly wash inside the tank using a pressure hose, drain the tank and clean any filters in the tank, pump, line and nozzles.

To rinse : After cleaning the tank as above, quarter fill the tank with clean water and circulate through the pump, lines, hoses and nozzles. Drain and repeat the rinsing procedure twice.

Compatibility

REBELION 800 HERBICIDE is compatible with Kenso Agcare Trifluralin 480 Selective Herbicide. Other products may be compatible with REBELION 800 HERBICIDE, please refer to your Kenso Agcare Pty Ltd representative.

As formulations of other manufacturers' products are beyond the control of KENSO AGCARE PTY LTD and water quality varies with location, all mixtures should be tested prior to mixing commercial quantities.

Application

Apply by ground rig only in a minimum of 50 L water/ha. Water rate selection should be based on soil type and stubble load. Stubble loads above 40 to 50% ground coverage can reduce weed control below acceptable levels. Water volumes greater than 70 L/ha are recommended in order to reduce the impact of stubble in direct drill or minimum tillage systems. Use a nozzle delivering spray quality in the medium to very coarse spray range. DO NOT apply by air.

Resistance Management

Large numbers of healthy surviving weeds can be an indication that resistance is developing. Efforts should be made to prevent seed set of the surviving weeds. DO NOT make more than 1 application of a Group J herbicide with the inhibition of fat acid synthesis mode of action to a crop in the same season.

If the user suspects that the target weed population is resistant to herbicides with this mode of action, REBELION 800 HERBICIDE or other Group J herbicides should not be used. Strategies to minimise the risk of herbicide resistance are available. The above recommendations should be incorporated into an Integrated Weed Management (IWM) Program that uses a diverse range of options to manage grass weeds and avoids over reliance on any one method of control. Consult your farm chemical supplier, consultant, local Department of Agriculture or Primary Industries, or local KENSO AGCARE PTY. LTD. representative for details.

Tank Mixes

Application of REBELION 800 HERBICIDE at less than 2.5 L/ha when tank mixing for Annual Ryegrass control WILL NOT be effective in the control of Group D resistant populations nor to delay the onset of herbicide resistance development. Tank mixing of herbicides is only effective in managing resistance where a lethal dose rate of each herbicide is applied. Whilst a tank mixture of 2 herbicides of differing modes of action at sub-lethal dose rates may still provide a high level of weed control, the risk of developing resistance to multiple modes of action is increased. Tank mixtures of REBELION 800 HERBICIDE with trifluralin are only recommended where targeting additional weed species, on sandy soils where soil moisture is limiting or where product movement into the seeding zone poses a crop safety risk.

Herbicide Resistance Warning

GROUP J HERBICIDE

REBELION 800 HERBICIDE is a member of the thiocarbamates group of herbicides. This product has the inhibitors of cell division mode of action. For weed resistance management, this product is a Group J herbicide. Some naturally occurring weed biotypes resistant to this product and other Group J herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by this product or other Group J herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, KENSO AGCARE PTY. LTD. accepts no liability for any losses that may result from the failure of this product to control resistant weeds. Advice as to strategies and alternative treatments that can be used should be obtained from your local supplier, consultant, local Department of Agriculture, Primary Industries Department or a KENSO AGCARE PTY. LTD. representative.

Precautions:

Re-entry Period

Do not enter treated areas until the spray has dried unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT sow susceptible crops within 9 months of herbicide application.

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

DO NOT allow spray to drift onto adjacent fallow land.

DO NOT apply on or near shrubs, trees, lawns or crops other than wheat or barley.

DO NOT drain or flush equipment on or near desirable trees or other plants, where their roots may extend, or in situations where by movement of soil or by seepage absorption of the herbicide may occur.