

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

KENSO AGCARE

TAEKWANDO 250 CS INSECTICIDE

ACTIVE CONSTITUENT: 250 g/L LAMBDA-CYHALOTHRIN

GROUP **3A** INSECTICIDE

For the control of certain insect pests in Cotton, Barley, Wheat and various field crops as per the Directions for Use.



Kenso Corporation (M) Sdn Bhd
Level 1, 98 Commercial Road,
Teneriffe QLD 4005
Phone (07) 3216 1188
www.kenso.com.au

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USE

CONTENTS: 5 Litres APVMA Approval No.: 85948/ 120508

KENSO AGCARE TAEKWANDO 250 CS INSECTICIDE

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Store in a locked room or place away from children, animals, food, feedstuffs, seed and fertilizers.

Triple-rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

SAFETY DIRECTIONS

Harmful if inhaled or swallowed. Will irritate the eyes and skin. Repeated exposure may cause allergic disorders. Facial skin contact may cause temporary facial numbness. Avoid contact with eyes and skin.

When opening the container, preparing the spray and using the prepared spray wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow-length PVC gloves and face shield.

If product on skin, immediately wash area with soap and water. Wash hands after use. After each day's use, wash gloves, face-shield and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Telephone Australia 13 11 26. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet, which can be obtained from www.kenso.com.au.

CONDITIONS OF SALE

Kenso Corporation (M) Sdn. Bhd. will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Kenso Corporation (M) Sdn. Bhd. is limited to the replacement of the goods or (at the option of Kenso Corporation (M) Sdn. Bhd.) the refund of the price paid and where possible sufficient part of the goods to enable proper examination being returned to Kenso Corporation (M) Sdn. Bhd. within thirty days of delivery.

In a Transport Emergency
Dial **000**
Police or Fire Brigade



9 328666 003274



Additional statements required by Globally Harmonised System of Classification and Labelling of Chemicals (GHS) and Safe Work Australia: **TOXIC IF SWALLOWED. MAY BE FATAL IF SWALLOWED AND ENTERS AIRWAYS. MAY CAUSE AN ALLERGIC SKIN REACTION. HARMFUL IF INHALED.**
Precautionary: Avoid breathing dust/fume/gas/mist/vapours/spray. Wash contacted area thoroughly after handling. Do not eat, drink or smoke when using this product. Use only outdoors or in a well-ventilated area. Contaminated work clothing should not be allowed out of the workplace. Wear protective gloves/protective clothing/eye protection/face protection. IF SWALLOWED: Immediately call a POISON CENTRE or doctor/physician. IF ON SKIN: Wash with plenty of soap and water. IF INHALED: Remove victim to fresh air and keep at rest in a position comfortable for breathing. Call a POISON CENTER or doctor/physician if you feel unwell. Specific treatment (see FIRST AID) on label. Rinse mouth. DO NOT induce vomiting. If skin irritation or rash occurs: Get medical advice/attention. Wash contaminated clothing before reuse. Store locked up.

Batch No.:

Date of Manufacture:

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

KENSO
aggreair

TAEKWANDO 250 CS

Insecticide

**ACTIVE CONSTITUENT: 250 g/L
LAMBDA-CYHALOTHRIN**

GROUP 3A INSECTICIDE

For the control of certain insect pests in Cotton, Barley, Wheat and various field crops as per the Directions for Use.

IMPORTANT: READ THIS LEAFLET BEFORE USE

APVMA Approval No.: 85948/120508



Kenso Corporation (M) Sdn. Bhd.
Level 1, 98 Commercial Road,
Teneriffe QLD 4005
Phone (07) 3216 1188

COMPATIBILITY

This product, when applied, as a water-based spray is compatible with the following actives: FLUAZIFOP-P, PIRIMIPHOS-METHYL 900SF, PROCYMIDONE*, Piri-Ken 500 WG, Speedy 250, Ken-Up and Para-Ken.

This product when used in an ultra low volume application is compatible with the following mineral spraying oils: DC-Tron Cotton*, DC-Tron*, Ulvapron*, Trycol*, Caltex Summer Spray Oil* and Omex*.

PRECAUTIONS

Human flagging is not supported unless flaggers are protected by engineering control such as vehicles with cabs.

Re-entry Period

DO NOT allow entry into treated areas until the spray has dried. If prior entry is necessary wear cotton overalls and chemical resistant gloves.

PROTECTION OF LIVESTOCK

Toxic to bees. DO NOT spray when bees are actively foraging. Risk is reduced by spraying in the early morning or late evening.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Dangerous to fish and aquatic invertebrates. DO NOT contaminate streams, rivers or waterways with Taekwando 250 CS or used container. Tail waters which flow from treated areas should be prevented from entering river systems. In case of spillage on floor or paved surfaces, soak up with sand, earth or synthetic absorbent and dispose of waste according to Australian Standards 2507, Storage and Handling of Pesticides. A strategy to minimise spray drift should be employed at all times when aerially applying sprays near sensitive areas. Such a strategy is illustrated by the cotton industry's Best Management Practice Manual.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Store in a locked room or place away from children, animals, food, feedstuffs, seed and fertilizers.

Triple-rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

SAFETY DIRECTIONS

Harmful if inhaled or swallowed. Will irritate the eyes and skin. Repeated exposure may cause allergic disorders. Facial skin contact may cause temporary facial numbness. Avoid contact with eyes and skin.

When opening the container, preparing the spray and using the prepared spray wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow-length PVC gloves and face shield. If product on skin, immediately wash area with soap and water. Wash hands after use. After each day's use, wash gloves, face-shield and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Telephone Australia 13 11 26. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet, which can be obtained from www.kenso.com.au.

CONDITIONS OF SALE

Kenso Corporation (M) Sdn. Bhd. will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Kenso Corporation (M) Sdn. Bhd. is limited to the replacement of the goods or (at the option of Kenso Corporation (M) Sdn. Bhd.) the refund of the price paid and where possible sufficient part of the goods to enable proper examination being returned to Kenso Corporation (M) Sdn. Bhd. within thirty days of delivery.

* Not a registered trademark of Kenso Corporation (M) Sdn Bhd

In a Transport Emergency
Dial **000**
Police of Fire Brigade



Kenso Corporation (M) Sdn. Bhd.
Level 1, 98 Commercial Road,
Teneriffe QLD 4005
Phone (07) 3216 1188

APVMA Approval No.: 85948/120508

DIRECTIONS FOR USE:

For ULV application: Taekwando 250 CS can be bulked up with spraying oils for all uses except those indicated in the critical comments in the Direction for Use table below.

Crop	Pest Controlled	State	Application Rate/ha	WHP	Critical Comments
Barley, Wheat	Pasture Webworm (<i>Hednota</i> spp)	NSW, Vic, Tas, SA and WA only	12 mL	14 days after harvest/ grazing	Pre-seeding the product can be tank mixed with knock down herbicides. Post crop emergence inspect crop regularly from sowing. Spray at first sign of damage. Use a minimum of 50 L water/ha. Apply at first sign of infestation before larvae are 10mm long. For best results apply at first sign of infestation before larvae are 10mm long. If larvae are larger than 10mm use the higher rate. Use a minimum of 50 L of water.
	Pink or Brown Cutworm (<i>Agrotis munda</i>)	All States	12 or 18 mL		
	Common Cutworm (<i>Agrotis infusa</i>)	NSW only	20 or 40 mL	14 days after harvest/ grazing	Treat as soon as possible after the autumn rains stimulate egg hatching and activity of existing larvae. This can be ascertained by monitoring soil populations in known areas. For best results spray when the larvae have surfaced to feed after rain. Preferably use a boom spray delivering 70 to 100 L water/ha. Use the lower rate until early June and the higher rate after mid-late June. DO NOT USE ULV APPLICATION FOR THIS PEST. If mites are present on an establishing crop, apply at first sign of crop emergence. Monitor crop regularly for re-infestation and re-spray if necessary.
	Blackhead Pasture Cockchafer (<i>Aphodius tasmaniae</i>)	NSW, Vic, Tas, SA and WA only			
	Redlegged Earth Mite (<i>Halotydeus destructor</i>)		9 mL †		
	Aphids (<i>Rhopalosiphum</i> spp) (Barley Yellow Dwarf Virus Vectors)		12 or 18 mL		To control aphids spray should be applied at 4 to 8 weeks after emergence to reduce aphid colonization and suppress Barley Yellow Dwarf Virus. Use the higher rate when greater than 15 aphids on 50% of tillers is expected during the season.
Broccoli, Brussel Sprouts, Cabbages, Cauliflowers, Forage brassica	Cabbage Cluster Caterpillar (<i>Crociodolomia pavonana</i>) Cabbage White Butterfly (<i>Pieris rapae</i>) Diamond Black Moth (<i>Plutella xylostella</i>)	Qld, NSW, Vic, WA and NT only	24 or 36 mL Add Agral* at 10mL/ 100L of spray volume	2 days after harvest/ grazing	Apply at first sign of infestation. For schedule spraying on a weekly basis, use the lower rate. For spraying as needed use the higher rate for longer persistence. Use a minimum of 500 L water/ha.
Canola	Cabbage Moth (<i>Plutella xylostella</i>) Cabbage White Butterfly (<i>Pieris rapae</i>)	All States	24 mL	7 days harvest/ grazing	Apply as soon as larvae reach threshold numbers. Check with local officer of the Department of Primary Industries for thresholds applicable to the particular growth stage of the crop.
	Rutherglen Bug and Grey Cluster Bug (<i>Nysius</i> spp.)		36 mL		Apply only near maturity when severe infestations are likely to down grade yields.
	Thrips (<i>Thrips tabaci</i>)	Qld, NSW, Vic, Tas, WA and NT only			
	Native Budworm (<i>Helicoverpa punctigera</i>)	NSW, Vic, Tas, SA and WA only	24 or 36 mL	7 days after harvest/ grazing	For best results, apply at hatching or soon after. Use higher rate if the crop is dense or the larvae are larger than 10mm.
	Redlegged Earth mites (<i>Halotydeus destructor</i>)	NSW, Vic, Tas, SA and WA only	9 ml †	7 days after harvest/ grazing	If mites are present on establishing crop, apply at the first sign of crop emergence. Monitor the crop regularly for reinfestation and respray if necessary.

Crop	Pest Controlled	State	Application Rate/ha	WHP	Critical Comments
Cotton	Native Budworm (<i>Helicoverpa punctigera</i>) Cotton Bollworm (<i>Helicoverpa armigera</i>)	Qld, NSW, NT and WA only	60 mL	21 days after harvest	Apply when egg laying is light less than 25 eggs/100 terminals and no larvae are present.
			70 mL		Apply when egg laying is moderate greater than 25 eggs/100 terminals and/or when less than 12 newly hatched larvae/100 terminals are present.
			85 mL		Apply when egg laying is heavy and continuous and/or when <i>H. punctigera</i> larvae are greater than 10mm in length. For <i>H. armigera</i> , apply only to larvae less than 5mm in length.
	Pink-spotted Bollworm (<i>Pectinophora scutigera</i>)	Qld and NT only	70 mL		Controlled with the <i>Helicoverpa</i> spp. program when used at this rate. If the pink-spotted bollworm is the only pest present, apply when more than 10 adult moths are caught in pheromone traps on two consecutive nights.
	Green Mirid (<i>Creontiades dilutus</i>) Brown Mirid (<i>C. pacificus</i>) Apple Dimpling Bug (<i>Campylomma liebknechti</i>) Broken backed Bug (<i>Taylorilygus pallidulus</i>) Cottonseed Bug (<i>Oxycarenus luctuosus</i>) Pale Cotton Stainer (<i>Dysdercus sidae</i>) Leafhoppers (<i>Austroasca viridigrisea</i> and <i>Amrasca terraereginae</i>)	Qld, NSW, NT and WA only	60 mL		Apply at the recommended threshold levels as indicated by field checks.
Faba Beans, Chick Peas, Vetch, Lentils	Native Budworm (<i>Helicoverpa punctigera</i>)	NSW, Vic, SA and WA only	24 or 36 mL	7 days after harvest/ grazing	For best results, apply at hatching or soon after. Use the higher rate if the crop is dense or the larvae are larger than 10 mm.
	Redlegged Earth Mite (<i>Halotydeus destructor</i>)	NSW, Vic, Tas, SA and WA only	9 mL †		If mites are present on an established crop, apply at first sign of crop emergence. Monitor crop regularly for reinfestation and respray if necessary. Control of Lucerne Flea will not be obtained with application.
Field Peas	Native Budworm (<i>Helicoverpa punctigera</i>)	NSW, Vic, SA and WA only	24 or 36 mL	7 days after harvest/ grazing	For best results, apply at hatching or soon after. Use higher rate if the crop is dense or the larvae are larger than 10 mm.
			Pea Weevil (<i>Bruchus pisorum</i>)	NSW and SA only	24 mL
		Vic and WA only	36 mL		
		Redlegged Earth Mite (<i>Halotydeus destructor</i>)	NSW, Vic, Tas, SA and WA only	9 mL †	7 days after harvest/ grazing

Crop	Pest Controlled	State	Application Rate/ha	WHP	Critical Comments
Lemons, Oranges	Fullers Rose Weevil (<i>Asynonychus cervinus</i>)	All States	300mL/100L as a directed spray	4 weeks (H)	Firstly ensure that the trees are skirted and all weeds under the trees are removed. Apply 250 mL spray solution to the tree trunk at about 300mm from the ground in a 100mm band. Deliver the spray through a U shaped wand fitted with 4 nozzles evenly spaced around the tree. Trees must be treated in the early stages of the adults weevils emerging from the grounds.
Lucerne	Lucerne Leaf Roller (<i>Merophyas divulsana</i>)	All States	24 or 36 mL	14 days after harvest/ grazing	For best results apply at hatching or soon after. Use higher rate if the crop is dense or the larvae are larger than 10mm. Apply the first spray when about 30% of the terminals are rolled.
	Native Budworm (<i>Helicoverpa punctigera</i>)				For best results apply at hatching or soon after. Use higher rate if the crop is dense or the larvae are larger than 10mm.
	Pea Aphid (<i>Acyrtosiphon pisum</i>)		24 mL		Good coverage, particularly the stems, is essential. Use hollow cone nozzles.
	Blackhead Pasture Cockchafer (<i>Aphodius tasmaniae</i>)	NSW, Vic, Tas, SA and WA only	20 or 40 mL		Treat as soon as possible after the autumn rains stimulate egg hatching and activity of existing larvae. This can be ascertained by monitoring soil populations in known areas. For best results spray when the larvae have surfaced to feed after rain. Preferably use a boom spray delivering 70 to 100 L water/ha. Use the lower rate until early June and the higher rate after mid-late June. DO NOT USE ULV APPLICATION FOR THIS PEST.
	Redlegged Earth Mite (<i>Halotydeus destructor</i>)		9 mL †		If mites are present on an establishing crop, apply at first sign of crop emergence. Monitor crop regularly for re-infestation and re-spray if necessary. Control of Lucerne Flea will not be obtained with this application.
Lupins	Native Budworm (<i>Helicoverpa punctigera</i>)	NSW, Vic, SA and WA only	24 mL	14 days after harvest/ grazing	Once crop has emerged, inspect regularly and apply at the first signs damage. Use a minimum of 50 L water/ha. DO NOT USE ULV APPLICATION FOR THIS PEST.
	Brown Pasture Looper (<i>Ciampa arietaria</i>)	NSW, Vic, Tas, SA and WA only	12 mL		For best results, apply at hatching or soon after when larvae are small. WA only: Environmental factors may cause populations of small caterpillars to decline, reducing damage potential. Spraying should commence once caterpillars are 12 mm in length.
	Redlegged Earth Mite (<i>Halotydeus destructor</i>)		9 mL †		If mites are present on an establishing crop, apply at the first sign of crop emergence. Monitor crop regularly for reinfestation and respray if necessary. Control of Lucerne Flea will not be obtained with application.
Navy Beans Mung Beans Beans	Native Budworm (<i>Helicoverpa punctigera</i>) Corn Earworm (<i>Helicoverpa armigera</i>)	Qld, NSW and NT only	60 or 70 mL	Harvest/ grazing 1 day if harvested green, 14 days if harvested dry	Apply when flower or pod feeding larvae reach a population of 1 to 2 per metre of row in navy beans and 1 per metre of row in mung beans. Use the higher rate if pest numbers are high or if larvae are larger than 10 mm. In northern NSW and Qld where corn earworm has established resistance to pyrethroids DO NOT apply to corn earworm larvae larger than 5 mm.
Onions bulb	Onion Thrips	All States	40mL	14 days (H)	Apply when thrips first appear. Apply via ground equipment in a minimum 300L water/ha. DO NOT exceed a maximum of 4 applications per crop with a minimum retreatment interval of 7 days between consecutive sprays.
Pasture	Pasture Webworm (<i>Hednota</i> spp.)	NSW, Vic, Tas, SA and WA only	12 mL	14 days after harvest/ grazing	Apply once larvae are present using adequate water to ensure good penetration.
	Brown Pasture Looper (<i>Ciampa arietaria</i>)	All States	12 mL		DO NOT USE ULV APPLICATION FOR THIS PEST.
	Pink or Brown Cutworm (<i>Agrotis munda</i>)	All States	12 to 18 mL		For best results apply at first sign of infestation before larvae are 10 mm long. If larvae are larger than 10 mm, use the higher rate. Use a minimum of 50 L water.
	Common Cutworm (<i>Agrotis infusa</i>)	NSW only			
	Blackhead Pasture Cockchafer (<i>Aphodius tasmaniae</i>)	NSW, Vic, Tas, SA and WA only	20 or 40 mL		Treat as soon as possible after the autumn rains stimulate egg hatching and activity of existing larvae. This can be ascertained by monitoring soil populations in known areas. For best results spray when the larvae have surfaced to feed after rain. Preferably use a boom spray delivering 70 to 100 L water/ha. Use the lower rate until early June and the higher rate after mid-late June. DO NOT USE ULV APPLICATION FOR THIS PEST.
	Redlegged Earth Mite (<i>Halotydeus destructor</i>)		9 mL †		If mites are present on an establishing crop, apply at first sign of crop emergence. Monitor crop regularly for re-infestation and re-spray if necessary. Control of Lucerne Flea will not be obtained with this application.
Potatoes	Vegetable Jassid (<i>Austroasca viridigrisea</i>)	All States	24 mL	7 days (H)	Apply only when numbers is excessive.
Sorghum	Sorghum Midge (<i>Contarinia sorghicola</i>)	Qld, NSW and NT only	18 or 36 mL	14 days after harvest/ grazing	Apply when midge numbers reach 1 to 2 per head. Use the higher rate for residual protection.
	Corn Earworm (<i>Helicoverpa armigera</i>)		60 or 70 mL		Apply when larval numbers reach 2 per head. Use the higher rate if pest pressure is severe. Best results are achieved on small larvae.

Crop	Pest Controlled	State	Application Rate/ha	WHP	Critical Comments
Sunflower	Rutherglen Bug and Grey Cluster Bug (<i>Nysius</i> spp.)	All States	36 mL	28 days after harvest	Apply when numbers reach 10 to 15 adults per plant at budding in dry land crops or 20 to 25 in irrigated crops. If <i>Helicoverpa armigera</i> are also present in northern NSW or Qld, use a minimum of 60 mL product.
	Native Budworm (<i>Helicoverpa punctigera</i>) Corn Earworm (<i>Helicoverpa armigera</i>)	Qld and Nth NSW only Sth NSW and Vic only	60 or 70 mL 48 or 60 mL	28 days after harvest	Apply when an average of 2 to 3 larvae are present per head or when larvae are damaging plants. Use the higher rate if pest numbers are high and/or <i>H. punctigera</i> larvae are larger than 10 mm in length. In Northern NSW and Qld, DO NOT apply to resistant <i>H. armigera</i> larvae larger than 5 mm in length. GENERAL COMMENTS: If flowering has started, application should be deferred until after flowering but before the heads turn down. If treatment is unavoidable during flowering, and bees are actively foraging in the crop, there will be minimal effect in the colony if spraying occurs early morning or late afternoon.
Soybeans	Native Budworm (<i>Helicoverpa punctigera</i>) Corn Earworm (<i>Helicoverpa armigera</i>)	Qld, NSW, Vic and NT only	60 or 70 mL	21 days after harvest/ grazing	Apply when flower or pod feeding larvae reach a population of 2 per metre of row in soybeans. Use the higher rate if pest numbers are high or if larvae are larger than 10 mm. In northern NSW and Qld DO NOT apply to resistant <i>H. armigera</i> larvae larger than 5 mm in length.
Tomatoes bush	Native Budworm (<i>Helicoverpa punctigera</i>)	All States	4 or 5 mL/100L or 30 or 36 mL/ha	1 day (H)	Treat plant on a 7 to 14 day schedule. Qld, Northern NSW: DO NOT apply to <i>H. armigera</i> larvae larger than 5mm in length. Other areas: For best results apply soon after egg lay. To help contain resistance, alternate sprays between different chemical groups. Check the crop every few days and follow the Summer Crop Insecticide Strategy. There may be phytotoxicity with some varieties especially Floradade.
	Tomato Grub (<i>Helicoverpa armigera</i>)	Vic, Tas, SA, WA only Qld, NSW, NT only	4 mL/100L or 60 mL/ha		
Tomatoes trellis	Native Budworm (<i>Helicoverpa punctigera</i>) Tomato Grub (<i>Helicoverpa armigera</i>)	All States	4 or 5 mL/100L	1 day (H)	

† Blue Oat Mites often co-occur with Redlegged Earth Mites and the 9 mL/ha rates of Taekwando 250 CS may be less effective against Blue Oat Mites.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHOLDING PERIODS:

Harvesting:

Mung Beans (if harvested green), Navy Beans (if harvested green), Tomatoes:

DO NOT HARVEST FOR 1 DAY AFTER APPLICATION.

Cabbages, Cauliflowers, Broccoli, Brussel Sprouts: **DO NOT HARVEST FOR 2 DAYS AFTER APPLICATION.**

Field peas, Canola, Faba Beans, Chick Peas, Lentils, Potatoes, Vetch: **DO NOT HARVEST FOR 7 DAYS AFTER APPLICATION.**

Lupins, Sorghum, Mung Beans (if harvested dry), Navy Beans (if harvested dry), Barley, Lucerne, Wheat, Pasture, Onions: **DO NOT HARVEST FOR 14 DAYS AFTER APPLICATION.**

Cotton, Soybeans: **DO NOT HARVEST FOR 21 DAYS AFTER APPLICATION.**

Lemons, Oranges, Sunflower: **DO NOT HARVEST FOR 4 WEEKS AFTER APPLICATION.**

Grazing:

Navy Beans (if harvested green), Mung Beans (if harvested green): **DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 1 DAY AFTER APPLICATION.**

Forage Brassicas: **DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 2 DAYS AFTER APPLICATION.**

Field peas, Canola, Faba Beans, Chick Peas, Vetch, Lentils: **DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION.**

Lupins, Sorghum, Navy Beans (if harvested dry), Mung Beans (if harvested dry), Barley, Wheat, Pasture, Lucerne: **DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 14 DAYS AFTER APPLICATION.**

Soybeans: **DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 21 DAYS AFTER APPLICATION.**

GENERAL INSTRUCTIONS INSECTICIDE RESISTANCE WARNING

GROUP 3A INSECTICIDE

For insecticide resistance management Kenso Agcare Taekwando 250 CS Insecticide is a Group 3A insecticide. Some naturally occurring insect biotypes resistant to Taekwando 250 CS and other Group 3A insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if Taekwando 250 CS or other Group 3A insecticides are used repeatedly. The effectiveness of Taekwando 250 CS on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, Kenso Agcare Pty Ltd accepts no liability for any losses that may result from the failure of Taekwando 250 CS to control insects.

Taekwando 250 CS may be subject to specific resistance management strategies. For further information contact your local supplier, Kenso Agcare representative or local agricultural department agronomist. *Helicoverpa* (*Heliothis armigera*) resistance in Northern New South Wales and Queensland: To help contain pyrethroid resistance in *Helicoverpa armigera*, the Summer Crop Insecticide Strategy as developed by AIRAC, Queensland Department of Primary Industries and the New South Wales Department of Agriculture and Fisheries should be adhered to. Failure to observe the strategy may result in widespread resistance affecting the future viability of summer cropping.

MIXING: SHAKE WELL BEFORE USE.

For ground or aircraft application with water: Taekwando 250 CS mixes readily with hard or soft water. Add the required quantity of product to water whilst under agitation to ensure thorough mixing. Agitate while spraying. It is not advisable to allow the mixed solution to stand longer than 24 hours before use. In extremely alkaline water (pH 9) spray immediately after mixing.

For ULV (ultra low volume) application with oil: It is recommended that Taekwando 250 CS be mixed with a mineral, spraying oil. See Compatibility section for a list of recommended mineral spraying oils. Add the required quantity of product to oil whilst under agitation to ensure thorough mixing. Agitate while spraying. It is not advisable to allow the mixed solution to stand longer than 24 hours before use.

APPLICATION

Good coverage is essential to ensure adequate control. The product may be applied by ground rig or aircraft. DO NOT apply if rain is expected within 6 hours. Acceptable "threshold" values for eggs and larval numbers may vary according to the stage of crop development and the pest management program undertaken. Alternative higher thresholds may be acceptable under certain circumstances.

Diluted with water: For ground rigs the volume of liquid applied should be 30 to 100 L/ha. Aerial application should be under conditions normally suitable for water-based insecticides. Apply in at least 10 to 20 L water/ha.

Mixed with oil: Apply the recommended rate of Taekwando 250 CS bulked with oil to total volume of 3 to 5 L/ha for cotton, sorghum and sunflowers. The total volume for all other crops should be 1.5 L/ha.

TIMING

This product is a contact and residual insecticide. Best result will be obtained if Taekwando 250 CS is applied as a protective treatment at regular intervals. However, if spraying frequency based on scouting, then for *Helicoverpa* spp. application at egg hatch will give optimum results.

CROP CHECKING

Frequent and thorough checking of whole plants, terminals, squares, flowers, bolls or fruiting bodies as required, should be made over a random sample of plants, representative of the whole crop area. Inspect crops after spraying to ensure a thorough kill has been obtained. However, note that a maximum kill may not be achieved until 48 hours after treatment. Then check at frequent intervals, not more than 2 days apart when insect pressure is heavy. Apply the recommended treatment as soon as a crop check indicates spraying is necessary.