

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

KENSO AGCARE

TEXUS

SELECTIVE HERBICIDE

Active Constituents: 250 g/L MCPA present as the ETHYL HEXYL ESTER
25 g/L DIFLUFENICAN
Solvents: 325 g/L HYDROCARBONS LIQUID
150 g/L N-METHYL-2-PYRROLIDONE

GROUP	F	I	HERBICIDE
-------	----------	----------	-----------

For the control of certain broadleaf weeds in winter cereals and clover as specified
in the DIRECTIONS FOR USE table.

Kenso Corporation (M) Sdn Bhd
Level 1, 98 Commercial Road,
Teneriffe QLD 4005
Phone (07) 3216 1188
www.kenso.com.au



IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USE

CONTENTS: 20 Litres APVMA Approval No.: 66669/ 121534

KENSO AGCARE TEXUS SELECTIVE HERBICIDE

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight.

Triple-rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

110L Mini Bulk Returnable Container (110L only)

Store the original sealed drum in a cool well-ventilated area. DO NOT store for prolonged periods in direct sunlight. DO NOT tamper with the non-return valve or the security seal. DO NOT contaminate the drum with water or any foreign matter. After each use of the product, please ensure that the non-return valve, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the drum have been used, please return the empty drum to the point of purchase.

Refillable containers (1000L only):

Store in the closed, original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight. Empty contents fully into application equipment. Close all valves.

SAFETY DIRECTIONS

Harmful if swallowed. Will damage the eyes. Will irritate the skin. Avoid contact with eyes and skin. When opening the container and preparing the spray, wear cotton overalls buttoned to the neck and wrist, washable hat, elbow length PVC gloves and face shield or goggles. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use wash gloves, face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a Doctor or Poisons Information Centre (Phone Australia 13 11 26). If swallowed, DO NOT induce vomiting. Give a glass of water. If in eyes, wash out immediately with water.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet, which can be obtained from the supplier.

CONDITIONS OF SALE

Kenso Corporation (M) Sdn. Bhd. will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Kenso Corporation (M) Sdn. Bhd. is limited to the replacement of the goods or (at the option of Kenso Corporation (M) Sdn. Bhd.) the refund of the price paid and where possible sufficient part of the goods to enable proper examination being returned to Kenso Corporation (M) Sdn. Bhd. within thirty days of delivery.

In a Transport Emergency
Dial **000** Police or Fire Brigade



Additional statements required by Globally Harmonised System of Classification and Labelling of Chemicals (GHS) and Safe Work Australia: **COMBUSTIBLE LIQUID. HARMFUL IF SWALLOWED. HARMFUL IN CONTACT WITH SKIN. CAUSES SKIN IRRITATION. CAUSES SERIOUS EYE IRRITATION. HARMFUL IF INHALED. MAY CAUSE RESPIRATORY IRRITATION.** *Precautionary:* Keep away from heat/sparks/open flames/hot surfaces. — No smoking. Avoid breathing dust/fume/gas/mist/vapours/spray. Wash contacted area thoroughly after handling. Do not eat, drink or smoke when using this product. Use only outdoors or in a well-ventilated area. Wear protective gloves/ protective clothing/eye protection/face protection. IF SWALLOWED: Call a POISON CENTER or doctor/physician if you feel unwell. IF ON SKIN: Wash with plenty of soap and water. IF INHALED: Remove victim to fresh air and keep at rest in a position comfortable for breathing. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. Call a POISON CENTER or doctor/physician if you feel unwell. Specific treatment (see FIRST AID on this label). Rinse mouth. If eye irritation persists: Get medical advice/attention. If skin irritation occurs: Get medical advice/attention. Take off contaminated clothing and wash before reuse. In case of fire: Use water fog, foam, dry agent (carbon dioxide, dry chemical powder) for extinction. Store in a well-ventilated place. Keep container tightly closed. Keep cool. Store locked up.

Batch No.:

Date of Manufacture:

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

KENSO
ag science

TEXUS

Selective Herbicide

Active Constituents: 250 g/L MCPA present as the ethyl hexyl ester
25 g/L DIFLUFENICAN
Solvents: 325 g/L HYDROCARBONS LIQUID
150 g/L N-METHYL-2-PYRROLIDONE

GROUP F I HERBICIDE

For the control of certain broadleaf weeds in winter cereals and clover as specified in the DIRECTIONS FOR USE table.

READ THIS LEAFLET BEFORE USE

APVMA Approval No.: 66669/121534



Kenso Corporation (M) Sdn. Bhd.
Level 1, 98 Commercial Road,
Teneriffe QLD 4005
Phone (07) 3216 1188

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple-rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

110L Mini Bulk Returnable Container

Store the original sealed drum in a cool well-ventilated area. DO NOT store for prolonged periods in direct sunlight. DO NOT tamper with the non-return valve or the security seal. DO NOT contaminate the drum with water or any foreign matter. After each use of the product, please ensure that the non-return valve, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the drum have been used, please return the empty drum to the point of purchase.

Refillable containers (1000L only):

Store in the closed, original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight. Empty contents fully into application equipment. Close all valves.

SAFETY DIRECTIONS

Harmful if swallowed. Will damage the eyes. Will irritate the skin. Avoid contact with eyes and skin. When opening the container and preparing the spray, wear cotton overalls buttoned to the neck and wrist, washable hat, elbow length PVC gloves and face shield or goggles. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use wash gloves, face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a Doctor or Poisons Information Centre (Phone Australia 13 11 26). If swallowed, DO NOT induce vomiting. Give a glass of water. If in eyes, wash it out immediately with water.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet which can be obtained from the supplier.

CONDITIONS OF SALE

Kenso Corporation (M) Sdn. Bhd. will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Kenso Corporation (M) Sdn. Bhd. is limited to the replacement of the goods or (at the option of Kenso Corporation (M) Sdn. Bhd.) the refund of the price paid and where possible sufficient part of the goods to enable proper examination being returned to Kenso Corporation (M) Sdn. Bhd. within thirty days of delivery.

® Not a registered trademark of Kenso Corporation (M) Sdn Bhd

In a Transport Emergency
Dial **000**
Police of Fire Brigade



Kenso Corporation (M) Sdn. Bhd.
Level 1, 98 Commercial Road,
Teneriffe QLD 4005
Phone (07) 3216 1188

APVMA Approval No.: 66669/121534

DIRECTIONS FOR USE

Restraints

DO NOT apply if crop or weeds are stressed due to dry or excessively moist conditions.
DO NOT apply to crops under stress due to disease or insect damage.
DO NOT apply to frost-affected crops or if frosts are imminent.
DO NOT apply when heavy rain is expected within 4 hours.

CROP	WEEDS CONTROLLED	STAGE OF WEED GROWTH	STATE	RATE/ HA	CRITICAL COMMENTS	
CEREALS Wheat, barley, oats, triticale, cereal rye (including cereals undersown with clover)	Wild radish	Up to the 2 leaf stage and not more than 60 mm in diameter	WA only	250 mL	CROP STAGE Cereals Up to 750 mL (3 leaf to fully tillered stage - Z13 to 30) Over 750 mL (5 leaf to fully tillered stage - Z15 to 30) Optimum results are achieved when sprayed at 3-5 leaf crop stage (generally 4-8 weeks post sowing). WA only: DO NOT apply to Barley or Kulin Wheat before the 5 leaf stage (Z15). Warning: Texus may cause transient crop yellowing of cereals. Some varieties of oats have not been tested. (Refer to " Crop Tolerance " section of General Instructions) Clover Application is recommended prior to the eighth trifoliolate leaf stage, however, applications prior to the third leaf stage may result in crop damage especially under stressed conditions and in sandy soils. DO NOT apply to Annual Medics or lucerne. Warning: Texus may cause transient crop yellowing of clover, and may affect growth and seed set of some varieties of clover. (Refer to " Crop Tolerance " section of General Instructions). WEED STAGE Apply when weeds are actively growing. In most situations the rate specified for each weed size will give satisfactory control. Under certain conditions such as: * high crop and weed density * late season germinations * abnormal weed growth (including early flowering), higher rates of product (up to the maximum rate of application specified for that weed) may be required. Texus will not effectively control: * regrowth of suppressed weeds; * transplanted weeds; * regrowth from rhizomes or roots; * weeds growing under stress from previous herbicide applications CONTINUED OVERLEAF	
		Up to the 4 leaf stage and not more than 120 mm in diameter	All States	500 mL		
		Up to the 5 leaf stage and not more than 150 mm in diameter		750 mL		
		Up to the 8 leaf stage and not more than 180 mm in diameter		1.0 L		
	Charlock, hedge mustard, Indian hedge mustard, shepherd's purse, turnip weed, wild turnip	Up to the 2 leaf stage and not more than 60 mm in diameter	Up to the 4 leaf stage and not more than 120 mm in diameter	All States		500 mL
						750 mL
						1.0 L
	London rocket	Up to the 5 leaf stage and not more than 120 mm in diameter	Ward's weed	Qld only		750 mL
				SA only		
	Capeweed	Up to the 2 leaf stage and not more than 60 mm in diameter	Up to the 4 leaf stage and not more than 120 mm in diameter	All States		500 mL
						1.0L
	Crassula	Up to the 2 leaf stage	Up to the 4 leaf stage	All States		500mL
						750mL
	Prickly lettuce	Up to the 2 leaf stage	Up to the 4 leaf stage	All States		500mL
						750mL
1.0L						
Dense-flower fumitory	Up to the 2 leaf stage		All States	750 mL		
				1.0L		
Corn gromwell, saffron thistle, toad rush			All States	1.0L		
				1.0L		
Deadnettle			NSW, Vic, SA only	750 mL		
				1.0L		
Sorrel	Up to the 2 leaf stage		Vic only	1.0 L		
Canola (rape)	Up to the 4 leaf stage		All States	500 mL		
Purple goosefoot	Up to the 6 leaf stage		Qld only	500 mL		
Turnip weed, wild turnip	Cotyledon to 2 leaf stage		NSW only (West of Newell Hwy.)	350 mL		
					SA only (Eyre peninsula north of the line between Venus Bay and Cowell)	

CROP	WEEDS CONTROLLED	STAGE OF WEED GROWTH	STATE	RATE/ HA	CRITICAL COMMENTS				
CEREALS Wheat, barley, oats, triticale, cereal rye	Fumitory	2 - 6 leaf stage	All States	500 mL + 200 mL terbutryn (500 g/L)	CONTINUED FROM PREVIOUS PAGE				
					GRAZING Efficacy on larger weeds will be improved by grazing with normal levels of stock after the 7 day withholding period. Refer to Protection of Livestock' for grazing precautions.				
CEREALS Wheat, barley, oats, triticale, cereal rye (including cereals undersown with clover)	SUPPRESSION OF THE FOLLOWING WEEDS								
	Saffron thistle	Up to the 6 leaf stage	All States	1.0 L	APPLICATION Activity of this product will be reduced if weeds are stressed. Optimum results will be obtained if good soil moisture exists at and after application. Where crop or weed density is high, water volume should be increased. WILD RADISH Texus will provide residual control of Wild Radish for up to 4 weeks after application. Effective residual activity of this product may be reduced where: * rates lower than 1.0 L/ha are used; * dry conditions prevail; * poor coverage of the soil surface is achieved; * crop is planted in non-wetting sand; * soils contain a high content of organic matter. Optimum results will be obtained if good soil moisture exists at and after application. Refer also to all Critical Comments relating to weed stage, grazing, application and wild radish above. * Reduced efficacy (suppression only) may be achieved on wild radish larger than 8 leaf or greater than 180 mm in diameter. DO NOT use this tank-mix if cereals are undersown with lucerne or annual medics. <u>Crop Stage</u> Texus 350 mL + LV MCPA 200 mL: Apply from 3 leaf to fully tillered (Zadok's Z13 to Z30). Texus 500 mL + LV MCPA 200 mL: Apply from 3 leaf to fully tillered (Zadok's Z13 to Z30). Texus 500 mL + LV MCPA 400 mL: Apply from 5 leaf stage to fully tillered (Zadok's Z15 to Z30). Optimum results are achieved when sprayed at 3-5 leaf crop stage (generally 4-8 weeks post sowing). WA only: DO NOT apply to Barley or Kulin Wheat before the 5 leaf stage (Z15). Warning: Texus may cause transient crop yellowing of cereals. Some varieties of oats have not been tested. (Refer to " Crop Tolerance " section of General Instructions) Observe instructions also on LV MCPA product label.				
						Chickweed, fireweed, hexham scent (King Island mellilot), iceplant, mouse-eared chickweed, night-scented stock, Paterson's curse, peppercress, skeleton weed, long storksbill, volunteer lupins	Up to the 4 leaf stage	All States	1.0 L
	Common sowthistle (milk thistle), cowvine, dock, doublegee (spiny emex), fat hen, horehound, hyssop loosestrife, marshmallow, rough poppy, scarlet pimpernel, stemless thistle, tree hogweed, variegated thistle, vetch (tares)	Up to the 2 leaf stage	All States	1.0 L					
					CEREALS	Wild radish	Up to the 4 leaf stage and not more than 120 mm in diameter	350 mL plus 200 mL LV MCPA (500 g/L)	
									Wheat, barley, oats, triticale, cereal rye
	Up to the 8 leaf stage and not more than 180 mm in diameter*	500 mL plus 400 mL LV MCPA (500 g/L)							

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS

CROP HARVEST: NOT REQUIRED WHEN USED AS DIRECTED

All crops: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION

GENERAL INSTRUCTIONS

Resistant Weeds Warning

Texus is a member of the phenoxy and nicotinilide groups of herbicides and acts by inhibiting carotenoid biosynthesis and disrupting plant cell growth. For weed resistance management Texus is both a Group F and a Group I herbicide. Some naturally occurring weed biotypes resistant to Texus and other Group F and I herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Texus or other Group F or Group I herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Kenso Corporation (M) Sdn Bhd accepts no liability for any losses that may result from the failure of Texus to control resistant weeds.

Tolerance

Some pre-emergence herbicides, such as atrazine, can cause stress to certain crops resulting in an increase in crop damage when using this product. Subterranean clover is particularly sensitive.

Cereals

After application some transient crop yellowing may occur. This usually appears as yellow or white banding on leaves.

Provided the crop is not under stress from pre-emergent herbicide, root disease, insect damage, frost, dry or excessively moist conditions, the development of the crop and subsequent growth will be unaffected.

Warning (Oats)

The tolerance of oat varieties Esk and Nile (the two main varieties grown in Tasmania) to Texus has not been tested.

Test a small area of crop before using Texus over large areas. Consult your local Kenso Agcare representative for advice on specific varieties.

Pasture

The tolerance of clover varieties to Texus can vary with rate of application, soil type, crop health, stage of growth and degree of moisture and temperature stress.

Warning

Texus may result in transient crop yellowing and suppression of growth with a resultant initial reduction in dry matter, particularly at rates in excess of 500 mL/ha and in areas of double spray. For this reason we recommend application prior to the 8 trifoliolate leaf stage. However, at the lower rates (500 mL/ha and less) and under normal growing conditions, subsequent growth and seed yield should not be affected.

Under normal growing conditions, the following varieties have shown acceptable levels of foliage tolerance to Texus applied at 500 mL/ha:

ArrowLeaf: Zulu

Balansa: Paradana

Berseem: Sacromonte

Persian: Kyambro, Lupers, Maral

White: Haifa

Subterranean clover: Daliak, Dalkeith, Denmark, Esperance, Geraldton, Goulburn, Karridale, Larissa, Leura, Mt.Barker, Nungarin, Rosedale, Seaton Park, Trikkala and Woogenellup.

The effects of Texus on clover seed yield have been tested on the following varieties. Under normal growing conditions they show acceptable levels of tolerance to Texus applied at 500 mL/ha:

Subterranean clover: Esperance, Goulburn, Larissa, Seaton Park and Trikkala.

Warning

Rose and Strawberry clover have shown increased sensitivity to Texus. Texus may affect the seed yield of subterranean clover variety Woogenellup.

Some pasture grasses, including Phalaris and Cocksfoot, may show some initial reduction in vegetative growth after application of Texus.

Care should be exercised if sensitive clover varieties or grasses are included in the pasture sward.

Varieties not listed should be tested before using Texus over large areas. Consult your local Kenso Agcare representative for advice on specific varieties.

Subsequent Crops

To reduce effect on subsequent susceptible crops (e.g. canola), ensure thorough cultivation of soil prior to the sowing of these crops.

Mixing

To ensure even mixing, half fill the spray tank with clean water and add the required amount of product. Agitate thoroughly then add the remainder of the water. Agitate again before spraying commences. Reseal part-used product container immediately after use. Spray mixtures containing Texus should not be left to stand overnight. Prolonged periods of exposure to cold temperatures could result in settling out of the product in the mixture.

Warning

The rubber components present in some spraying units may be affected by exposure to the solvents in Texus and some other agricultural products. To reduce this risk it is recommended that the spray unit be thoroughly washed with a boom cleaner and fresh water after use. Contact the spray unit manufacturer to determine the suitability of the rubber components for use with agricultural products.

Application

Boom Sprayer

A minimum of 50 L of water per hectare should be used, however, for optimum results water rates of 70-100 L/ha are recommended. Increase the water volume if weed infestation is heavy or crop cover is dense. Complete coverage of weeds is essential.

Aircraft (NSW, Vic, SA only)

Apply in a minimum of 30 L water per hectare. Effective weed control will only be achieved where good coverage of leaf surface is achieved.

Compatibility

The following products are physically compatible with Texus as a two-way mixture in the spray tank but should only be used for the crops specified:

Crop	Texus	Compatible Product
Wheat, triticale and cereal rye	Wheat, triticale and cereal rye	Wheat, triticale and cereal rye
Cereals (including undersown) All rates Chlorpyrifos (500 g/L), dimethoate, Thiodan®.	Cereals (including undersown) All rates Chlorpyrifos (500 g/L), dimethoate, Thiodan®.	Cereals (including undersown) All rates Chlorpyrifos (500 g/L), dimethoate, Thiodan®.
Cereals (not undersown)	Up to 500 mL/ha All rates	Ken-Met 600, Ken-Chlor 750, LV MCPA 500, Ken-Gran 750 Bromoxynil 200 g/L, 2,4-D Amine 500 Herbicide, Ken-Trel 300, Tordon® 50-D, KA Dicamba 700 (up to 115 g only), Eclipse® Achieve®
Wheat, barley, triticale, and cereal rye only (not undersown)		Token 240
Wheat only (not undersown)		
Clover	Up to 750 mL/ha	Sextant, Fusilade®
Subterranean clover		Simazine (500 g/L), simazine (500 g/L) + paraquat (200 g/L)
	Up to 1.0 L/ha	2,4-DB amine (500 g/L)

When mixing with other herbicides, crop yellowing may be enhanced. When mixing with Ken-Grass 375, Wildcat, Puma Progress or Tristar Advance some reduction in the efficacy and speed of action of these products may occur. If the crop is stressed, the application of the herbicide tank-mixtures may cause yield reduction. When mixing with Cadence a temporary wilting may be evident in some crops after application. Growers should seek advice before spraying recently released cereal varieties. Use the recommended rates for both herbicides in the tank-mixture as well as the surfactant recommendation of the grass herbicide. If another herbicide is applied as a tank mix, observe the plantback restrictions on that label. DO NOT add surfactant when mixing Texus and Ken-Met 600.

Simazine: Refer to the simazine label for correct application rates, especially with regard to soil types.

This product may be mixed in the spray tank with one of the following insecticides according to the directions for use on this product: Hallmark® 50EC, Ken-Tac 100, Karate®, Decis Options®, and Tal-ken 100.

DO NOT use crop oils with Texus or Texus tank mixtures with other products in cereals.

As formulations of other manufacturer's products are beyond the control of Kenso Corporation (M) Sdn Bhd, all mixtures should be tested prior to mixing commercial quantities.

WEEDS LIST

Common name	Scientific name	Common name	Scientific name
Canola (rape)	<i>Brassica napus</i>	Canola (rape)	<i>Brassica napus</i>
Capeweed	<i>Arctotheca calendula</i>	Capeweed	<i>Arctotheca calendula</i>
Charlock	<i>Sinapis arvensis</i>	Charlock	<i>Sinapis arvensis</i>
Chickweed	<i>Stellaria media</i>	Chickweed	<i>Stellaria media</i>
Common sowthistle (milk thistle)	<i>Sonchus oleraceus</i>	Common sowthistle (milk thistle)	<i>Sonchus oleraceus</i>
Corn gromwell	<i>Buglossoides arvense</i>	Corn gromwell	<i>Buglossoides arvense</i>
Cowvine	<i>Ipomoea lonchophylla</i>	Cowvine	<i>Ipomoea lonchophylla</i>
Crassula	<i>Crassula</i> spp.	Crassula	<i>Crassula</i> spp.
Deadnettle	<i>Lamium amplexicaule</i>	Deadnettle	<i>Lamium amplexicaule</i>
Dense-flower fumitory	<i>Fumaria densiflora</i>	Dense-flower fumitory	<i>Fumaria densiflora</i>
Dock	<i>Rumex</i> spp.	Dock	<i>Rumex</i> spp.
Doublegee (spiny emex)	<i>Emex australis</i>	Doublegee (spiny emex)	<i>Emex australis</i>
Fat hen	<i>Chenopodium album</i>	Fat hen	<i>Chenopodium album</i>
Fireweed	<i>Senecio</i> spp.	Fireweed	<i>Senecio</i> spp.
Fumitory	<i>Fumaria</i> spp.	Fumitory	<i>Fumaria</i> spp.
Hedge mustard	<i>Sisymbrium officinale</i>	Hedge mustard	<i>Sisymbrium officinale</i>
Hexham scent (King Island melilot)	<i>Mellilotus indicus</i>	Hexham scent (King Island melilot)	<i>Mellilotus indicus</i>
Horehound	<i>Marrubium vulgare</i>	Horehound	<i>Marrubium vulgare</i>
Hyssop loosestrife	<i>Lythrum hyssopifolia</i>	Hyssop loosestrife	<i>Lythrum hyssopifolia</i>
Iceplant	<i>Mesembryanthemum</i> spp.	Iceplant	<i>Mesembryanthemum</i> spp.
Indian hedge mustard	<i>Sisymbrium orientale</i>	Indian hedge mustard	<i>Sisymbrium orientale</i>
London rocket	<i>Sisymbrium irio</i>	London rocket	<i>Sisymbrium irio</i>
Long storksbill	<i>Erodium botrys</i>	Long storksbill	<i>Erodium botrys</i>
Marshmallow	<i>Malva parviflora</i>	Marshmallow	<i>Malva parviflora</i>
Mouse-eared chickweed	<i>Cerastium glomeratum</i>	Mouse-eared chickweed	<i>Cerastium glomeratum</i>
Night-scented stock	<i>Matthiola longipetala</i>	Night-scented stock	<i>Matthiola longipetala</i>
Paterson's curse	<i>Echium plantagineum</i>	Paterson's curse	<i>Echium plantagineum</i>
Peppercress	<i>Lepidium</i> spp.	Peppercress	<i>Lepidium</i> spp.
Prickly lettuce	<i>Lactuca serriola</i>	Prickly lettuce	<i>Lactuca serriola</i>
Purple goosefoot	<i>Scleroblitum atriplicinum</i>	Purple goosefoot	<i>Scleroblitum atriplicinum</i>
Rough poppy	<i>Papaver hybridum</i>	Rough poppy	<i>Papaver hybridum</i>
Saffron thistle	<i>Carthamus lanatus</i>	Saffron thistle	<i>Carthamus lanatus</i>
Scarlet pimpernel	<i>Anagallis arvensis</i>	Scarlet pimpernel	<i>Anagallis arvensis</i>
Shepherd's purse	<i>Capsella bursa-pastoris</i>	Shepherd's purse	<i>Capsella bursa-pastoris</i>
Skeleton weed	<i>Chondrilla juncea</i>	Skeleton weed	<i>Chondrilla juncea</i>
Sorrel	<i>Rumex</i> spp.	Sorrel	<i>Rumex</i> spp.
Stemless thistle	<i>Onopordum acaulon</i>	Stemless thistle	<i>Onopordum acaulon</i>
Toad rush	<i>Juncus bufonius</i>	Toad rush	<i>Juncus bufonius</i>
Tree hogweed	<i>Polygonum patulum</i>	Tree hogweed	<i>Polygonum patulum</i>
Turnip weed	<i>Rapistrum rugosum</i>	Turnip weed	<i>Rapistrum rugosum</i>
Variegated thistle	<i>Silybum marianum</i>	Variegated thistle	<i>Silybum marianum</i>
Vetch (tares)	<i>Vicia sativa</i>	Vetch (tares)	<i>Vicia sativa</i>
Volunteer lupins	<i>Lupinus</i> spp.	Volunteer lupins	<i>Lupinus</i> spp.
Ward's weed	<i>Carrichtera annua</i>	Ward's weed	<i>Carrichtera annua</i>
Wild radish	<i>Raphanus raphanistrum</i>	Wild radish	<i>Raphanus raphanistrum</i>
Wild turnip	<i>Brassica tournefortii</i>	Wild turnip	<i>Brassica tournefortii</i>
Wireweed (hogweed)	<i>Polygonum aviculare</i>	Wireweed (hogweed)	<i>Polygonum aviculare</i>

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions, or from spraying equipment, that may cause drift onto nearby susceptible plants/crops, cropping lands or pastures. Avoid spray drift and vapour movement onto susceptible crops such as cotton, tobacco, tomatoes, vines, lupins, fruit trees and ornamentals.

PROTECTION OF LIVESTOCK

Grazing Precaution

Sprayed weeds may become more palatable to stock and a higher intake of some weeds may result in stock poisoning and death from causes such as nitrate poisoning. Care should be taken especially where capeweed, Paterson's curse and variegated thistles predominate in the pasture. Avoid grazing with young or breeding stock. Do not graze horses or pigs on Paterson's curse. If in doubt, contact your nearest Department of Agriculture.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Dangerous to fish. DO NOT contaminate streams, rivers or waterways with the chemical or used containers.