

# POISON

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

*KENSO AGCARE*

# POLLUX

## FUNGICIDE

ACTIVE CONSTITUENT: 250 g/L FLUTRIAFOL

**GROUP 3 FUNGICIDE**

For the control of certain fungal diseases on wheat, barley and canola when mixed with fertilizer or applied as a foliar spray as specified in the "Directions for Use" table.



Kenso Corporation (M) Sdn Bhd  
Level 1, 98 Commercial Road,  
Teneriffe QLD 4005  
Phone (07) 3216 1188  
[www.kenso.com.au](http://www.kenso.com.au)

**IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USE**

**CONTENTS: 20 Litres APVMA Approval No.: 60027/ 121695**

# KENSO AGCARE POLLUX FUNGICIDE

## STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Store in a locked room or a place away from children, animals, food, feedstuffs, seed and fertilizers. Triple rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

### 110L Mini Bulk Returnable Container (110L only)

Store the original sealed drum in a cool well-ventilated area. DO NOT store for prolonged periods in direct sunlight. DO NOT tamper with the non-return valve or the security seal. DO NOT contaminate the drum with water or any foreign matter. After each use of the product, please ensure that the non-return valve, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the drum have been used, please return the empty drum to the point of purchase.

## SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes. May irritate the nose and throat and skin. Avoid contact with eyes and skin. When opening the container, preparing the spray, wear cotton overalls buttoned to the neck and wrist and washable hat and elbow-length PVC gloves and face shield and goggles. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, goggles and contaminated clothing.

## FIRST AID

If poisoning occurs, contact a Doctor or Poisons Information Centre (Phone 13 11 26).

## SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet which can be obtained from [www.kenso.com.au](http://www.kenso.com.au).

## CONDITIONS OF SALE

Kenso Corporation (M) Sdn. Bhd. will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Kenso Corporation (M) Sdn. Bhd. is limited to the replacement of the goods or (at the option of Kenso Corporation (M) Sdn. Bhd.) the refund of the price paid and where possible sufficient part of the goods to enable proper examination being returned to Kenso Corporation (M) Sdn. Bhd. within thirty days of delivery.

In a Transport Emergency  
Dial **000** Police or Fire Brigade



9 328666 000945



Additional statements required by Globally Harmonised System of Classification and Labelling of Chemicals (GHS) and Safe Work Australia: **HARMFUL IF SWALLOWED. HARMFUL IF INHALED.** *Precautionary:* Avoid breathing dust/ fume/gas/mist/vapours/spray. Wash contacted area thoroughly after handling. Do not eat, drink or smoke when using this product. Use only outdoors or in a well-ventilated area. IF SWALLOWED: Call a POISON CENTER or doctor/ physician if you feel unwell. IF INHALED: Remove victim to fresh air and keep at rest in a position comfortable for breathing. Call a POISON CENTER or doctor/physician if you feel unwell. Rinse mouth.

Batch No.:

Date of Manufacture:

**POISON**  
KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



Fungicide


ACTIVE CONSTITUENT: 250 g/L FLUTRIAFOL

GROUP **3** FUNGICIDE

For the control of certain fungal diseases on wheat, barley and canola when mixed with fertilizer or applied as a foliar spray as specified in the Directions for Use table.

READ THIS LEAFLET BEFORE USE

APVMA Approval No.: 60027/121695



**SAFETY DIRECTIONS**

Harmful if swallowed. Will irritate the eyes. May irritate the nose and throat and skin. Avoid contact with eyes and skin. When opening the container, preparing the spray, wear cotton overalls buttoned to the neck and wrist and washable hat and elbow-length PVC gloves and face shield and goggles. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, goggles and contaminated clothing.

**FIRST AID**

If poisoning occurs, contact a Doctor or Poisons Information Centre (Phone 13 11 26).

**SAFETY DATA SHEET**

Additional information is listed in the Safety Data Sheet which can be obtained from www.kenso.com.au.

**CONDITIONS OF SALE**

Kenso Corporation (M) Sdn. Bhd. will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Kenso Corporation (M) Sdn. Bhd. is limited to the replacement of the goods or (at the option of Kenso Corporation (M) Sdn. Bhd.) the refund of the price paid and where possible sufficient part of the goods to enable proper examination being returned to Kenso Corporation (M) Sdn. Bhd. within thirty days of delivery.

In a Transport Emergency  
Dial **000**  
Police of Fire Brigade



**Kenso Corporation (M) Sdn. Bhd.**  
Level 1, 98 Commercial Road,  
Teneriffe QLD 4005  
Phone (07) 3216 1188

APVMA Approval No.: 60027/121695

**DIRECTIONS FOR USE:**

**A. Pollux Fungicide as a fertilizer treatment**

Crop	Disease	Rate Per Hectare	WHP (days)	Critical Comments																		
Wheat	Stripe Rust ( <i>Puccinia striiformis</i> )	200 or 400mL	4 weeks grazing	For control of stripe rust of wheat and control of powdery mildew and scald of barley, use the high rate where conditions conducive to severe disease are expected. Application: Spray product evenly onto granulated fertilizer in a mechanical mixer or auger. Apply at a rate which will deposit the specific rate of the product on the quantity of fertilizer required to be sown in furrow per hectare, using the table below as guide. <b>Fertilizer Rate      Volume of Product per 100kg of Fertilizer (mL)</b> <b>Per hectare                      200mL Rate                      400mL Rate</b>																		
	Septoria Tritici blotch ( <i>S. tritici</i> ), take-all ( <i>Geaumannomyces graminis</i> )	400mL																				
Barley	Powdery mildew ( <i>Erysiphe graminis</i> )	200 or 400mL		<table border="1"> <tr> <td>50</td> <td>400</td> <td>800*</td> </tr> <tr> <td>60</td> <td>333</td> <td>667*</td> </tr> <tr> <td>70</td> <td>286</td> <td>571</td> </tr> <tr> <td>80</td> <td>250#</td> <td>500</td> </tr> <tr> <td>100</td> <td>200#</td> <td>400</td> </tr> <tr> <td>120</td> <td>167#</td> <td>333</td> </tr> </table>	50	400	800*	60	333	667*	70	286	571	80	250#	500	100	200#	400	120	167#	333
	50				400	800*																
60	333	667*																				
70	286	571																				
80	250#	500																				
100	200#	400																				
120	167#	333																				
	Scald ( <i>Rhynchosporium secalis</i> )																					

\*Application volumes over 600mL per 100kg of fertilizer may adversely effect the flow rate and handling characteristic of dusty/poorly granulated fertilizers.  
#Where application volume is low, product may be diluted with water to improve coverage.

Crop	Disease	Rate Per Hectare	WHP (days)	Critical Comments										
Canola	Blackleg ( <i>Leptosphaeria macularis</i> )	400mL	4 weeks grazing	Treatment will reduce blackleg infection in both susceptible and tolerant canola varieties. <b>Application:</b> Spray undiluted product evenly onto granulated fertilizer and immediately mix thoroughly in a mechanical mixer or auger. Apply at a rate which will deposit 400mL of the product on the quantity of fertilizer required to be sown in furrow per hectare, using the table below as guide. <b>Fertilizer Rate      Volume of Product per 100kg of Fertilizer (mL)</b> <b>Per hectare                      200mL Rate                      400mL Rate</b>										
				<table border="1"> <tr> <td>60</td> <td>667*</td> </tr> <tr> <td>70</td> <td>571</td> </tr> <tr> <td>80</td> <td>500</td> </tr> <tr> <td>100</td> <td>400</td> </tr> <tr> <td>120</td> <td>333</td> </tr> </table>	60	667*	70	571	80	500	100	400	120	333
60	667*													
70	571													
80	500													
100	400													
120	333													

<b>GROUP</b>	<b>3</b>	<b>FUNGICIDE</b>
--------------	----------	------------------

**Fungicide Resistance Warning**

Pollux Fungicide is a member of the DMI group of fungicides. For fungicide resistance management Pollux Fungicide is a Group 3 Fungicide. Some naturally occurring individual fungi resistant to Pollux Fungicide and other Group 3 Fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by Pollux Fungicide and other Group 3 Fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, Kenso Corporation (M) Sdn. Bhd. accepts no liability for any losses that may result from the failure of Kenso Agcare Pollux Fungicide to control resistant fungi. Resistant should not be assumed without first reviewing the method of application, the coverage and the time of application.

**PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT**

DO NOT contaminate streams, rivers or waterways with the chemical or used container.

**STORAGE AND DISPOSAL**

Store in the closed, original container in a dry, cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Store in a locked room or a place away from children, animals, food, feedstuffs, seed and fertilizers. Triple rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

*110L Mini Bulk Returnable Container (110L only)*

Store the original sealed drum in a cool well-ventilated area. DO NOT store for prolonged periods in direct sunlight. DO NOT tamper with the non-return valve or the security seal. DO NOT contaminate the drum with water or any foreign matter. After each use of the product, please ensure that the non-return valve, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the drum have been used, please return the empty drum to the point of purchase.

**B. Pollux Fungicide Applied as a Foliar Spray**

Crop	Disease	Rate Per Hectare	WHP (days)	Critical Comments
Wheat	Stripe rust ( <i>Puccinia striiformis</i> )	250-500mL <b>Ground Applications:</b> Apply in 100-150mL water plus 200mL "Ken-Wett 1000" per 100L of spray solution. <b>Aerial Application:</b> Apply in a minimum of 15L water plus 200mL "Ken-Wett 1000" per 100L of spray solution.	7 weeks harvest or grazing	Apply when stripe rust appears and before disease reaches 1% of leaf area. The high rate of application gives a longer period of protection than the low rate. In crops with high yield potential and where conditions favourable to severe disease persist, use a single application at the higher rate, or two applications 3-4 weeks apart at the low rate.
	Leaf rust ( <i>Puccinia striiformis</i> ) Septoria nodorum blotch ( <i>Septoria nodorum</i> ) Septoria Tritici blotch ( <i>S. Tritici</i> )			Apply after flag leaves have begun to emerge and before disease reaches 1% of flag leaf area. The high rate of application gives a longer period of protection than the low rate. In crops with high yield potential, and where conditions favourable to severe diseases persist during flowering, use a single application at the higher rate or two applications 3-4 weeks apart at the low rate.
Barley	Powdery mildew ( <i>Erysiphe graminis</i> )		10 weeks harvest or grazing	Apply when powdery mildew first appears. The high rate of application gives a longer period of protection than the low rate. Where conditions favourable to severe disease persist, a single application at the higher rate, or two applications 3-4 weeks apart at the low rate.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**WITHHOLDING PERIODS:**

**A. Pollux Fungicide Applied in-furrow as a fertilizer**

**TREATMENT:**

**CANOLA, WHEAT and Barley: DO NOT GRAZE OR CUT FOR STOCK FOR 4 WEEKS AFTER SOWING.**

**B. Pollux Fungicide Applied as a Foliar Spray**

**DO NOT HARVEST EARLIER THAN THE NUMBER OF WEEKS SHOWN AFTER APPLICATION.**

**BARLEY -10 WEEKS**

**WHEAT – 7 WEEKS**

**DO NOT GRAZE OR CUT FOR STOCK FOOD EARLIER THAN THE NUMBER OF WEEKS SHOWN AFTER APPLICATION.**

**BARLEY – 10 WEEKS**

**WHEAT – 7 WEEKS**

**GENERAL INSTRUCTIONS**

**Fertilizer Application:**

Ensure that treated fertilizer is dry before sowing with the crop.

Sowing: Treated fertilizer should be sown in the same furrow as the seed, not broadcast. Note that the flow rate of fertilizer may be changed after treatment, and the seeder should be calibrated with the treated material to ensure that the corrected rates of fertilizer and fungicide are delivered. DO NOT mix seed with treated fertilizer.

**Foliar Application:**

Check the crop carefully and regularly. Spray only if there is sufficient time in the season for the rust to develop to a level on the variety in question when it will cause significant yield loss.