

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

KENSO AGCARE

KENTROLE 250

HERBICIDE

ACTIVE CONSTITUENTS: 250 g/L AMITROLE
220 g/L AMMONIUM THIOCYANATE

GROUP **34** HERBICIDE

For the Control of Weeds in Orchards, Vineyards, Potatoes, Pine plantations,
Irrigation Ditches and Drains, Roadsides, Wheat and Barley, and for General
Industrial Situations as per the Directions for Use.

Kenso Corporation (M) Sdn Bhd
Level 1, 98 Commercial Road,
Teneriffe QLD 4005
Phone (07) 3216 1188
www.kenso.com.au



IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USE

CONTENTS: 20 Litres APVMA Approval No.: 92432/135489

KENSO AGCARE KENTROLE 250 HERBICIDE

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight.

Triple-rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush, or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

Refillable containers:

Store the closed original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Do not inhale spray mist. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet, which can be obtained from www.kenso.com.au.

CONDITIONS OF SALE

Kenso Corporation (M) Sdn. Bhd. will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection

with the supply of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Kenso Corporation (M) Sdn. Bhd. is limited to the replacement of the goods or (at the option of Kenso Corporation (M) Sdn. Bhd.) the refund of the price paid and where possible sufficient part of the goods to enable proper examination being returned to Kenso Corporation (M) Sdn. Bhd. within thirty days of delivery.

In a Transport Emergency
Dial **000**
Police or Fire Brigade



9 328666 003991



Additional statements required by Globally Harmonised System of Classification and Labelling of Chemicals (GHS) and Safe Work Australia: **HARMFUL IF SWALLOWED. HARMFUL IN CONTACT WITH SKIN. SUSPECTED OF DAMAGING FERTILITY OR THE UNBORN CHILD. MAY CAUSE DAMAGE TO ORGANS THROUGH PROLONGED OR REPEATED EXPOSURE.** *Precautionary:* Obtain special instruction before use. Do not handle until all safety precautions have been read and understood. Do not breathe dust/fume/gas/mist/vapours/spray. Wash contacted areas thoroughly after handling. Do not eat, drink or smoke when using this product. Wear protective gloves/protective clothing/eye protection/face protection. IF SWALLOWED: Call a POISON CENTRE or doctor/physician if you feel unwell. IF ON SKIN: Wash with plenty of soap and water. IF exposed or concerned: Get medical advice/attention. Specific treatment (see FIRST AID on this label). Rinse mouth. Take off contaminated clothing and wash before reuse. Store locked up.

Batch No.:

Date of Manufacture:

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

KENSO
agcare

KENTROLE 250

Herbicide

ACTIVE CONSTITUENT: 250 g/L AMITROLE
220 g/L AMMONIUM THIOCYANATE

GROUP **34** HERBICIDE

For the Control of Weeds in Orchards, Vineyards, Potatoes, Pine plantations, Irrigation Ditches and Drains, Roadsides, Wheat and Barley, and for General Industrial Situations as per the Directions for Use

READ THIS LEAFLET BEFORE USE

APVMA Approval No.: 92432/135489



Kenso Corporation (M) Sdn. Bhd.
Level 1, 98 Commercial Road,
Teneriffe QLD 4005
Phone (07) 3216 1188

GENERAL INSTRUCTIONS

Before opening carefully read Safety Directions, First Aid Instructions, Directions for Use, Storage and Disposal and Protection Statements.

APPLICATION INFORMATION

Thoroughly wet all surfaces by aiming to apply 1700 - 2200 L/ha with power spraying equipment, 900 - 1100 L/ha (1 litre per 10m²) using a knapsack and 170 L/ha when using a misting machine.

Note: Misting machines are not recommended for use in Tasmania.

MIXING INSTRUCTIONS

To improve the efficiency of this product it is recommended that a surfactant be added to the spray mixture at the following rates:

Power Spraying: 250 mL/100 L spray mixture

Knapsack Spraying: 5 mL/15 L knapsack

Misting Machine: 60 mL/15 L spray mixture (Not recommended in Tasmania)

Note: The rates indicated in the Directions for Use table are per 100 L of water for power spraying. Refer to the chart below for the required amounts to suit other equipment.

CONVERSION CHART FOR RECOMMENDATIONS TO SUIT VARIOUS EQUIPMENT

Recommended Rate per 100 L of water	Per 15L of knapsack	Per 5L Misting
4.5 L	1.1 L	1.3 L
2.2 L	570 mL	630 mL
2.0 L	510 mL	560 mL
1.7 L	430 mL	480 mL
1.1 L	280 mL	310 mL
570 mL	110 mL	120 mL
280 mL	55 mL	60 mL

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

Drift Warning: This product is a non-selective herbicide and may damage crops, pastures and some ornamentals. However, it is safer to use near plants susceptible to phenoxy type herbicides. DO NOT apply under weather conditions or from spraying equipment that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

Equipment Maintenance and usage: DO NOT use the spray machine for any other purpose unless thoroughly cleaned by several rinsings of all parts with clean water.

SAFETY DIRECTIONS

DO NOT inhale spray mist. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet, which can be obtained from www.kenso.com.au.

CONDITIONS OF SALE

Kenso Corporation (M) Sdn. Bhd. will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Kenso Corporation (M) Sdn. Bhd. is limited to the replacement of the goods or (at the option of Kenso Corporation (M) Sdn. Bhd.) the refund of the price paid and where possible sufficient part of the goods to enable proper examination being returned to Kenso Corporation (M) Sdn. Bhd. within thirty days of delivery.

In a Transport Emergency
Dial **000**
Police of Fire Brigade



Kenso Corporation (M) Sdn. Bhd.
Level 1, 98 Commercial Road,
Teneriffe QLD 4005
Phone (07) 3216 1188

APVMA Approval No.: 92432/135489

RESISTANT WEEDS WARNING:

GROUP **34** HERBICIDE

Kenso Agcare Kentrole 250 Herbicide is a member of the triazole group of herbicides. Kenso Agcare Kentrole 250 Herbicide has the inhibition of lycopene cyclase mode of action. For weed resistance management Kenso Agcare Kentrole 250 Herbicide is a Group 34 herbicide. Some naturally occurring weed biotypes resistant to Kenso Agcare Kentrole 250 Herbicide and other Group 34 herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. The effectiveness of Kenso Agcare Kentrole 250 Herbicide on resistant individuals could be significantly reduced. Since the occurrence of resistant weeds is difficult to detect prior to use, Kenso Corporation (M) Sdn. Bhd. accepts no liability for any losses that may result from the failure of Kenso Agcare Kentrole 250 Herbicide to control resistant weeds. Kenso Agcare Kentrole 250 Herbicide may be subject to specific resistance strategies. For further information contact your local supplier, Kenso Corporation (M) Sdn. Bhd. representative or local Department of Agriculture agronomist.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight.

Triple-rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. Do not burn empty containers or product.

For refillable containers (1000L)

Store the closed original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

DIRECTION FOR USE

Crop / Situation	Weeds Controlled	State	Rate / ha	Critical Comments
Vineyards and Orchards	Couch Grass	All states	1.1 L/100 L water	Apply when weeds are small and actively growing. Repeat application in 6 - 8 weeks. Apply as a directed spray to weeds only.
	Broadleaf Weeds and Grasses		570 mL/100 L water	
Aquatic Areas Drains and Channels, Margins of Streams, Lakes and Dams	Cumbungi, Phragmites and Nutgrass		2.3 L/100 L water	Apply during flowering period between January and May. Re-treatment of Nutgrass will be necessary.
	Water Couch		4.5 L/100 L water 2.2 L/100 L water	Apply as a single treatment between February and May. Apply in November to December with a further application at the same rate 6 – 8 weeks later.
	Water Hyacinth		280 mL/100 L water	Apply immediately prior to flowering.
Pinus radiata Plantations	Rye Grass, Wild Oats, Yorkshire Fog Grass, Silver Grass, Sand Brome, Capeweed, Clovers, Mouse-ear Chickweed, Sowthistle and Spear Thistle seedlings	NSW, SA, WA, Tas, Vic only	5.6 – 8 L/ha plus 2.8 – 4 L/ha flowable atrazine (500 g/L)	Use highest rates where grasses are a problem. Spray evenly. Pre-planting: Graze area heavily and apply the tank mixture not more than 2 weeks before planting. Post-planting: Apply tank mixture immediately after or within one month of planting. DO NOT spray pine seedlings by using a directed spray.
Non-crop areas around Buildings, Commercial and Industrial areas, Domestic and Public Service areas, Right- of- Way	Blackberry	Vic, Tas, SA only	2 L/100 L water	Apply twice. First treatment in December or January and regrowth sprayed in autumn. Respraying in the following year is essential. DO NOT apply to blackberry bushes carrying mature or developing fruit.
	Furze or Gorse, Ox- eye Daisy			Apply during summer or before full flowering. Respraying will be necessary.
	Fennel, Hoary Cress, African Daisy, Apple of Sodom, Illyrian Thistle			Apply when weeds are actively growing, immediately prior to flowering.
	Russian (creeping) Knapweed, Watsonia, Artichoke Thistle, Prairie Ground Cherry, Onion Weed, Khaki Weed, Chilean Cestrum (Green Cestrum), Oxalis (Soursob), Couch, Kikuyu, Fog Grass, Sorrel	All states	1.1 L/100 L water	Apply when weeds are actively growing, immediately prior to flowering. Respraying will be necessary to destroy regrowth and seedlings.
Potatoes Pre-Harvest Preparation	Rye Grass, Winter Grass, Capeweed, Paterson's Curse	Vic, Tas only	5.5 - 11 L	Apply approximately 4 - 5 weeks before harvest after the potato haulms have dried off. Use the higher rates when rye grass is in clumps or other weeds more than 30 cm high are present. For boom spraying apply at least 100 litres of spray mixture per hectare. If other broadleaf weeds such as Clovers, Variegated Thistle and Cruciferous weeds are also present add a low volatile ester 400 2,4-D herbicide. Warning: If used on sandy soils do not plant follow-up crops within 3 months.
Wheat and Barley	Rye Grass, Wild Oats	All states	2.8 L 5.6 L	Pre-planting. Sow the crop at a depth of 5 cm not less than 5 days after spraying. A mechanical cultivation 4 to 5 days after spraying will assist results. Apply when grasses are in the 2 - 3 leaf stage and 5 cm high. Apply when grasses are in the 6-leaf stage and 10 cm high. DO NOT use on sandy soils in cropping situations.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WITHHOLDING PERIOD

ORCHARDS AND VINES: DO NOT APPLY LATER THAN 56 DAYS BEFORE HARVEST
OTHER CROPS: NOT REQUIRED WHEN USED AS DIRECTED