

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

KENSO AGCARE

**DICAMBA
M**

SELECTIVE HERBICIDE

ACTIVE CONSTITUENTS: 340 g/L MCPA (present as the dimethylamine salt)
+ 80 g/L DICAMBA (present as the dimethylamine salt)

GROUP 4 HERBICIDE

For the control of certain broadleaf weeds in winter cereals,
pastures, turf and non-crop areas.

Kenso Corporation (M) Sdn Bhd
Level 1, 98 Commercial Road,
Teneriffe QLD 4005
Phone (07) 3216 1188
www.kenso.com.au



IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USE

CONTENTS: 20 Litres APVMA Approval No.: 92191/ 134805

KENSO AGCARE DICAMBA M SELECTIVE HERBICIDE

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well ventilated area away from children, animals, food, feedstuffs, seed and fertilizers. DO NOT store for prolonged periods in direct sunlight. Triple rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available bury the empty container 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. Do not burn empty containers or product.

SAFETY DIRECTIONS

Harmful if absorbed by skin contact, inhaled or swallowed. Will irritate the eyes and skin. DO NOT inhale vapour. Repeated exposure may cause allergic disorders. When opening the container and preparing the spray wear cotton overalls buttoned to the neck and wrist, washable hat, elbow length PVC gloves, face shield or goggles. When using the prepared spray wear cotton overalls buttoned to the neck and wrist, washable hat and elbow length PVC gloves. If product in eyes, wash out immediately with water. If product on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet, which can be obtained from www.kenso.com.au.

CONDITIONS OF SALE

Kenso Corporation (M) Sdn. Bhd. will not accept any responsibility

whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Kenso Corporation (M) Sdn. Bhd. is limited to the replacement of the goods or (at the option of Kenso Corporation (M) Sdn. Bhd.) the refund of the price paid and where possible sufficient part of the goods to enable proper examination being returned to Kenso Corporation (M) Sdn. Bhd. within thirty days of delivery.

In a Transport Emergency
Dial **000** Police or Fire Brigade



Additional statements required by Globally Harmonised System of Classification and Labelling of Chemicals (GHS) and Safe Work Australia: **HARMFUL IF SWALLOWED. CAUSES SKIN IRRITATION. CAUSES SERIOUS EYE DAMAGE.** *Precautionary:* Wash contacted areas thoroughly after handling. Do not, drink or smoke when using this product. Wear protective gloves, protective clothing and eye or face protection. **IF SWALLOWED:** Call a POISON CENTER or doctor/physician if you feel unwell. **IF ON SKIN:** Wash with plenty of soap and water. **IF ON EYES:** Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. Immediately call a POISON CENTER or doctor/physician. Specific treatment (see FIRST AID on this label). Rinse mouth. If skin irritation occurs: Get medical advice/attention. Take off contaminated clothing and wash it before reuse.

Batch No.:

Date of Manufacture:

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

KENSO
agcare

DICAMBA M

Selective Herbicide

ACTIVE CONSTITUENTS: 340 g/L MCPA (present as the dimethylamine salt) + 80 g/L DICAMBA (present as the dimethylamine salt)

GROUP 4 HERBICIDE

For the control of certain broadleaf weeds in winter cereals, pastures, turf and non-crop areas.

READ THIS LEAFLET BEFORE USE

APVMA Approval No.: 92191/134805



Kenso Corporation (M) Sdn. Bhd.
Level 1, 98 Commercial Road,
Teneriffe QLD 4005
Phone (07) 3216 1188

GENERAL INSTRUCTIONS

RESISTANCE WEEDS WARNING

GROUP 4 HERBICIDE

Kenso Agcare Dicamba M Selective Herbicide ("Dicamba M") contains members of the Benzoic acid and Phenoxy groups of herbicides. Dicamba M has the disruptors of plant cell growth mode of action. For weed resistance management Dicamba M is a Group 4 Herbicide. Some naturally-occurring weed biotypes resistant to Dicamba M and other Group 4 herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Dicamba M or any other Group 4 Herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Kenso Agcare Pty Ltd accepts no liability for any losses that may result from the failure of Dicamba M to control resistant weeds.

MIXING

Add the required amount of Dicamba M directly to the tank and mix well.

APPLICATION

For Boom Application: Apply by a properly calibrated boom spray using not less than 50L of water per hectare unless indicated otherwise in the Critical Comments column.

For Aerial Application: Apply at least 25L total spray volume per hectare. The aircraft should fly as low as practical under the prevailing conditions to minimise drift.

For High Volume Application: Apply minimum 1500L per hectare.

For Knapsack Application: A 15L knapsack treats 150 square metres.

NOTE: Spray when weeds are young and actively growing. A temporary wilting may be evident in crops after application. If seasonal conditions are unusually late or dry seek further advice as damage may result from spraying under these conditions. Growers should seek advice before spraying recently released cereal varieties. There is a nil tolerance of pea seed contaminates in cereal grain and less than 100% control of Volunteer Peas could result in dockage.

COMPATIBILITY

Dicamba M is compatible with Terbutryn 500g/L SC.

SAFETY DIRECTIONS

Harmful if absorbed by skin contact, inhaled or swallowed. Will irritate the eyes and skin. DO NOT inhale vapour. Repeated exposure may cause allergic disorders. When opening the container and preparing the spray wear cotton overalls buttoned to the neck and wrist, washable hat, elbow length PVC gloves, face shield or goggles. When using the prepared spray wear cotton overalls buttoned to the neck and wrist, washable hat and elbow length PVC gloves. If product in eyes, wash out immediately with water. If product on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet, which can be obtained from www.kenso.com.au.

CONDITIONS OF SALE

Kenso Corporation (M) Sdn. Bhd. will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Kenso Corporation (M) Sdn. Bhd. is limited to the replacement of the goods or (at the option of Kenso Corporation (M) Sdn. Bhd.) the refund of the price paid and where possible sufficient part of the goods to enable proper examination being returned to Kenso Corporation (M) Sdn. Bhd. within thirty days of delivery.

In a Transport Emergency
Dial **000**
Police of Fire Brigade



Kenso Corporation (M) Sdn. Bhd.
Level 1, 98 Commercial Road,
Teneriffe QLD 4005
Phone (07) 3216 1188

APVMA Approval No.: 92191/134805

WITHHOLDING PERIOD:

DO NOT GRAZE OR CUT FOR STOCK FOOD 7 DAYS AFTER APPLICATION.

HARVEST WITHHOLDING PERIOD:

NOT REQUIRED WHEN USED AS DIRECTED.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

Spray equipment must be calibrated accurately prior to working on crops. Minimise spray drift by using low pressure and nozzles which do not give a fine droplet size. DO NOT use in high wind. DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures. Avoid spray drift onto susceptible crops such as cotton, tobacco, tomatoes, vines, lupins, fruit trees and ornamentals. DO NOT apply this product on or near desirable trees or plants or in locations where the chemical may be washed or moved into contact with their roots. Keep container sealed (airtight) when not in use. All spray equipment must be thoroughly washed out after use and must not be used for spraying horticulture crops.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well ventilated area away from children, animals, food, feedstuffs, seed and fertilizers. DO NOT store for prolonged periods in direct sunlight. Triple rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available bury the empty container 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. Do not burn empty containers or product.

RESTRAINTS

- DO NOT spray when rain seems likely to occur within 4 hours.
- DO NOT spray when weeds are wet with dew or rain or under stress from drought, low soil fertility, extreme cold or water logging.
- DO NOT spray outside crop growth stages recommended as crop damage may result.
- DO NOT apply to crops or grass pastures undersown with clover, lucerne or medics.
- DO NOT apply to legume pastures.
- DO NOT apply to plants suffering from stress.

SPRAY DRIFT RESTRAINTS

- Specific definitions for terms used in this section of the label can be found at apvma.gov.au/spraydrift
- DO NOT allow bystander to come into contact with the spray cloud.
- DO NOT apply in a manner that may cause an unacceptable impact to native vegetation, agricultural crops, landscaped gardens and aquaculture production, or cause contamination of plant or livestock commodities, outside the application site from spray drift.
- Wherever possible, correctly use application equipment designed to reduce spray drift and apply when the wind direction is away from these sensitive areas.
- DO NOT apply unless the wind speed is between 3 and 20 kilometres per hour at the application site during the time of application.
- DO NOT apply if there are hazardous surface temperature inversion conditions present at the application site during the time of application. Surface temperature inversion conditions exist most evenings one to two hours before sunset and persist until one to two hours after sunrise.

DIRECTIONS FOR USE

CROP & SITUATION	WEEDS	STATE	RATE	CRITICAL COMMENTS
Barley, Commercial Rye, Oats, Triticale, Wheat From early tillering stage (Z21)	Capeweed, Charlock, Common ice-plant, seedling Docks, Doublegee, (Threecornered Jack, Spiny emex), Hexham scent (Melilotus), Mustards, Soldier thistle, Sorrel seedlings, Tares, Tree hogweed, Turnip weed (<i>Rapistrum rugosum</i>), Vetches, Ward's weed, Wild radish, Wild turnip (<i>Brassica tournefortii</i>), Wireweed (Hogweed)	All states	1 L/ha	Spray when most weeds are in the 2-4 leaf stage. In some instances suppression only of wireweed (hogweed), capeweed and wild radish will occur.
	Seedling Cotton thistle, Fat hen	Tas only		
Barley, Commercial Rye, Oats, Triticale, Wheat From early tillering stage (Z21) until the fully tillered stage and before jointing occurs (Z30)	Cotton thistle to bud formation	Tas only	1.7 L/ha	Spray when most weeds are still in the rosette or seedling stage. N.B. Wild radish may need to be resprayed (with 2,4-D or MCPA) if future germinations occur after spraying. In some instances, suppression only of Wireweed (hogweed), Hoary cress and Climbing buckwheat will be obtained.
	Mintweed, New Zealand spinach	Qld, NSW only		
	Capeweed, Charlock, Chickweed, Climbing buckwheat (Black bindweed), Clover, Common ice-plant, seedling Docks, Doublegee (Threecornered Jack, Spiny emex), Fat hen, Hexham scent (Melilotus), seedling Hoary cress, Mouse-eared Chickweed, Mustards, Narrow-leaved plantain, Saffron thistle, Spurry, Swinecress, Tares, Tree hogweed, Turnip weed (<i>Rapistrum rugosum</i>), Variegated thistle, Vetch, Wild radish, Wild turnip (<i>Brassica tournefortii</i>), Wireweed (Hogweed)	Qld, NSW Tas, WA, Vic only	1.4 L/ha	
Barley, Commercial Rye, Triticale From early tillering stage (Z21) until the fully tillered stage before jointing occurs (Z30)	Volunteer peas	SA only	1 L/ha plus 60 mL/ha Dicamba 500	Apply when most peas are 5-7 cm high.
Oats, Wheat From early tillering stage (Z21) until the fully tillered stage and before jointing occurs (Z30)			1L/ha plus 120 mL/ha Dicamba 500	

CROP & SITUATION	WEEDS	STATE	RATE			CRITICAL COMMENTS
			Per Hectare	Per 100L	Per 15L Knapsack	
Grass pastures	Bathurst burr, Caltrop (Yellow vine), Chicory, Crofton weed, Cudweed, Dandelion, Deadnettle, Fleabanes, Fumitory, Groundsel, Heliotrope, Inkweed, Mintweed, Nettles, Noogoora burr, Pennycress, Potato weed, Ragwort, Ripplewort, Shepherd's purse, Smartweed, Soft rush, Stinkwort, Thistles including Cockspur, Scotch (Spear), Slender, Star, Variegated, Wireweed (Hogweed)	All states	2.8-4L	190-270mL	60mL	Use the higher rate on larger weeds.
	Cotton thistle	Tas only				
	Buttercups, Hoary cress seedlings, Horehound, Khaki weed, Purple-top, Veined verbena	All states	4-7L	470mL	100mL	When necessary a second treatment may be made after 4 weeks.
Turf	Bindy-eye, Capeweed, Catsear, Chickweed, Clovers, Cotula, Creeping oxalis, Cudweeds, Dandelion, Dock, Fleabanes, Fumitory, Jo-Jo (Onhunga), Lamb's tongue, Medics, Mustards, Narrow-leaved plantain, Pearlwort, Pennywort, Peppergrass, Radish, Scarlet pimpernel, Slender thistle, Soldier thistle, Spear thistle, Toad rush, Variegated thistle, Wild turnip, Wireweed (Hogweed).	All states	Golf Tee, Golf Green: 100mL in 15L water per 150m ² Bowling Green: 900mL in 200L water per 1400m ² Large Turf Area: (Field, Park, Golf Fairway) 6.5L in 250-400L water per 10,000m ² (1 hectare)			Apply to actively growing weeds and moist soil. DO NOT use on Lippia, Strawberry clover or Buffalo grass. DO NOT mow turf for two days before or after application or fertilize within two weeks of spraying.
	Fat hen, Purple-top, Red flowered mallow, Staggerweed, Swinecress	Qld, NSW, Tas, SA, WA only				
Non-Crop Situations	Artichoke, Bindweed, Fennel, Hemlock, Thistle seedlings	All states	5.2L	350mL	80mL	A repeat spray may be necessary.

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.