

# POISON

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

*KENSO AGCARE*

# BROMOXYNIL 200

**SELECTIVE HERBICIDE**

ACTIVE CONSTITUENT: 200 g/L BROMOXYNIL present as the N-OCTANOYL ESTER  
SOLVENT: 616 g/L LIQUID HYDROCARBON

**GROUP C HERBICIDE**

For the control of weeds in wheat, oats, barley, cereal rye and triticale (alone or undersown), linseed, clover and lucerne pastures, turf, fallow, non-crop, roadsides and rights-of-way as per the Directions for Use table.

Kenso Corporation (M) Sdn Bhd  
Level 1, 98 Commercial Road,  
Teneriffe QLD 4005  
Phone (07) 3216 1188  
[www.kenso.com.au](http://www.kenso.com.au)



**IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USE**

**CONTENTS: 20 Litres APVMA Approval No.: 66395/ 53558**

# KENSO AGCARE BROMOXYNIL 200 SELECTIVE HERBICIDE

## STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight.

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If not available bury the containers below 500 mm in a disposal pit specifically set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

### 110L Mini Bulk Returnable Container (110L only)

Store the original sealed drum in a cool well-ventilated area. DO NOT store for prolonged periods in direct sunlight. DO NOT tamper with the non-return valve or the security seal. DO NOT contaminate the drum with water or any foreign matter. After each use of the product, please ensure that the non-return valve, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the drum have been used, please return the empty drum to the point of purchase.

### Refillable containers (1000L only):

Store in the closed, original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight. Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

## SAFETY DIRECTIONS

Product is poisonous if inhaled or swallowed. May irritate the eyes and skin. Avoid contact with eyes and skin. DO NOT inhale spray mist. When preparing spray wear elbow-length PVC gloves and face shield. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and face shield or goggles and contaminated clothing.

## FIRST AID

If poisoning occurs, contact a Doctor or Poisons Information Centre (Phone Australia 13 11 26). If swallowed, DO NOT induce vomiting. Give a glass of water.

## SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet, which can be obtained from [www.kenso.com.au](http://www.kenso.com.au).

## CONDITIONS OF SALE

Kenso Corporation (M) Sdn. Bhd. will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Kenso Corporation (M) Sdn. Bhd. is limited to the replacement of the goods or (at the option of Kenso Corporation (M) Sdn. Bhd.) the refund of the price paid and where possible sufficient part of the goods to enable proper examination being returned to Kenso Corporation (M) Sdn. Bhd. within thirty days of delivery.

In a Transport Emergency  
Dial **000** Police or Fire Brigade



Additional statements required by Globally Harmonised System of Classification and Labelling of Chemicals (GHS) and Safe Work Australia: **COMBUSTIBLE LIQUID. HARMFUL IF SWALLOWED. MAY CAUSE AN ALLERGIC SKIN REACTION. TOXIC IF INHALED. SUSPECTED OF DAMAGING THE UNBORN CHILD.** *Precautionary:* Obtain special instructions before use. Do not handle until all safety precautions have been read and understood. Keep away from heat/sparks/open flames/hot surfaces — No smoking. Avoid breathing dust/fume/gas/mist/vapours/spray. Wash contacted areas thoroughly after handling. Do not eat, drink or smoke when using this product. Use only outdoors or in a well-ventilated area. Contaminated work clothing should not be allowed out of the workplace. Wear protective gloves/ protective clothing/ eye protection/ face protection. IF SWALLOWED: Call a POISON CENTER or doctor/physician if you feel unwell. IF ON SKIN: Wash with plenty of soap and water. IF INHALED: Remove victim to fresh air and keep at rest in a position comfortable for breathing. IF exposed or concerned: Get medical advice/attention. Call a POISON CENTER or doctor/physician. Specific treatment (see FIRST AID on this label). Rinse mouth. If skin irritation or rash occurs: Get medical advice/attention. Wash contaminated clothing before reuse. In case of fire: Use water fog, foam, dry agent (carbon dioxide, dry chemical powder) for extinction. Store in a well-ventilated place. Keep container tightly closed. Keep cool. Store locked up.

Batch No.:

Date of Manufacture:

**POISON**  
**KEEP OUT OF REACH OF CHILDREN**  
**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

**KENSO**  
**BROMOX-  
 YNIL200**  
**Selective Herbicide**

**ACTIVE CONSTITUENT: 200g/L BROMOXYNIL present as the N-OCTANOYL ESTER**  
**SOLVENT: 616g/L LIQUID HYDROCARBON**

**GROUP C HERBICIDE**

For the control of weeds in wheat, oats, barley, cereal rye and triticale (alone or undersown), linseed, clover and lucerne pastures, turf, fallow, non-crop, roadsides and rights-of-way as per the Directions for Use table.

**READ THIS LEAFLET BEFORE USE**

**APVMA Approval No.: 66395/53558**



**Kenso Corporation (M) Sdn. Bhd.**  
 Level 1, 98 Commercial Road,  
 Teneriffe QLD 4005  
 Phone (07) 3216 1188

**STORAGE AND DISPOSAL**

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If not available bury the containers below 500 mm in a disposal pit specifically set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

**110L Mini Bulk Returnable Container**

Store the original sealed drum in a cool well-ventilated area. DO NOT store for prolonged periods in direct sunlight. DO NOT tamper with the non-return valve or the security seal. DO NOT contaminate the drum with water or any foreign matter. After each use of the product, please ensure that the non-return valve, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the drum have been used, please return the empty drum to the point of purchase.

**Refillable containers (1000L only):**

Store in the closed, original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight. Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

**SAFETY DIRECTIONS**

Product is poisonous if inhaled or swallowed. May irritate the eyes and skin. Avoid contact with eyes and skin. DO NOT inhale spray mist. When preparing spray wear elbow-length PVC gloves and face shield. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and face shield or goggles and contaminated clothing.

**FIRST AID**

If poisoning occurs, contact a Doctor or Poisons Information Centre (Phone Australia 13 11 26). If swallowed, DO NOT induce vomiting. Give a glass of water.

**SAFETY DATA SHEET**

Additional information is listed in the Safety Data Sheet which can be obtained from the supplier.

**CONDITIONS OF SALE**

Kenso Corporation (M) Sdn. Bhd. will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Kenso Corporation (M) Sdn. Bhd. is limited to the replacement of the goods or (at the option of Kenso Corporation (M) Sdn. Bhd.) the refund of the price paid and where possible sufficient part of the goods to enable proper examination being returned to Kenso Corporation (M) Sdn. Bhd. within thirty days of delivery.

® is not a registered trademarks of Kenso Corporation (M) Sdn. Bhd.

In a Transport Emergency  
 Dial **000**  
 Police of Fire Brigade



**Kenso Corporation (M) Sdn. Bhd.**  
 Level 1, 98 Commercial Road,  
 Teneriffe QLD 4005  
 Phone (07) 3216 1188

**APVMA Approval No.: 66395/53558**

**DIRECTIONS FOR USE:**

**RESTRAINTS:** DO NOT apply by mister machines. DO NOT apply if crops or weeds are stressed due to excessively dry or moist conditions. DO NOT apply to diseased or frost affected crops or turf or if frosts are imminent. DO NOT apply when rain is expected within 3 hours. DO NOT apply to lucerne crops or clover based pastures or turf in temperatures above 20°C or when temperatures above 20°C may follow for some days after application. DO NOT apply when windspeed is less than 3 or more than 20km per hour at application site. DO NOT apply during surface temperature inversion conditions at application site. DO NOT apply with smaller than COARSE spray quality according to ASAE S572 definition for standard nozzles.

SITUATION AND CROP	WEEDS CONTROLLED	WEED STAGE	STATE	RATE	CRITICAL COMMENTS
Wheat, Barley, Cereal rye, Oats and Triticale: including those undersown with Clover, Lucerne or Medic (Vic only); Linseed; Clover or Lucerne Pastures; Lotus seed or forage crops (Qld only)	Amsinckia ( <i>Amsinckia</i> spp.), Black bindweed (Climbing buckwheat) ( <i>Fallopia convolvulus</i> ), Capeweed ( <i>Arctotheca calendula</i> ), Chamomile ( <i>Matricaria matricarioides</i> ), Common peppercress ( <i>Lepidium africanum</i> ), Corn gromwell (Ironweed, Sheepweed) ( <i>Buglossoides arvensis</i> ), Fat hen ( <i>Chenopodium album</i> ), Field madder ( <i>Sherardia arvensis</i> ), Lesser swinecress ( <i>Coronopus didymus</i> ), Mountain sorrel ( <i>Oxalis acetosella</i> ), Purple calandrinia ( <i>Calandrinia menziesii</i> ), Saffron thistle ( <i>Carthamus lanatus</i> ), Shepherd's purse ( <i>Capsella bursa-pastoris</i> ), Three-horned bedstraw ( <i>Gallium tricomutum</i> ), Tree hogweed ( <i>Polygonum patulum</i> ), Variegated thistle ( <i>Silybum marianum</i> )	Up to the 4 leaf stage or when plants are no more than 35mm in diameter	All States	1.4L/ha	<b>CROP STAGE:</b> <b>Wheat, Barley, Cereal, Rye, Oats and Triticale:</b> 3 leaf to full tillered (Z13-30). <b>Linseed:</b> 50 to 150mm high. <b>Lucerne and Clover:</b> After seedlings have one trifoliolate leaf or before there is 150mm regrowth in established stands. <b>Lotus:</b> Apply post-emergence at least 4 weeks after sowing (crop seedlings with a minimum of 2-6 true leaves). <b>APPLICATION:</b> Refer to General Instructions. Avoid application when maximum daily temperatures above 20°C occur, or are likely to occur for a few days after application. Good coverage of the spray on the weed is essential. Apply only in the Autumn and Winter (mid-April to end of August). For optimum control, apply only when the weeds are actively growing and before the weeds are shielded by the crop.
		Up to the 6 leaf stage or when plants are no more than 50mm in diameter		2L/ha	
Wheat, Barley, Cereal rye, Oats and Triticale: including those undersown with Clover, Lucerne or Medic (Vic only); Linseed; Clover or Lucerne Pastures; Lotus seed or forage crops (Qld only)	Mexican poppy ( <i>Argemone ochroleuca</i> ), Three cornered Jack (Doublegee, Spiny emex) ( <i>Emex australis</i> ), Turnip weed (Giant mustard) ( <i>Rapistrum rugosum</i> ), Wild radish ( <i>Raphanus raphanistrum</i> ), Wild mustard ( <i>Sisymbrium</i> spp.), Wild turnip ( <i>Brassica tournefortii</i> ), Wireweed## ( <i>Polygonum aviculare</i> ), Paterson's curse# (Salvation Jane) ( <i>Echium plantagineum</i> )	Up to the 4 leaf stage or when plants are no more than 50mm in diameter	Qld, NSW, Vic, SA, WA, ACT only	2L/ha	<b>CAUTION:</b> <b>Lucerne and Clover:</b> This product will cause slight leaf burning of undersown lucerne and clover. DO NOT apply to Persian clover or Berseem clover. <b>Linseed:</b> DO NOT spray linseed by aerial application. <b>Lotus:</b> Some initial scorching of leaves may occur in some cases. <b>Medics:</b> Application to some cultivars of Medics may result in unacceptable crop damage in some situations. # Effective control of Paterson's curse and Fumitories will only be obtained under good growing conditions. ## Where Wireweed is growing under stress, it has been found to be less susceptible.
	Pheasant's eye ( <i>Adonis microcarpa</i> ), Birds eye (Carrot Weed) ( <i>Cotula australis</i> )	Up to the 4 true leaf stage Greater than the 4 true leaf stage	SA only	700mL/ha 1.4L/ha	DO NOT use where undersown medics occur.
Fireweed ( <i>Senecio</i> spp.)	Seedlings	Qld, NSW, ACT only	1.4L/ha 2.8L/ha	Apply during the Autumn-Winter period when weeds are young and actively growing. Not effective on mature plants.	
				Early flowering stage	

SITUATION AND CROP	WEEDS CONTROLLED	WEED STAGE	STATE	RATE	CRITICAL COMMENTS
Wheat, Barley, Cereal rye, Oats and Triticale: including those undersown with Clover, Lucerne or Medics; Clover Seed Crops	Amsinckia ( <i>Amsinckia</i> spp.), Capeweed ( <i>Arctotheca calendula</i> ), Chamomile ( <i>Matricaria matricarioides</i> ), Charlock ( <i>Sinapisarvenses</i> ), Corn gromwell (Ironweed, Sheepweed) ( <i>Buglossoides arvensis</i> ), Fat hen ( <i>Chenopodium album</i> ), Field madder ( <i>Sherardia arvensis</i> ), Hexham scent (King Island Melilot) ( <i>Melilotus indicus</i> ), Horned poppy ( <i>Glaucium flavum</i> ), Indian hedge mustard ( <i>Sisymbrium orientale</i> ), Lesser swinecress ( <i>Coronopus didymus</i> ), Mexican poppy ( <i>Argemone ochroleuca</i> ), Mintweed ( <i>Salvia reflexa</i> ), Purple calandrinia ( <i>Calandrinia menziesii</i> ), Paterson's curse (Salvation Jane) ( <i>Echium plantagineum</i> ), Rough poppy ( <i>Papaver hybridum</i> ), Saffron thistle ( <i>Carthamus lanatus</i> ), Shepherd's purse ( <i>Capsella bursa-pastoris</i> ), Slender thistle ( <i>Carduus tenuiflorus</i> ), Turnip weed (Giant mustard) ( <i>Rapistrum rugosum</i> ), Wild radish ( <i>Raphanus raphanistrum</i> )	Up to the 5 leaf stage or when plants are no more Than 50mm in diameter	WA only	750mL/ha PLUS 440mL Ken-Amine 625/ha OR PLUS 700mL of MCPA 500/ha	<b>CROP STAGE:</b> <b>Cereals:</b> 3 leaf to fully tillered. (Z13-30). <b>Clover:</b> Apply after clover seedlings have at least three trifoliolate leaves <b>APPLICATION:</b> Refer to General Instructions. Avoid application when maximum daily temperatures above 20°C occur, or are likely to occur for a few days after application. Apply only in the Autumn and Winter (mid-April to end of August). Apply only when weeds are actively growing and before the weeds are shielded by the crop. <b>CAUTION:</b> Some crop damage and/or yield loss may occur in Olympic and Shortim wheat cultivars. DO NOT apply to Persian clover or Berseem clover. A slight burning of clover may occur. DO NOT use 2,4-D or MCPA tank mix if wheat, oats, barley or triticale are undersown with lucerne, clover or medics unless some damage is acceptable. This product may be tank mixed with Ken-Grass 375 for control of Annual rye grass and wild oats in wheat and barley only. DO NOT use 2,4-D or MCPA tank mix if using a tank mix with Ken-Grass 375.
			Vic, SA only	1.4/ha PLUS 440mL Ken-Amine 625/ha OR PLUS 700mL of MCPA 500/ha	
			Vic only		
	Common peppercress ( <i>Lepidium africanum</i> ), Tree hogweed ( <i>Polygonum patulum</i> )		Vic only		
	Black bindweed (Climbing Buckwheat) ( <i>Fallopia convolvulus</i> )	Up to the 8 leaf stage	Vic, SA only		
Wheat, Oats, Barley, Triticale alone or with undersown lucerne and clover	Common sowthistle ( <i>Sonchus oleraceus</i> )	Up to 4 leaves	Qld, Northern NSW	1.4L/ha	<b>APPLICATION:</b> Refer to General Instructions. Avoid application when maximum daily temperatures above 20°C occur, or are likely to occur for a few days after application. Good coverage of the spray on the weed is essential. Apply only in the Autumn and Winter (mid-April to end of August). For optimum control, apply only when the weeds are actively growing and before the weeds are shielded by the crop. <b>CAUTION:</b> <b>Lucerne and Clover:</b> This product will cause slight leaf burning of undersown lucerne and clover. DO NOT apply to Persian clover or Berseem clover. Strictly for use on small weeds early in crop before shading out. <b>CROP STAGE:</b> <b>Wheat, Barley, Oats and Triticale:</b> 3 leaf to mid tillering (Z13-23). Use a minimum spray volume of 70L/ha. If Group B resistance is suspected, or weeds have grown through a previous application of chlorsulfuron apply 2.1L/ha.

SITUATION AND CROP	WEEDS CONTROLLED	WEED STAGE	STATE	RATE	CRITICAL COMMENTS
Wheat, Oats, Barley, Triticale alone or with undersown lucerne and clover	Common sowthistle ( <i>Sonchus oleraceus</i> )	4-8 leaves	Qld, Northern NSW	2.1L/ ha	<b>CROP STAGE:</b> <b>Wheat, Barley, Oats and Triticale:</b> 3 leaf to full tillering (Z13-30). This product will cause slight leaf burning of undersown lucerne and clover.
Wheat, Barley, Oats and Triticale	Three cornered Jack (Doublegee, Spiny emex) ( <i>Emex australis</i> ), Wireweed ( <i>polygonum aviculare</i> ), Varigated thistle ( <i>Silybum marianum</i> ) Fumitories ( <i>Fumaria</i> spp.)	Up to the 4 leaf stage or when plants are no more than 50mm in diameter	Qld, NSW, Tas, SA, ACT only	1.4/ha PLUS 440mL Ken-Amine 625/ha OR PLUS 700mL of MCPA 500/ha	<b>CROP STAGE:</b> 5 leaf to fully tillered (Z15-30): Apply under good growing conditions. <b>BOOM SPRAYING:</b> Use a minimum of 220L spray per hectare on Black bindweed when crop is greater than 300mm high. <b>CAUTION:</b> Some crop damage and/or yield loss may occur in Olympic and Shortim wheat cultivars. DO NOT mix with 2,4-D or MCPA tank mix if wheat, oats, barley or triticale are undersown with lucerne, clover or medics unless some damage is acceptable. DO NOT apply to medics (WA only). This product may be tank mixed with Ken-Grass 375 for control of Annual Rye Grass and Wild Oats in Wheat and Barley only. DO NOT use 2,4-D or MCPA tank mix if using a tank mix with Ken-Grass 375. <b>APPLICATION:</b> Refer to General Instructions. Avoid application when maximum daily temperatures above 20°C occur, or are likely to occur for a few days after application.
	Wild mustard ( <i>Sisymbrium</i> spp.), Wild turnip ( <i>Brassica tournefortii</i> ) Cleavers		Qld, NSW, ACT, Vic only Qld, Vic only		
	Three cornered Jack (Doublegee, Spiny emex) ( <i>Emex australis</i> ), Wireweed ( <i>Polygonum aviculare</i> ) Fumitories ( <i>Fumaria</i> spp.)	4-5 leaf stage	WA only Qld, Vic, NSW, ACT, SA, Tas only Qld, NSW, Vic, SA, ACT only		
	Black bindweed (Climbing buckwheat) ( <i>Fallopia convolvulus</i> ), Chamomile ( <i>Matricaria matricarioides</i> ), Fat hen ( <i>Chenopodium album</i> ), Field madder ( <i>Sherardia arvensis</i> ), Lesser swinecress ( <i>Coronopus didymus</i> ), Mountain sorrel ( <i>Oxalis acetosella</i> ), Purple calandrinia ( <i>Calandrinia menziesii</i> ), Paterson's curse (Salvation Jane) ( <i>Echium plantagineum</i> ), Shepherd's purse ( <i>Capsella bursa-pastoris</i> ), Three-horned bedstraw ( <i>Gallium tricometum</i> ), Variegated thistle ( <i>Silybum marianum</i> )	5-8 leaf stage	All States		
	Fallow & Non Crop	Cowvine or Peachvine ( <i>Ipomoea lonchophylla</i> ) Climbing buckwheat ( <i>Fallopia convolvulus</i> ) Volunteer conventional cotton ( <i>Gossypium hirsutum</i> ) Volunteer Roundup Ready® cotton ( <i>Gossypium hirsutum</i> )	Up to 6 leaves Up to 8 leaves Cotyledon to mature, vining plants Cotyledon to 6 leaves Cotyledon to 6 leaves		
Pastures	Adonis (Pheasant's Eye), Bird's Eye (Carrot Weed)	Up to 4 leaves More than 4 leaves	SA only	700mL/ ha 1.4L/ ha	Apply in May-June. Medics may be severely scorched at the low rate and killed at the high rate. DO NOT apply to polymorpha medics. Avoid application to clover-based pastures in temperatures over 20°C or when temperatures above 20°C may follow for some days after application; as seedling mortality may occur and established plants may be damaged. Apply with low volume boom spray during Autumn/Winter, when weeds are young and actively growing. Not effective on mature plants.
	Doublegee, Wild radish Matricaria	Up to 4 leaves Up to 3 leaves	WA only	1.5L/ ha 750mL/ ha	
	Fireweed	Seedlings Early flowering	Qld, NSW, ACT only	1.4L/ha 2.8L/ha	

SITUATION AND CROP	WEEDS CONTROLLED	WEED STAGE	STATE	RATE	CRITICAL COMMENTS
Pastures, Roadsides and Rights of Way	African daisy ( <i>Senecio pterophorus</i> )	Up to and including the flowering stage	SA only	300mL per 100L water	The plants, including the stems, must be thoroughly wet to the point of run-off. Apply when the plants are actively growing. Nufarm Activator® should be added at 125mL per 100L of spray. A follow-up treatment may be necessary for large, well established plants. Ensure an overall spray coverage on weed seedlings.
	Boneseed/Bitou-bush ( <i>Chrysanthemoides monilifera</i> )	Seedlings only	Vic, Tas only	160mL per 100L water	
SITUATION AND CROP	WEEDS CONTROLLED	STATE	RATE		CRITICAL COMMENTS
<b>Turf:</b> Established Couch(including Queensland blue), Buffalo, Bent, Paspalum, Kikuyu, Fescue, and Rye lawns	Bindy-eye ( <i>Calotis hispidula</i> ), Jo-jo (Onehunga) ( <i>Soliva pterosperma</i> ), Cudweed ( <i>Gnaphalium</i> spp.)	All States	6.0L/ 500L water per ha	<b>High Volume:</b> 300mL per 100L	60mL/ 100m <sup>2</sup> per 15L knap-sack

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

#### WITHHOLDING PERIOD

**CROP HARVEST: NOT REQUIRED WHEN USED AS DIRECTED.**

**GRAZING/STOCKFOOD: DO NOT GRAZE TREATED CROPS OR CUT FOR STOCKFOOD FOR 8 WEEKS AFTER APPLICATION.**

#### GENERAL INSTRUCTIONS

Before opening, carefully read Directions for Use, Precautionary and Protection Statements, Storage and Disposal, Safety Directions and First Aid Instructions. Use only at recommended growth stages of crops and weeds. DO NOT use on crops other than recommended.

#### MIXING INFORMATION

To ensure even mixing, half fill the spray tank with clean water and add the required amount of this product. Agitate thoroughly, then add the remainder of water. Agitate again before spraying commences.

#### BOOM SPRAYER

USE ONLY low boom equipment set up to provide good coverage of weeds within the crop canopy. USE ONLY coarse spray quality according to ASAE S572 definition for standard nozzles. Apply 50-200L (WA: 50-100L) of water per hectare. Calibrate the machine before use to ensure that the correct application rate is obtained. Check height of boom above the crop to ensure that the spray is evenly distributed. A minimum of 160L and preferably 200L of water per hectare is recommended where the weed infestation is heavy or the crop cover is dense.

#### AERIAL APPLICATION

Apply a minimum of 22L of water per hectare. Markers should always move upwind before aircraft passes overhead. Application by microneaire equipment may give variable results.

#### EQUIPMENT MAINTENANCE

After use, wash equipment thoroughly and rinse several times prior to re-use. Wash sprayer thoroughly with clean water after use. If using tank mixes with Ken-Amine 625 or MCPA 500 apply with equipment that will not be used in 2,4-D and MCPA susceptible crops, particularly cotton and vines.

#### COMPATIBILITY

This product may be mixed in the spray tank with one of the following herbicides or insecticides in wheat and barley only. Ken-Grass 375, Omethoate, Kensban 500, dimethoate, endosulfan, Ken-Amine 625 Herbicide (2,4-D Amine), MCPA 500 Herbicide, MCPA 750, Ken-Chlor 750 WDG, Ken-Met 600 WG, Brodal\*, Glyphosate products (Ken-Up 450 CT, Credit®/Bonus®, Weedmaster® Duo and Weedmaster® 490 CT Plus, Ken-Up Aquatic 360, Roundup PowerMAX, Dicamba 500, LV MCPA 500, Bromoxynil 200 is compatible with Fluroken 200 when needed for control of mixed weed populations in grain sorghum. As formulations of other manufacturers products are beyond the control of Kenso Agcare, all mixtures should be tested prior to mixing commercial quantities.

#### GROUP C HERBICIDE

#### RESISTANT WEEDS WARNING

Bromoxynil 200 Selective Herbicide (Bromoxynil 200) is a member of the nitriles group of herbicides. Bromoxynil 200 has the inhibitors of photosynthesis at photosystem II mode of action. For weed resistance management Bromoxynil 200 is a Group C herbicide. Some naturally occurring weed biotypes resistant to Bromoxynil 200 and other Group C herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by this product or other Group C herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Kenso Agcare Australia Limited accepts no liability for any losses that may result from the failure of Bromoxynil 200 to control resistant weeds.

#### PRECAUTION

**RE-ENTRY PERIOD:** DO NOT allow entry into treated areas until the spray has dried, unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

#### PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

**Drift Warning:** DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

**Note:** Maximum daily temperatures at or above 20°C at application, or on days following treatment may result in seedling mortality and established plants may be damaged.

#### PROTECTION OF LIVESTOCK

Low hazard to bees. May be applied at any time as recommended in the Directions for Use.

#### PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Dangerous to fish. DO NOT contaminate streams, rivers or waterways with the chemical or used containers.