

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

KENSO AGCARE

CAPTAN 800 WG FUNGICIDE

ACTIVE CONSTITUENT: 800 g/kg CAPTAN

GROUP M4 FUNGICIDE

For the control of certain diseases in a range of fruit crops,
turf & ornamentals as per directions for use.



Kenso Corporation (M) Sdn Bhd
Level 1, 98 Commercial Road,
Teneriffe QLD 4005
Phone (07) 3216 1188
www.kenso.com.au

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USE

CONTENTS: 5 Kilograms APVMA Approval No.: 66892/ 54638

KENSO AGCARE CAPTAN 800 WG FUNGICIDE

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots in compliance with relevant Local, State or Territory government regulations. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Harmful if inhaled or swallowed. Will damage eyes. May irritate the skin, nose and throat. Avoid contact with eyes and skin. If product or spray in eyes, wash it out immediately with water. DO NOT inhale dust or spray mist. When opening the container, mixing and loading and preparing spray wear cotton overalls buttoned to the neck and wrist [or equivalent clothing], elbow-length PVC/nitrile gloves and face shield or goggles. When using the prepared spray wear cotton overalls buttoned to the neck and wrist [or equivalent clothing] and elbow-length rubber gloves. Repeated exposure may cause allergic disorders. Sensitive workers should use protective clothing. Wash hands after use after each day's use, wash gloves, goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a Doctor or Poisons Information Centre (Phone Australia 13 11 26).

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet which can be obtained from www.kenso.com.au.

CONDITIONS OF SALE

Kenso Corporation (M) Sdn. Bhd. will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply of these goods other than

responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Kenso Corporation (M) Sdn. Bhd. is limited to the replacement of the goods or (at the option of Kenso Corporation (M) Sdn. Bhd.) the refund of the price paid and where possible sufficient part of the goods to enable proper examination being returned to Kenso Corporation (M) Sdn. Bhd. within thirty days of delivery.

In a Transport Emergency
Dial **000** Police or Fire Brigade



Additional statements required by Globally Harmonised System of Classification and Labelling of Chemicals (GHS) and Safe Work Australia: **MAY CAUSE AN ALLERGIC SKIN REACTION. CAUSES SERIOUS EYE DAMAGE. TOXIC IF INHALED. SUSPECTED OF CAUSING CANCER.**
Precautionary: Do not handle until all safety precautions have been read and understood. Avoid breathing dust/fume/ gas/mist/vapours/spray. Use only outdoors or in well-ventilated area. Contaminated work clothing should not be allowed out of workplace. Wear protective gloves/protective clothing/eye or face protection. IF ON SKIN: Wash with plenty of soap and water. IF INHALED: Remove person to fresh air and keep comfortable for breathing. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. IF exposed or concerned: Get medical advice/attention. Immediately call a POISON CENTER/doctor/physician. Specific treatment (see FIRST AID on this label). If skin irritation or rash occurs: Get medical advice/attention. Take off contaminated clothing and wash it before reuse. Store in a well-ventilated place. Keep container tightly closed. Store locked up.

Batch No.:

Date of Manufacture:

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

KENSO
enggard
CAPTAN
800 WG

Fungicide

ACTIVE CONSTITUENT: 800 g/kg CAPTAN

GROUP **M4** FUNGICIDE

For the control of certain diseases in a range of fruit crops, turf & ornamentals as per directions for use.

IMPORTANT: READ THE LEAFLET BEFORE USE

APVMA Approval No.: 66892/54638



Kenso Corporation (M) Sdn. Bhd.
Level 1, 98 Commercial Road,
Teneriffe QLD 4005
Phone (07) 3216 1188

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight.

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots in compliance with relevant Local, State or Territory government regulations. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Harmful if inhaled or swallowed. Will damage eyes. May irritate the skin, nose and throat. Avoid contact with eyes and skin. If product or spray in eyes, wash it out immediately with water. DO NOT inhale dust or spray mist. When opening the container, mixing and loading and preparing spray wear cotton overalls buttoned to the neck and wrist [or equivalent clothing], elbow-length PVC/nitrile gloves and face shield or goggles. When using the prepared spray wear cotton overalls buttoned to the neck and wrist [or equivalent clothing] and elbow-length rubber gloves. Repeated exposure may cause allergic disorders. Sensitive workers should use protective clothing. Wash hands after use after each day's use, wash gloves, goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a Doctor or Poisons Information Centre (Phone Australia 13 11 26).

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet which can be obtained from www.kenso.com.au

CONDITIONS OF SALE

Kenso Corporation (M) Sdn. Bhd. will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Kenso Corporation (M) Sdn. Bhd. is limited to the replacement of the goods or (at the option of Kenso Corporation (M) Sdn. Bhd.) the refund of the price paid and where possible sufficient part of the goods to enable proper examination being returned to Kenso Corporation (M) Sdn. Bhd. within thirty days of delivery.

® is not a registered trademark of Kenso

In a Transport Emergency
Dial **000**
Police of Fire Brigade



Kenso Corporation (M) Sdn. Bhd.
Level 1, 98 Commercial Road,
Teneriffe QLD 4005
Phone (07) 3216 1188

APVMA Approval No.: 66892/54638

DIRECTIONS FOR USE:

Restraints: Do not apply by aircraft
Tree and Vine crops:

Rate					Critical Comments	
In the following table, all rates are given for dilute spraying. For concentrate spraying, refer to the Mixing/Application section.						
Crop	Disease	States	Rate/100L	WHP	For all uses in this table apply by dilute or concentrate spraying equipment. Apply the same total amount of product to the target crop whether applying this product by dilute or concentrate spraying methods.	
Apples	Black spot (<i>Venturia inaequalis</i>)	All States	Dilute spraying 125 g/100 L Concentrate spraying Refer to the Mixing/Application section	7 days		Apply from pink stage onwards following a copper spray at green tip, then at 7 day intervals until petal fall. After petal fall apply as a cover spray at 10-14 day intervals. The risk of russet may be increased in some varieties. Where this is of concern use alternative materials between full bloom and second cover. Apply to point of run-off. Apply no more than five applications per season. Do not use at concentrate spray rates greater than 5X the dilute application rate.
Pears	Black spot (<i>Venturia pirina</i>)	All States		7 days		Apply from pink stage onwards following a copper spray at green tip, then at 7 day intervals until petal fall. After petal fall apply as a cover spray at 10-14 day intervals. The risk of russet may be increased in some varieties. Where this is of concern use alternative materials between full bloom and second cover. Apply to point of run-off. Apply no more than five applications per season. Do not use at concentrate spray rates greater than 5X the dilute application rate.
Grapes	Black spot (<i>Elsinoe ampelina</i>)	All States		7 days		Apply at 50% bud-burst and repeat application 10-14 days later. Apply to point of run-off. Apply no more than five applications per season. Do not use at concentrate spray rates greater than 5X the dilute application rate.

Rate					Critical Comments	
In the following table, all rates are given for dilute spraying. For concentrate spraying, refer to the Mixing/Application section						
Crop	Disease	States	Rate/100L	WHP	For all uses in this table apply by dilute or concentrate spraying equipment. Apply the same total amount of product to the target crop whether applying this product by dilute or concentrate spraying methods.	
Grapes	Grey mould (<i>Botrytis cinerea</i>)	All States	Dilute spraying 125 g/100 L Concentrate spraying Refer to the Mixing/Application section	7 days		Apply in a program of sprays applied at early flowering, 80-100% cap fall, pre-bunch closure, veraison and pre-harvest. Apply to point of run-off. Apply no more than five applications per season. Do not use at concentrate spray rates greater than 5X the dilute application rate.
	Downy mildew (<i>Plasmopara viticola</i>)					Apply from flowering onwards, at intervals of 10-14 days. Apply to point of run-off. Apply no more than five applications per season. Do not use at concentrate spray rates greater than 5X the dilute application rate.
	Phomopsis Cane and Leaf Blight (<i>Phomopsis viticola</i>)					Commence application at early budburst and repeat at 7 to 14 day intervals until prior to flowering to maintain a cover of new grape foliage. Apply no more than 5 applications per season. Use the shorter application interval where there is rapid shoot growth and conditions are favourable for Phomopsis infection (wet weather). Adjust spray application equipment to obtain a thorough coverage of grape buds and shoots. Do not use at concentrate spray rates greater than 5X the dilute application rate.
Stone Fruit (Except apricots)	Blossom blight and Brown rot (<i>Sclerotinia laxa</i> , <i>S. fructicola</i>)	All States		7 days	Apply at any of the following times as part of a spray program to control blossom blight and brown rot - pink bud, 10% blossom, full bloom, petal fall and shuck fall and pre-harvest sprays at 6, 3 and 1 week prior to harvest. Apply to point of run-off. Apply no more than five applications per season. Do not apply to apricots as damage to fruit and foliage may occur. Do not use at concentrate spray rates greater than 2X the dilute application rate.	

DIRECTIONS FOR USE

Non Tree and Vine Crops:

Crop	Disease	States	Rate/100L	WHP	Critical Comments
Strawberries	Grey mould (<i>Botrytis cinerea</i>) Gloeosporium fruit rot	All States	125g	1 day	Apply every 10 days commencing at blossom stage. Apply to point of run-off. Apply no more than five applications per season.
	Phytophthora fruit rot (<i>Phytophthora nicotianae</i> var. <i>parasitica</i>) Black spot (Anthracnose), (<i>Colletotrichum acutatum</i>) Scorch (<i>Diplocarpon earlianum</i>) Leaf blight (<i>Dendrophoma obscurans</i>)		200g		
Turf	Brown patch (<i>Rhizoctonia solani</i>)	All States	100-150g in 25-50 litres of water for each 100 m ² of turf	-	For preventative treatment from September to April, apply every 5-6 weeks. Apply no more than five applications per season.
Seedlings (Ornamentals & Tobacco)	Damping-off (<i>Pythium</i> spp)	All States	125g	-	Use 3 litres of solution per m ² of seedbed. Apply weekly-fortnightly when weather conditions favouring the disease persists, paying particular attention to bases of the seedlings. When treating tobacco seedbeds wait until seedlings are at least 5cm high.
Roses	Black spot (<i>Diplocarpon rosae</i>)	All States	125g	-	Apply to thoroughly wet both surfaces of foliage. Apply every 7-14 days. Use the shorter interval during showery weather. Apply no more than five applications per season.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

THIS PRODUCT SHOULD NOT BE APPLIED BY AIRCRAFT.

WITHHOLDING PERIODS

STRAWBERRIES: DO NOT HARVEST FOR 1 DAY AFTER APPLICATION.

APPLES, GRAPES, PEARS, STONE FRUIT: DO NOT HARVEST FOR 7 DAYS AFTER APPLICATION.

DO NOT USE TREATED PRODUCE FOR STOCK FOOD WITHIN 7 DAYS OF TREATMENT.

GENERAL INSTRUCTIONS

Fungicide Resistance Warning

GROUP M4 FUNGICIDE

For fungicide resistance management the product is a Group M4 fungicide.

Some naturally occurring individual fungi resistant to Captan 800 WG Fungicide and other Group M4 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by Captan 800 WG Fungicide or other Group M4 fungicides, thus resulting in a reduction in efficacy and possible yield loss.

Since the occurrence of resistant fungi is difficult to detect prior to use, Kenso Agcare Corporation (M) Sdn Bhd accepts no liability for any losses that may result from the failure of Captan 800 WG Fungicide to control resistant fungi.

Mixing

Captan 800 WG mixes readily with water and should not be pre-mixed. Slowly add the required amount of product to the water under agitation in the spray vat. Maintain agitation throughout spraying. After a stoppage, thoroughly re-mix before recommencing spraying.

Compatibility and Crop Safety

Mixtures of Captan 800 WG with commonly used wettable powders and suspension concentrate insecticides and fungicides should not cause any problems. Apart from the products listed below mixtures with emulsifiable concentrates are not recommended.

If tank mixes are to be used observe all directions, precautions and limitations on all products to be used. Caution should also be exercised when spraying any mixtures on varieties of fruit trees or vines where there has been no previous experience of mixtures with Captan 800 WG.

Captan 800 WG. is compatible with any one of the following products: Anvil®, Bayfidan®, Topaz®, Pennncap-M®, Gusathion® 200SC with or without calcium nitrate, Bugmaster® Flowable, Savage 200, Hymal, Insegar®, Kumulus®, Lebaycid®, Kensban 500, Mavrik® Aquaflow, Omite ®300WG (apples only), Pirimor® WG, Pyranica®, Saboteur®, Sanmite®, Systhane®.

A mixture of Captan 800 WG with more than one of these products or with any other product may be ineffective or may cause serious damage. The use of such a mixture is not recommended without prior advice, and would therefore be entirely at the user's risk. If other mixtures or 3 way mixtures of any of these products are being considered consult your Kenso Agcare dealer or representative for advice.

Mixtures of Captan 800 WG with Kensban 500 or Nimrod® may leave an oily residue on the inside of spray vats which may be difficult to remove.

Captan 800 WG should not be combined with Bordeaux mixture, spray lime, lime sulphur or oils. Do not apply Captan 800 WG in combination with or within 14 days before or after winter or summer oil sprays. Combinations of Captan and sulfur should not be used on crops sensitive to sulfur.

Captan 800 WG mixed with Folidol® M500 may be phytotoxic to many varieties of apples and pears. If it is necessary to use both Captan 800 WG and Folidol M500 in a spray program, sprays of these products must be at least 7 days apart.

On pears tank mixtures of Captan 800 WG plus Omite® 300WG and Captan 800 WG and Gusathion ® (without calcium nitrate) plus Omite® 300WG have been found to be phytotoxic. Do not tank mix these products, for use on pears.

Caution should be taken when applying chemical mixtures to all crops. Varieties within crop groups may react differently therefore a small area should be tested prior to application to crop.

Application

For ground application - these rates are for high volume spraying to point of run-off.

DO NOT apply by aircraft.

Dilute Spraying:

Use a sprayer designed to apply high volumes of water up to the point of run-off and matched to the crop being sprayed. Set up and operate the sprayer to achieve even coverage throughout the crop canopy. Apply sufficient water to cover the crop to the point of run-off. Avoid excessive run-off. The required water volume may be determined by applying different test volumes, using different settings on the sprayer, from industry guidelines or expert advice. Add the amount of product specified in the Directions for Use table for each 100 L of water. Spray to the point of run-off. The required dilute spray volume will change and the sprayer set up and operation may also need to be changed, as the crop grows.

Concentrate Spraying:

Use a sprayer designed and set up for concentrate spraying (that is a sprayer which applies water volumes less than those required to reach the point of run-off) and matched to the crop being sprayed. Set up and operate the sprayer to achieve even coverage throughout the crop canopy using your chosen water volume.

Determine an appropriate dilute spray volume (See Dilute Spraying above) for the crop canopy, this is needed to calculate the concentrate mixing rate. The mixing rate for concentrate spraying can then be calculated in the following way:

Example only

1. Dilute spray volume as determined above: For example 1000 L/ha.
2. Your chosen concentrate spray volume: For example 500L/ha.
3. The concentration factor in this example is: 2 x (ie. 1000L / 500L = 2).
4. If the dilute label rate is 125 g/100L, then the concentrate rate becomes 2 x 125, that is 250 g / 100L of concentrate spray.

The chosen spray volume, amount of product per 100L of water, and the sprayer set up and operation may need to be changed as the crop grows. Do not use at a concentration rate higher than that specified in the Critical Comments as crop safety has not been assessed. For further information on concentrate spraying, users are advised to consult relevant industry guidelines, undertake appropriate competency training and follow industry Best Practices.

PRECAUTION

Re-entry Periods

Do not allow entry into treated areas until the spray has dried. When prior entry is necessary wear cotton overalls buttoned to the neck and wrists (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Dangerous to fish. DO NOT apply directly to water. DO NOT apply under meteorological conditions or from spraying equipment which could be expected to cause spray to drift onto water, or to areas where surface water is present. DO NOT contaminate surface water with the concentrated product, spray mix, rinsate, drainage water from glasshouses or the used container.