

# POISON

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

*KENSO AGCARE*

# MCPA 750

## SELECTIVE HERBICIDE

ACTIVE CONSTITUENT: 750 g/L MCPA present as the DIMETHYLAMINE SALT

**GROUP 4 HERBICIDE**

For the selective control of broadleaf weeds in cereals, linseed, pastures, sugar cane, rice, poppies and turf as per Directions for Use table



Kenso Corporation (M) Sdn Bhd  
Level 1, 98 Commercial Road,  
Teneriffe QLD 4005  
Phone (07) 3216 1188  
[www.kenso.com.au](http://www.kenso.com.au)

**IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USE**

**CONTENTS: 20 Litres APVMA Approval No.: 62385/ 123094**

# KENSO AGCARE MCPA 750 SELECTIVE HERBICIDE

## STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. DO NOT store in direct sunlight. Triple rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush, or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

**Refillable containers (110L):** Store the original sealed drum in a cool, well ventilated area. Do not store for prolonged periods in direct sunlight. Do not tamper with the non-return valve or the security seal. Do not contaminate the drum with water or any other foreign matter.

After each use of the product, please ensure the non-return valve, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the drum have been used, please return the empty drum to the point of purchase.

**Refillable containers (1000L):** Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

## SAFETY DIRECTIONS

Harmful if swallowed. Attacks eyes. Will irritate the skin. When opening the container, preparing spray and using the prepared spray wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and goggles. If product on skin, immediately wash area with soap and water. If product or spray in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms, and face thoroughly with soap and water. After each day's use wash gloves, goggles and contaminated clothing.

## FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766.

## SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet, which can be obtained from [www.kenso.com.au](http://www.kenso.com.au).

Batch No.:

Date of Manufacture:

## CONDITIONS OF SALE

Kenso Corporation (M) Sdn. Bhd. will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Kenso Corporation (M) Sdn. Bhd. is limited to the replacement of the goods or (at the option of Kenso Corporation (M) Sdn. Bhd.) the refund of the price paid and where possible sufficient part of the goods to enable proper examination being returned to Kenso Corporation (M) Sdn. Bhd. within thirty days of delivery.

In a Transport Emergency  
Dial **000** Police or Fire Brigade



Additional statements required by Globally Harmonised System of Classification and Labelling of Chemicals (GHS) and Safe Work Australia: **HARMFUL IF SWALLOWED. HARMFUL IN CONTACT WITH SKIN. CAUSES SKIN IRRITATION. CAUSES SERIOUS EYE DAMAGE. HARMFUL IF INHALED.**  
**Precautionary:** Avoid breathing dust/fume/gas/mist/vapours/spray. Wash contacted areas thoroughly after handling. Do not eat, drink or smoke when using this product. Use only outdoors or in a well-ventilated area. Wear protective gloves, protective clothing and eye or face protection. IF SWALLOWED: Call a POISON CENTER or doctor/physician if you feel unwell. IF ON SKIN: Wash with plenty of soap and water. IF INHALED: Remove person to fresh air and keep comfortable for breathing. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. Immediately call a POISON CENTER or doctor/physician. Specific treatment (see FIRST AID on this label). Rinse mouth. If skin irritation occurs: Get medical advice/attention. Take off contaminated clothing and wash it before reuse.

**POISON**  
KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

**KENSO**  
*agcare*

**MCPA**  
**750**

**Selective Herbicide**

**ACTIVE CONSTITUENT: 750 g/L MCPA present as the DIMETHYLAMINE SALT**

**GROUP 4 HERBICIDE**

For the selective control of broadleaf weeds in cereals, linseed, pastures, sugar cane, rice, poppies and turf as per Directions for Use table.

**DIRECTIONS FOR USE LEAFLET**

**APVMA Approval No.: 62385/123094**



**Kenso Corporation (M) Sdn. Bhd.**  
Level 1, 98 Commercial Road,  
Teneriffe QLD 4005  
Phone (07) 3216 1188

**STORAGE AND DISPOSAL**

Store in the closed, original container in a cool, well-ventilated area. DO NOT store in direct sunlight. Triple rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

**Refillable containers (110L) :**

Store the original sealed drum in a cool, well ventilated area. Do not store for prolonged periods in direct sunlight. Do not tamper with the non-return valve or the security seal. Do not contaminate the drum with water or any other foreign matter. After each use of the product, please ensure the non-return valve, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the drum have been used, please return the empty drum to the point of purchase.

**Refillable containers (1000L) :**

Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

**SAFETY DIRECTIONS**

Harmful if swallowed. Attacks eyes. Will irritate the skin. When opening the container, preparing spray and using the prepared spray wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and goggles. If product on skin, immediately wash area with soap and water. If product or spray in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms, and face thoroughly with soap and water. After each day's use wash gloves, goggles and contaminated clothing.

**FIRST AID**

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766.

**SAFETY DATA SHEET**

Additional information is listed in the Safety Data Sheet, which can be obtained from www.kenso.com.au.

**CONDITIONS OF SALE**

Kenso Corporation (M) Sdn. Bhd. will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Kenso Corporation (M) Sdn. Bhd. is limited to the replacement of the goods or (at the option of Kenso Corporation (M) Sdn. Bhd.) the refund of the price paid and where possible sufficient part of the goods to enable proper examination being returned to Kenso Corporation (M) Sdn. Bhd. within thirty days of delivery.

\* are not registered trademarks of Kenso Corporation (M) Sdn Bhd

In a Transport Emergency  
Dial **000**  
Police of Fire Brigade



**Kenso Corporation (M) Sdn. Bhd.**  
Level 1, 98 Commercial Road,  
Teneriffe QLD 4005  
Phone (07) 3216 1188

**APVMA Approval No.: 62385/123094**

**DIRECTIONS FOR USE**

**RESTRAINTS:** DO NOT apply if crop or weeds are stressed due to dry or excessively moist conditions.  
DO NOT apply if rain is likely within 6 hours.

**1. FIELD CROPS**

Situation & Crop	Weeds Controlled	State	Rate	Critical Comments	
<b>Wheat, Barley, Oats, Triticale</b>	Refer to Weed Table	SA, Tas only	460mL-1.45L/ha	Apply through the tillering stage after the crop plants have 5 leaves and before they commence jointing.	
		Qld only		Apply from early tillering (4-5 fully emerged main stem leaves plus one or more tillers) until before the boot stage (visible swelling of the head at the top of the main stem).	
		NSW, ACT only		Apply 460mL rate after crop has reached 5 leaf stage and weeds marked # in weed table have 3-6 leaves. Higher rates after the first node can be felt at the base of a tiller and before swelling of the head can be felt in a tiller.	
		Vic only		Apply 460mL rate after crop has reached 2-3 leaf stage and weeds marked # in weed table have 2-4 leaves. Higher rate after the crop plants have 5 leaves to fully tillered.	
<b>Wheat, Barley, Triticale, Cereal Rye</b>	Capeweed, Doublegee, Erodium, Mustard, Rapistrum, Wild Turnip, Wild Radish	WA only	270mL/ha plus 350mL/ha Kenso Agcare Diuron 500	CROP: 3-4 leaf stage. (Zadoks 13-14)	
			330mL/ha plus 400mL/ha Kenso Agcare Diuron 500	CROP: 4-5 leaf stage. (Zadoks 14-15)	
				Apply by aircraft or boom with 20-100L/ha water, when most weeds have germinated and are in 2-5 leaf stage. Crop and weeds should be dry at time of application. Some temporary yellowing of crop may occur after application. Undersown sub-clovers may be slightly retarded. DO NOT apply to undersown medics.	
		NSW, ACT, Vic only	330mL plus 500mL/ha Kenso Agcare Diuron 500	Apply by boom spray or aircraft when the crop is at the 3-5 leaf stage (NSW and SA only) and 5 leaf stage (Vic only). Weeds must be in the 2-5 leaf stage. Do not apply to undersown medic or lucerne. Do not use on Kamilaroi, Kite, Olympic or Shortim Wheat.	
<b>Wheat, Barley, Oats, Triticale</b>	Soldier Thistle	SA only	230mL plus 500mL/ha Kenso Agcare Diuron 500	Do not use on very light sandy soils.	
			460mL plus 200mL/ha Ken-Trel 300	Apply to crop at tillering stage after the crop has 5 leaves and before the crop commences jointing.	
<b>Wheat, Barley, Oats, Triticale</b>	Skeleton Weed		660mL plus 500mL/ha Ken-Trel 300	Apply to rosettes and from 5-15cm diameter when the crop is between the 5 leaf and late tillering growth stages. It is essential that all the Skeleton weed is emerged and minimum of 5cm diameter which is usually not before August. This rate will provide control of Skeleton Weed until harvest and will substantially reduce seed numbers the following season.	
				This rate will provide control of Skeleton Weed until harvest and will substantially reduce seed numbers the following season.	
<b>Linseed</b>	Refer to Weed Table	Qld only	460mL-1.45L/ha	Apply when crop is 10-15cm tall with at least 170L/ha of water. Do not spray after buds appear. Some wilting effect may be seen following application but crop usually recovers quickly. Do not apply by aircraft.	
		NSW, ACT, Vic, SA, Tas only	460mL-730mL/ha		
<b>Sugar Cane</b>	Blue Top, Chinese Burr, Flannel Weed, Gambia Pea, Bell Vine, Streaked Rattle Pod, Bindweed, Pink Convolvulus, Cupids Flower, Merremia Vine, Morning Glory	Qld only	930mL/ha	POST-EMERGENT: Apply as directed inter-row spray. Do not exceed rates with phenoxy sensitive varieties.	
			As above plus Fat Hen, Noogoora Burr	1.4L/ha	
			All of the above	6.7L/ha	PRE-EMERGENT: Apply over the row in a 45cm band within 24 hours of planting.
<b>Flax</b>	Refer to Weed Table	Vic, Tas only	325mL/ha	Apply to crop when 8-20cm high.	

Situation & Crop	Weeds Controlled	State	Rate	Critical Comments
<b>Rice</b>	Dirty Dora, Docks, Sedges, Starfruit, Buddah (Butter pea)	All States	485mL – 970mL/ha	Apply by aircraft at early/mid tillering stage of rice but not later than panicle initiation. Use 485 mL/ha at 35 days after aerial sowing (approx. 2L per rice plant). Use 970mL/ha at 45 days or more after cereal sowing (3 tillers or more per rice plant).  The crop should be sprayed before flowering of the weeds.  WATER MANAGEMENT: Before herbicide application. Lower water level in the bays to at least two thirds of the weed growth for direct contact with Kenso Agcare MCPA 750 Selective Herbicide.  Weeds covered by water will not be controlled. Raise water depth to normal levels 24 hours following spraying.
<b>Oilseed poppies (Papaver somniferous)</b>	Fat hen (Chenopodium album) Spear thistle (Cirsium vulgare)	Tas and NSW	1L/ha	Use only in consultation with the poppy contracting company.  Apply at least one week after petal fall after poppy seed has been set.  Apply by aircraft in 30-40L of water/ha. This treatment will control late germinating weeds of they are green and at the bud or early flowering stage. Spraying at this time will twist weeds below the crop canopy.  DO NOT apply in horticultural areas where there is a risk of spray or vapour drift over susceptible crops.

**2. PASTURES, SEED CROPS & TURF**

Situation & Crop	Weeds Controlled	State	Rate	Critical Comments
<b>Grass Pastures</b>	Refer to Weed Table	All States	460mL-2.7L/ha	Apply to established grass based pastures. Damage may occur to Legumes if present. See "Legume Tolerance" below. NOTE: Capeweed, paterson's curse and variegated thistle may be poisonous to stock after spraying.
		SA only	660mL plus 400mL/ha Kenso Agcare Diuron 500	Use this mixture only if spraying is delayed until early winter. Rosettes should not exceed 20cm diameter.
		SA only	460mL-1.45L/ha	Use low rate on seedlings with no more than 2 true leaves. Use 100 litres of water for best results. If clover present do not exceed 575mL.
		NSW, ACT only	2L/ha	Apply in Autumn when horehound plants are at the seedling stage. If sub clover or medic plants are present they will be damaged by this application.
<b>Grass Seed Crops</b>	Refer to Weed Table	Qld, SA, Tas, Vic, WA only	460mL-2.7L/ha	Apply to established grass seed crops.
<b>Sub-clover seed crops</b>	Refer to Weed Table	Vic, SA only	460mL-960mL/ha	Do not exceed rate specified. Spray in Autumn and early Winter.
<b>Pastures - Spray/Graze Technique</b>	Caltrop, Capeweed, Charlock, Mustards, Paterson's Curse, Shepherds Purse, Saffron, Slender, Spear or Variegated Thistle, Turnip Weed, Wild Radish, Wild Turnip	NSW, ACT, Vic only	230-930mL/ha	Spray actively growing 6-8 weeks old weeds. Introduce stock 7-10 days after spraying, preferably sheep as cattle are less effective. Use stocking rate at least 5 times heavier than normal until weeds have been reduced but before survival of desirable pasture species is threatened. Lucerne and medics may be damaged and should be grazed short before spraying. Refer "Legume Tolerance" under General Instructions. PRECAUTION: An increased quantity of poisonous plants may be eaten by stock using Spray/Graze, eg. caltrop, capeweed, paterson's curse and variegated thistle and deaths could result from causes such as nitrate poisoning. With paterson's curse preferably graze stock soon destined for slaughter and avoid extended periods of grazing. Avoid grazing with young or breeding stock. Do not graze horses or pigs on paterson's curse. Observe Withholding Period of at least 7 days after spraying.
		Tas only	460mL/ha	
		WA only	610mL-1.25L/ha	
<b>Turf</b>	Refer to Weed Table	All States	330mL/ha or 2.8mL per 100 sq. metres	Newly laid - Delay until well established or at least two months old. Apply in high volume to actively growing weeds. See "Legume Tolerance". Buffalo grass is more easily damaged than other grasses.
			930mL-1.8L/ha or 10-19mL per 100 sq. metres	Established - Apply in high volume to actively growing weeds. Do not mow for 2 days before application. Some transitory damage may occur to fine turf grasses and strawberry clover. See "Legume Tolerance". Buffalo grass is more easily damaged than other grasses.



### 3. SPOT SPRAYING

Situation & Crop	Weeds	State	Mixing Rates/Critical Comments
<b>High Volume Spraying</b>	Refer to Weed Table	All States	Add 1/10th of rate on Weed Table to 150 litres of water. Each 150 litres of mix will cover 1000m <sup>2</sup> (1/10th Ha.) eg. If rate in Weed Table is 1.4L use 140mL/150L water.
<b>Knapsack Application</b>			Add 1/100th of rate on Weed Table to 10 litres of water. Each 10 litres of mix will cover 100m <sup>2</sup> (1/100th Ha.) eg. If rate in Weed Table is 1.4L use 14mL/10L water.

### 4. NON CROP AREAS, FALLOW LAND, INDUSTRIAL & COMMERCIAL AREAS, RIGHTS-OF-WAY

Situation & Crop	Weeds Controlled	State	Rate	Critical Comments
<b>Non Crop Areas, Fallow Land, Industrial &amp; Commercial Areas, Rights of Way</b>	Refer to Weed Table	All States	Refer to Weed Table	Apply at rates shown in Weed Table when weeds only are present and no crop or pasture is involved.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

#### WITHHOLDING PERIOD:

**HARVEST: OILSEED POPPIES: DO NOT HARVEST FOR 5 WEEKS AFTER APPLICATION.**

**ALL OTHER CROPS: NOT REQUIRED WHEN USED AS DIRECTED.**

**GRAZING: CEREAL PASTURE: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION.**

#### WEED TABLE:

Where weeds are to be sprayed in a crop or pasture use only the rates given for particular crop or situation indicated under the Directions for Use table.

In most cases this will give control, however some hard to kill weeds or those in advanced stages of growth may only be suppressed. The rates listed in the Weed Table below are for use where weeds only are present and no crop or pasture is involved. Use at these rates in a crop or pasture other than for spot spraying may cause damage.

This product should only be used in those states where a rate or range of rates is indicated for the particular weeds listed.

#### APPLICATION RATE PER HECTARE

#: refer to Critical Comments for Wheat, Barley, Oats, Triticale

WEEDS CONTROLLED	Vic	NSW, ACT	SA	Tas	Old	WA	Critical Comments
Ball Mustard	-	-	1.15L	-	-	-	Apply at rosette stage.
Bathurst Burr	1.35L	660mL-1.35L	1.35L	1.35L	1.35L	1.35L	Spray young seedlings only.
Black Bindweed	-	970mL-1.35L	-	-	930mL-1.35L	-	Apply at young stage prior to flowering.
Californian Burr	-	660mL-1.35L	-	1.45L	-	-	Spray young seedlings only.
Cape Tulip	-	-	-	-	-	1.35L	Apply just prior to flowering. Gives mild suppression only.
Capeweed	-	1.45L	1.45L	1.45L	1.45L	-	Apply at seedling stage.
Carrot Weed	1.35L	-	-	-	1.35L	-	Apply at seedling stage.
Charlock	460mL	660mL	460mL	660mL-1.45L	460mL	-	Spray up to rosette stage.
Cutleaf Mignonette	-	-	1.45L	-	-	-	Apply at young rosette stage.
Dandelion	1.35L	1.35L	1.35L	1.35L	1.35L	1.35L	Apply at young rosette stage.
Deadnettle	-	1.45L	-	-	1.45L	-	Moderately susceptible. Apply at seedling stage.
Docks	960mL	-	1.45L	1.45L	-	-	Apply at rosette stage and before flower heads emerge. Only the top growth of old established plants will be affected.
Fat Hen	1.35L	660mL-1.35L	-	1.35L	1.35L	-	Spray up to rosette stage.
Fennel	1.80L	1.80L	1.80L	1.80L	1.80L	-	Apply at young stage.
Field Bindweed	1.35L	-	1.35L	1.35L	1.35L	1.35L	Apply at bud stage, repeating over several seasons.
Field Cress	-	-	-	1.45L	-	-	Apply at young stage.
Flatweed or Catsear	-	-	-	1.45L	-	930mL	Apply when weeds are 10-15cm high.
Fumitory (red)	930mL	930mL	930mL	-	930mL	-	Apply at young stage prior to flowering.
Hoary Cress	3.0L	-	1.45L	-	3.0L	-	Apply before flowering.
Horehound	-	2.0L	1.45L	-	-	-	Apply at rosette stage.
Hedge Mustard	730mL #	-	460mL	460mL	460mL	825mL	Apply at rosette stage.
Hexham Scent or Melilotus	960mL	800mL	930mL	-	930mL	-	Apply at young seedling stage.
Lesser Swinecress	-	-	-	1.45L	-	-	Apply at young stage.
Lincoln Weed	-	-	460mL	-	-	-	Apply at rosette stage.
London Rocket	-	-	-	-	-	825mL	-
Lupins	-	460mL-960mL	-	-	-	825mL	Spray up to 10cm high.
Mintweed	1.35L	1.35L	-	-	1.35L	-	Spray young seedlings only.
Mustards	730mL #	660mL #	730mL	730mL	-	825mL	Spray up to rosette stage.
Noogoora Burr	1.35L	660mL-1.35L	1.35L	-	1.35L	1.35L	Spray young seedlings only.
Opium Poppy	-	-	-	1.45L	-	-	Apply at young stage.
Paterson's Curse	-	660mL-960mL	-	-	2.70L	-	Apply early rosette stage.
Pimpernel	-	-	460mL	-	-	-	Apply at rosette stage.
Plantains	-	1.35L	-	1.35L	1.35L	-	Apply at young stage.
Rapeseed	-	660mL-960mL	-	-	-	-	Spray up to rosette stage.
Rough Poppy	-	460mL-960mL	-	-	-	-	Spray up to rosette stage.
Safflower	-	460mL-660mL	-	-	-	-	Spray up to rosette stage.
Shepherd's Purse	-	-	-	1.45L	-	-	Apply at rosette stage.
Skeleton Weed	-	960mL-1.35L	-	-	-	-	Spray rosettes before aerial growth commences.
Sorrel	-	-	1.45L	-	-	-	Apply to rosette stage.
Stinkwort	1.35L	660mL-960mL	1.35L	-	-	1.35L	Apply when weeds are 10-15cm high.
Sunflower (seedlings)	-	690mL-1L	-	-	-	-	Spray up to rosette stage.

WEEDS CONTROLLED	Vic	NSW, ACT	SA	Tas	Old	WA	Critical Comments
Thistles:							
- Artichoke	1.05L	-	1.45L	-	1.35L	460mL	Apply at rosette stage.
- Nodding	-	1.65L	-	-	-	-	Spray early rosettes. Repeat spray required in following Spring or Autumn.
- Saffron	960mL-1.65L	660mL-1.35L	930mL-1.65L	930mL-1.65L	930mL-1.65L	960mL-1.65L	Spray up to rosette stage. (September WA).
Scotch or Cotton	1.45L	-	460mL-930mL	1.45L	-	-	Apply at seedling stage.
Slender	2.0L	660mL-1.35L	660mL	2.0L	-	2.0L	Spray young seedlings only.
Soldier	730mL	-	730mL	-	-	-	Spray young seedlings. Use higher rate for larger weeds.
Spear	-	960mL-1.35L	-	1.45L	-	-	Apply at rosette stage.
Star	-	-	730mL-930mL	-	-	-	Apply at seedling stage. Use higher rate for larger weeds.
Variegated	960mL	660mL-1.35L	460mL-930mL	960mL	960mL	960mL	Spray young seedlings only. Use 1.35L/ha (SA) for well developed weeds.
Turnip Weed	730mL #	660mL #	660mL	-	-	-	Spray up to rosette stage.
Wild Artichoke	-	-	1.45L	-	-	-	Apply at rosette stage.
Wild Radish	930mL	660mL #	930mL	1.35L	460mL-730mL	800mL	Spray up to rosette stage.
Wild Turnip	730mL #	660mL #	460mL	1.35L	460-730mL	600mL	Spray up to rosette stage.

#### GENERAL INSTRUCTIONS

Weeds should be sprayed while actively growing and at their most susceptible stage. As a general guide, annuals should be sprayed when young and perennials just prior to appearance of bud stage. Extremes of cold or drought are unfavorable. The best conditions are when the soil is moist, weather fine and rain unlikely within 6 hours of application.

#### APPLICATION/MIXING INFORMATION

This product mixes readily with water. Half fill the spray tank with clean water and add the required amount of product. Agitate thoroughly before adding the remaining water.

BOOM SPRAYING - Use 30-120 litres water per hectare.

- Use minimum of 170 litres water per hectare for LINSEED.

AERIAL SPRAYING - Use 10-90 litres water per hectare.

#### EQUIPMENT MAINTENANCE AND USAGE

Equipment that has been used for this chemical should not be used for the application of other materials to sensitive plants unless it has been cleaned with Spraymate Tank & Equipment Cleaner or well washed out with a 2% solution of cloudy ammonia followed by several clear water rinses.

#### COMPATIBILITY/TANK MIXES

This product can be tank mixed with Diflufenican 500, Ken-Met 600/Associate\*, Dicamba 500, Bromicide\*, Kenso Agcare Diuron 500, Ken-Trel 300, Sertin\*, Speedy, Flowable Terbutryn, Tordon\*, and Tribunil\*.

The effectiveness of Imidan\*, Le Mat\*, Metasystox and Dimethoate may be reduced by alkaline hydrolysis if tank mixed with this product. For these situations the addition of Spraymate LI-700\* is recommended. When tank mixing add the required amount of compatible product (and surfactant if recommended) when the spray tank is nearly full and while continuously agitating. If left overnight, agitate thoroughly before spraying.

### GROUP 4 HERBICIDE

#### RESISTANT WEEDS WARNING

Kenso Agcare MCPA 750 Selective Herbicide ('MCPA 750') is a member of the Phenoxy group of herbicides. MCPA 750 has the Disruptors of plant cell growth mode of action. For weed resistance management MCPA 750 is a Group 4 herbicide.

Some naturally-occurring weed biotypes resistant to MCPA 750 and other disruptors of plant cell growth herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by MCPA 750 or other disruptors of plant cell growth herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Kenso Corporation (M) Sdn Bhd accepts no liability for any losses that may result from the failure of MCPA 750 to control resistant weeds.

#### RE-ENTRY PERIOD

Hand weeding should not be performed for 7 days after spray application unless workers wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), chemical resistant gloves and footwear.

#### PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT spray in high winds. DO NOT spray crops or weeds outside the stages indicated in "Critical Comments" as damage, loss of yield or inadequate weed control may result.

**Drift Warning:** DO NOT apply under weather conditions or from spraying equipment that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures. Avoid spray drift onto susceptible crops such as cotton, tobacco, tomatoes, vines, fruit trees, vegetables, legume crops and pastures, oilseed crops and susceptible trees (eg. kurrajongs, belahs, eucalypts).

**Legume Tolerance:** The use of MCPA is preferred to Ken-Amine or Ester 800 in situations where legumes are undersown in the crop.

**NB:** The tolerance of pasture legumes to applied product can vary with species, variety, growing conditions, stage of growth and companion crop cover. DO NOT spray cereal crops undersown with lucerne as damage may result.

#### PROTECTION OF LIVESTOCK

Low hazard to bees. May be applied at any time as recommended in the Directions for Use.

#### PROTECTION OF WILDLIFE, FISH, CRUSTACEAN AND ENVIRONMENT

DO NOT contaminate dams, rivers or streams with the product or used container.