

**CAUTION**  
KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

*KENSO AGCARE*

**METOKEN  
GOLD**  
HERBICIDE

ACTIVE CONSTITUENT: 960 g/L S-METOLACHLOR

**GROUP 15 HERBICIDE**

For the controls of certain annual grasses and broadleaf weeds in certain crops as specified in the Directions for Use table.



Kenso Corporation (M) Sdn Bhd  
Level 1, 98 Commercial Road,  
Teneriffe QLD 4005  
Phone (07) 3216 1188  
[www.kenso.com.au](http://www.kenso.com.au)

**IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USE**

**CONTENTS: 20 Litres APVMA Approval No.: 68625/ 127089**

# KENSO AGCARE METOKEN GOLD HERBICIDE

## STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight.

**Refillable containers:** Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

**Other containers:** Triple-rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

## SAFETY DIRECTIONS

Will damage eyes. Repeated exposure may cause allergic disorders. Sensitive workers should use protective clothing. Avoid contact with eyes. When opening the container and preparing spray, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length pvc gloves, and goggles. When using the prepared spray, wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow-length PVC gloves and face shield or goggles. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves, and face shield or goggles, and contaminated clothing.

## FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766.

## SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet which can be obtained from [www.kenso.com.au](http://www.kenso.com.au).

## CONDITIONS OF SALE

Kenso Corporation (M) Sdn. Bhd. will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Kenso Corporation (M) Sdn. Bhd. is limited to the replacement of the goods or (at the option of Kenso Corporation (M) Sdn. Bhd.) the refund of the price paid and where possible sufficient part of the goods to enable proper examination being returned to Kenso Corporation (M) Sdn. Bhd. within thirty days of delivery.

In a Transport Emergency  
Dial **000** Police or Fire Brigade



Additional statements required by Globally Harmonised System of Classification and Labelling of Chemicals (GHS) and Safe Work Australia: **COMBUSTIBLE LIQUID. MAY CAUSE AN ALLERGIC SKIN REACTION.**

**Precautionary:** Keep away from heat/sparks/open flames/hot surfaces and other ignition sources. No smoking. Avoid breathing dust/fume/gas/mist/vapours/ spray. Contaminated work clothing should not be allowed out of the workplace. Wear protective gloves, protective clothing and eye or face protection. IF ON SKIN: Wash with plenty of soap and water. Specific treatment (see FIRST AID on this label). If skin irritation or rash occurs: Get medical advice/attention. Take off contaminated clothing and wash it before reuse. In case of fire: Use water fog, foam, dry agent (carbon dioxide, dry chemical powder) to extinguish. Store in a well-ventilated place.

Batch No.:

Date of Manufacture:

**CAUTION**  
KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

**KENSO**  
agcare

**METOKEN  
GOLD**

**Herbicide**

**ACTIVE CONSTITUENT: 960 g/L  
S-METOLACHLOR**

**GROUP 15 HERBICIDE**

For the controls of certain annual grasses and broadleaf weeds in certain crops as specified in the Directions for Use table.

**READ THIS LEAFLET BEFORE USE**

**APVMA Approval No.: 68625/127089**



**Kenso Corporation (M) Sdn. Bhd.**  
Level 1, 98 Commercial Road,  
Teneriffe QLD 4005  
Phone (07) 3216 1188

**PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS**

DO NOT apply under weather conditions, or from spray equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures. DO NOT plant crops other than maize, sweet corn, tobacco, cotton, soybeans, sunflowers, peanuts, broccoli, Brussels sprouts, cabbages, cauliflowers or Concep II treated sorghum within 6 months of application of Kenso Agcare Metoken Gold Herbicide.

**PROTECTION OF LIVESTOCK**

DO NOT graze or cut sweet potato leaves for stock food.

**PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT**

TOXIC TO FISH. DO NOT contaminate streams, rivers or waterways with the chemical or used containers. DO NOT apply under meteorological conditions or from spraying equipment which could be expected to cause spray to drift onto adjacent areas, particularly wetlands, waterbodies or watercourses.

**STORAGE AND DISPOSAL**

Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight.

**Refillable containers:** Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

**Other containers:** Triple-rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

**SAFETY DIRECTIONS**

Will damage eyes. Repeated exposure may cause allergic disorders. Sensitive workers should use protective clothing. Avoid contact with eyes. When opening the container and preparing spray, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length pvc gloves, and goggles. When using the prepared spray, wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow-length PVC gloves and face shield or goggles. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves, and face shield or goggles, and contaminated clothing.

**FIRST AID**

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766.

**SAFETY DATA SHEET**

Additional information is listed in the Safety Data Sheet which can be obtained from [www.kenso.com.au](http://www.kenso.com.au).

**CONDITIONS OF SALE**

Kenso Corporation (M) Sdn. Bhd. will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Kenso Corporation (M) Sdn. Bhd. is limited to the replacement of the goods or (at the option of Kenso Corporation (M) Sdn. Bhd.) the refund of the price paid and where possible sufficient part of the goods to enable proper examination being returned to Kenso Corporation (M) Sdn. Bhd. within thirty days of delivery.

In a Transport Emergency  
Dial **000** Police of Fire Brigade

® Not a registered trademark of Kenso Corporation



**Kenso Corporation (M) Sdn. Bhd.**  
Level 1, 98 Commercial Road,  
Teneriffe QLD 4005  
Phone (07) 3216 1188

**APVMA Approval No.: 68625/127089**

**DIRECTIONS FOR USE**

**Restraints:** DO NOT apply by aircraft.  
DO NOT apply to waterlogged soils.  
DO NOT apply if heavy rains or storms that are likely to cause run-off are forecast within 2 days of application.  
DO NOT irrigate to the point of run-off for at least 2 days after application.

Crop	Weeds Controlled	State	Rate	Critical Comments
<b>Broccoli, Brussels Sprouts, Cabbages, Cauliflowers</b>	Barnyard Grass, Blackberry Nightshade, Chickweed (suppression), Deadnettle, Fat Hen, Love Grass, Pigeon Grass, Pigweed (suppression), Potato Weed, Shepherd's Purse, Stinging Nettle, Summer Grass, Wireweed (suppression)	Qld, NSW, Vic, Tas, NT only	1.5 to 2 L per ha	Apply immediately after transplanting. Sufficient irrigation to wet the top 3 to 4 cm of soil should be applied within 24 hours. Use rates towards the higher end of the range where Blackberry Nightshade, Chickweed, Fat Hen, Love Grass, Pigeon Grass, Pigweed, Potato Weed and Wireweed are the major problem. <b>See General Instructions for Warning - Brassica Crops.</b>
	Crowsfoot Grass, Fierce Thornapple (suppression), Liverseed Grass, Winter Grass	NSW only		Apply immediately after transplanting. Sufficient irrigation to wet the top 3 to 4 cm of soil should be applied within 24 hours. Use rates towards the higher end of the range where Apple of Peru, Common Sowthistle, Fierce Thornapple, Liverseed Grass, Redroot Amaranth, Twiggy Turnip and Nettleleaf Goosefoot are the major problems.
	Apple of Peru, Awnless Barnyard Grass, Fierce Thornapple (suppression), Liverseed Grass, Yellow Weed, Redroot Amaranth (suppression)	Qld only		Apply immediately after transplanting. Sufficient irrigation to wet the top 3 to 4 cm of soil should be applied within 24 hours. Use rates towards the higher end of the range where Apple of Peru, Common Sowthistle, Fierce Thornapple, Liverseed Grass, Redroot Amaranth, Twiggy Turnip and Nettleleaf Goosefoot are the major problems. <b>WARNING:</b> Qld only. Stunting and reduced yield may occur in broccoli in red soils and light alluvial soils. <b>See General Instructions for Warning - Brassica Crops.</b>
<b>Brassica leafy vegetables</b> including Chinese white cabbage, Chinese flowering cabbage, Chinese broccoli, Indian mustard cabbage, Chinese cabbage, Mizuna greens, Red mustard, Curled mustard, Chinese flat cabbage, Mustard greens, Turnip greens, Rutabaga greens, and Mibuna greens)	Annual Ryegrass, Common Cotula, Common Sowthistle (suppression), Crowsfoot Grass, Deadnettle, Fat Hen, Fiddle Dock, Fierce Thornapple (suppression), Italian Ryegrass, Liverseed Grass, Love Grass, Nettleleaf Goosefoot (suppression), Pigeon Grass, Pigweed (suppression), Potato Weed, Redroot Amaranth (suppression), Shepherd's Purse, Stinging Nettle, Summer Grass, Twiggy Turnip (suppression), Winter Grass, Wireweed (suppression), Yellow Weed	All States	1.0 to 2 L per ha	Apply one treatment only, immediately after transplanting. Apply in a minimum of 60 L water per hectare. Apply sufficient irrigation after application to wet the upper 3 to 4 cm of soil with 24 hours. Use rates towards the higher end of the range where blackberry nightshade, chickweed, fat hen, love grass, pigweed, potato weed and wire weed are major problems (Refer Brassica use pattern on product label). <b>See general instructions for warning - brassica crops</b>
	Annual Ryegrass, Apple of Peru, Awnless Barnyard Grass, Barnyard Grass, Blackberry Nightshade, Chickweed (suppression), Common Cotula, Common Sowthistle (suppression), Crowsfoot Grass, Deadnettle, Fat Hen, Fiddle Dock, Fierce Thornapple (suppression), Italian Ryegrass, Liverseed Grass, Love Grass, Nettleleaf Goosefoot (suppression), Pigeon Grass, Pigweed (suppression), Potato Weed, Redroot Amaranth (suppression), Shepherd's Purse, Stinging Nettle, Summer Grass, Twiggy Turnip (suppression), Winter Grass, Wireweed (suppression), Yellow Weed	All States	0.15 to 0.25 L per ha	Apply at or immediately after planting and before crops and weeds emerge. Apply to moist soil. Use rates toward the higher end of the range where longer residual control is required. <b>See General Instructions for use in zero-tillage systems.</b>
<b>Canola and mustard</b> (oilseed cultivars) ( <i>Brassica juncea</i> )	Toad Rush	All States	0.15 to 0.25 L per ha	Apply at or immediately after planting and before crops and weeds emerge. Apply to moist soil. Use rates toward the higher end of the range where longer residual control is required. <b>See General Instructions for use in zero-tillage systems.</b>
<b>Cotton</b>	Barnyard Grass, Crowsfoot Grass, Liverseed Grass, Love Grass, Pigeon Grass, Summer Grass, Wandering Jew	Qld, NSW, WA only	1 L per ha	<b>Pre-emergent:</b> Apply before, at or immediately after planting and before crops and weeds have germinated using a low volume boom sprayer applying 60 to 120 L of mixture per ha. Rain or irrigation is necessary within 10 days of spraying to thoroughly wet the top 3 to 4 cm of soil. If rain does not occur or irrigation is unavailable, incorporate chemical to a depth of 3 to 4 cm. Use rates towards the higher end of the range on heavy soils or where a heavy grass population is expected. DO NOT throw untreated soil onto treated areas as this will reduce weed control. DO NOT use on soils where the sand content exceeds 70%. Use lower rate where the sand content of the soil is within the range 50% to 70%. <b>See General Instructions for Warning - Cotton.</b>

Crop	Weeds Controlled	State	Rate	Critical Comments
<b>Culinary herbs:</b> basil, bay leaves, borage, chives, coriander, dill, fennel, lemon balm, lemon grass, kaffir lime leaves, marigold flowers, marjoram (oregano), mints, nasturtium leaves, parsley, rosemary, sage, salad burnett, sorrel, tarragon, thyme <b>Savoury root vegetable:</b> galangal <b>Leafy vegetables:</b> rucola (rocket), chervil, mizuna. <b>Teas:</b> lemon verbena <b>Spice:</b> turmeric. <b>Edible flowers:</b> dianthus, roses.	Annual Ryegrass, Apple of Peru, Awnless Barnyard Grass, Barnyard Grass, Blackberry Nightshade, Chickweed (suppression), Common Cotula, Common Sowthistle (suppression), Crowsfoot Grass, Deadnettle, Fat Hen, Fiddle Dock, Fierce Thornapple (suppression), Italian Ryegrass, Liverseed Grass, Love Grass, Nettleleaf Goosefoot (suppression), Pigeon Grass, Pigweed (suppression), Potato Weed, Redroot Amaranth (suppression), Shepherd's Purse, Stinging Nettle, Summer Grass, Twiggy Turnip (suppression), Winter Grass, Wireweed (suppression), Yellow Weed	All States	1.5 - 2.0 L/ha	Apply before, at, or immediately after planting/transplanting and before crops and weeds have germinated.  Sufficient irrigation to wet the top 3-4 cm of soil should be applied within 24 hours.  Do not use more than once per season
	<b>Green Beans, Navy Beans</b>	Blackberry Nightshade, Common Sowthistle, Fat Hen, Potato Weed, Barnyard Grass, Crowsfoot Grass, Liverseed Grass, Love Grass, Pigeon Grass, Summer Grass and suppression of Pigweed	All States	1.5 to 2L per ha
	Powell's Amaranth	Tas only	2 L per ha	<b>Pre-emergent:</b> Apply before, at or immediately after planting and before crops and weeds have germinated. For sufficient Amaranth control, the Metoken Gold application should be followed by 1 L/ha Basagran® (2 leaf Amaranth) or 2 L/ha Basagran® (4 leaf Amaranth). Rain or irrigation is necessary within 10 days of spraying to thoroughly wet the top 3 to 4 cm of soil. If rain does not occur or irrigation is unavailable, incorporate chemical to a depth of 3 to 4 cm. DO NOT throw untreated soil onto treated areas as this will reduce weed control.
<b>Maize, Sweet Corn</b>	Barnyard Grass, Crowsfoot Grass, Liverseed Grass, Love Grass, Pigeon Grass, Summer Grass, Wandering Jew	All States	1 to 2 L per ha	<b>Pre-emergent:</b> Apply before, at or immediately after planting and before crops and weeds have germinated using a low volume boom sprayer applying 60 to 120 L of mixture per ha. Rain or irrigation is necessary within 10 days of spraying to thoroughly wet the top 3 to 4 cm of soil. If rain does not occur or irrigation is unavailable, incorporate chemical to a depth of 3 to 4 cm. Use rates towards the higher end of the range on heavy soils or where a heavy grass population is expected. DO NOT throw untreated soil onto treated areas as this will reduce weed control. DO NOT use on soils where the sand content exceeds 70%. Use lower rate where the sand content of the soil is within the range 50% to 70%.
	Blackberry Nightshade, Caltrop, Common Thornapple, Mintweed, Needle Burr, Noogoora Burr, Pigweed, Redroot Amaranth, Slim Amaranth		1 to 2 L plus 2.5 L of Flowable Gesaprim (500 g/L) per ha	
<b>Pastures -</b> Trikkala Sub Clover, Haifa, White Clover, Maral Persian Clover, Paradana Balansa Clover, Redquin Red Clover, Phalaris and Cocksfoot	Toad Rush	NSW, Vic, SA, WA, Tas, NT only	200 to 250 mL per ha	Apply at or immediately after planting and before pasture species and weeds emerge. Application should be made to a moist seed bed. Sufficient rain to thoroughly wet the top 3 to 4 cm of soil should occur within 10 days after spraying. Use rates towards the higher end of the range in wet areas or when longer residual control is required. DO NOT use in soils where prolonged waterlogging is anticipated otherwise unacceptable crop retardation will occur. Avoid damage to pasture species by avoiding overlaps and double spraying in headlands where White, Persian and Balansa Clovers are sown and if the 250 mL rate is being used.

Crop	Weeds Controlled	State	Rate	Critical Comments
<b>Peanuts</b>	Barnyard Grass, Crowsfoot Grass, Liverseed Grass, Love Grass, Pigeon Grass, Summer Grass, Wandering Jew	Qld, NSW, WA, NT only	1 to 2 L per ha	Apply before pre-irrigation where soil crusting is likely. Apply before, at or immediately after planting and before weeds germinate. Rain or irrigation is necessary within 10 days of spraying to thoroughly wet the top 3 to 4 cm of soil. If rain does not occur or irrigation is unavailable, incorporate chemical to a depth of 3 to 4 cm. DO NOT throw untreated soil onto treated areas as this will reduce weed control. Use rates toward the higher end of the range on heavy soils or where a heavy grass population is expected.
<b>Soybeans, Sunflowers</b>		All States		
<b>Rhubarb</b>	Annual Ryegrass, Apple of Peru, Awnless Barnyard Grass, Barnyard Grass, Blackberry Nightshade, Chickweed (suppression), Common Cotula, Common Sowthistle (suppression), Crowsfoot Grass, Deadnettle, Fat Hen, Fiddle Dock, Fierce Thornapple (suppression), Italian Ryegrass, Liverseed Grass, Love Grass, Nettleleaf Goosefoot (suppression), Pigeon Grass, Pigweed (suppression), Potato Weed, Redroot Amaranth (suppression), Shepherd's Purse, Stinging Nettle, Summer Grass, Twiggy Turnip (suppression), Winter Grass, Wireweed (suppression), Yellow Weed	All States	1.0 to 2 L per ha	Pre-emergent: Apply one (1) application only, immediately before or after transplanting crowns and before weeds have germinated. Sufficient irrigation to wet the soil through the weed zone should be applied within 24 hours. Further weed germination may occur following re-hilling due to exposure of untreated soil.
<b>Spinach and silverbeet</b>	Annual Ryegrass, Apple of Peru, Awnless Barnyard Grass, Barnyard Grass, Blackberry Nightshade, Chickweed (suppression), Common Cotula, Common Sowthistle (suppression), Crowsfoot Grass, Deadnettle, Fat Hen, Fiddle Dock, Fierce Thornapple (suppression), Italian Ryegrass, Liverseed Grass, Love Grass, Nettleleaf Goosefoot (suppression), Pigeon Grass, Pigweed (suppression), Potato Weed, Redroot Amaranth (suppression), Shepherd's Purse, Stinging Nettle, Summer Grass, Twiggy Turnip (suppression), Winter Grass, Wireweed (suppression), Yellow Weed	All States	1.5 – 2.0 L/ha	Apply with a boom spray, in a minimum of 60 L water/ha. Apply immediately before, at, or immediately after sowing. If transplanting the crop, apply immediately before transplanting. DO NOT apply herbicide over the transplants. Crop retardation may occur where the herbicide is used on soils that have low organic matter content and which contain more than 60% of fine sand and silt. On these soil types, use rates towards the low end of the rate range, and apply to moist soil. Irrigation after application on these types of soils should be limited to no more than 25 mm. Also, on these soils, DO NOT use more than once per year (as on product label for brassicas); For other relevant instructions, see Broccoli (etc) section of Directions for Use table.
Spring onions ( <i>Allium fistulosum</i> ) and Shallots ( <i>Allium cepa</i> var. <i>aggregatum</i> )	Annual Ryegrass, Apple of Peru, Awnless Barnyard Grass, Barnyard Grass, Blackberry Nightshade, Chickweed (suppression), Common Cotula, Common Sowthistle (suppression), Crowsfoot Grass, Deadnettle, Fat Hen, Fiddle Dock, Fierce Thornapple (suppression), Italian Ryegrass, Liverseed Grass, Love Grass, Nettleleaf Goosefoot (suppression), Pigeon Grass, Pigweed (suppression), Potato Weed, Redroot Amaranth (suppression), Shepherd's Purse, Stinging Nettle, Summer Grass, Twiggy Turnip (suppression), Winter Grass, Wireweed (suppression), Yellow Weed	All States	1 L/ha	DIRECT SEEDED CROPS: Apply one application before, at or immediately after planting and before crops and weeds have germinated;  TRANSPLANTED CROPS: Apply one application before planting and before weeds have germinated.

Crop	Weeds Controlled	State	Rate	Critical Comments
<b>Sorghum treated with Concep II Sorghum Seed Safener</b>	Barnyard Grass, Crowsfoot Grass, Liverseed Grass, Love Grass, Pigeon Grass, Summer Grass, Wandering Jew	Qld, NSW, WA, NT only	1 to 2 L per ha	<b>Pre-emergent:</b> Apply before, at or immediately after planting and before crops and weeds have germinated using a low volume boom sprayer applying 60 to 120 L of mixture per ha. DO NOT apply to sorghum seed not previously treated with Concep II Sorghum Seed Safener. Rain or irrigation is necessary within 10 days of spraying to thoroughly wet the top 3 to 4 cm of soil. If rain does not occur or irrigation is unavailable, incorporate chemical to a depth of 3 to 4 cm. Use rates toward the higher end of the range on heavy soils or where a heavy grass population is expected. DO NOT throw untreated soil onto treated areas as this will reduce weed control. Use the lower rate on sandy soils.
	Blackberry Nightshade, Caltrop, Common Thornapple, Mintweed, Needle Burr, Noogoora Burr, Pigweed, Redroot Amaranth, Slim Amaranth		1 to 2 L plus 2.5 L of Flowable Gesaprim (500 g/L) per ha	
<b>Sugar Cane</b> (plant and ratoon)	Awnless Barnyard Grass, Barnyard Grass, Bellvine, Billygoat Weed (Blue Top), Blackberry Nightshade, Caltrop, Common Sowthistle (Milk Thistle), Common Thornapple, Crowsfoot Grass, Green Summer Grass, Guinea Grass, Liverseed Grass, Mexican Clover, Mintweed, Needle Burr, Passionfruit vines, Pigeon Grass, Pigweed, Redroot Amaranth, Square Weed, Star of Bethlehem, Summer Grass, Wild Rose	Southern Qld only (Bundaberg South)	1.1 to 1.45 L plus 2.7 to 3.6 L Flowable Gesaprim 500 SC Or 1.5 to 2 kg of Atrazine 900 WG	Apply as a pre- or post emergent application to crop. Apply once per year only. Application should be made to a moist soil before weeds and grasses have germinated. If conditions remain dry for a period of 10 days after spraying, irrigation or a shallow cultivation (2.5 cm) may assist results. DO NOT throw untreated soil onto treated areas as this will reduce weed control. In Southern Queensland, use rates toward the higher end of the range on heavy soils where a high grass population is expected. In Northern Queensland, application must be made to moist soil and rainfall or irrigation should occur within 24 hours of application. Use rates toward the higher end of the range where high Green Summer Grass population is expected. Where broadleaf weeds and grasses have emerged and are in the 2 to 4 true leaf stage, knockdown herbicides such as Primatol® Z (2 L/ha), a flowable diuron (500 g/L) at 1 L/ha or Para-Ken 250 (1 L/ha), applied as a post directed spray, can be added. Where broadleaf weeds only have emerged and are at the 2 to 4 true leaf stage then a 500 g/L 2,4-D amine at 2 L/ha should be added. In all cases add a non-ionic surfactant. Follow all instructions and restrictions on the Atrazine labels.
		Northern Qld only (Mackay North)	1.45 to 1.8 L plus 3.6 to 4.5 L Flowable Gesaprim 500 SC Or 2 to 2.5 kg of Atrazine 900 WG	
<b>Sweet Potatoes</b>	Black Pigweed, Red Shank, Potato Weed, Barnyard Grass, Crowsfoot Grass, Liverseed Grass, Love Grass, Pigeon Grass, Summer Grass, suppression of Evening Primrose	Qld, NSW, Vic, SA, NT only	1.5 L per ha	<b>Pre-emergent:</b> Apply immediately after transplanting before weeds have germinated. Sufficient irrigation to wet the soil through the weed zone should be applied within 24 hours. Further weed germination may occur following rehilling due to exposure of untreated soil.
<b>Barley, Oats</b>	Suppression of Annual Ryegrass	NSW, ACT, Vic, Tas, SA, WA only	375 to 500 mL per ha	Apply to moist soils immediately after planting and before crop and weeds emerge. Use the lower rate on light soils. Application should be made to a moist seed bed and sufficient rain to thoroughly wet the top 3 to 4 cm of soil should occur within 10 days after spraying. Damage may occur if crop is sown too shallow (less than 4 cm). Damage can occur on light sandy soil types after heavy rain. <b>See General Instructions for use in zero-tillage systems.</b>

Crop	Weeds Controlled	State	Rate	Critical Comments
<b>Tobacco</b> (transplanted in field)	Barnyard Grass, Love Grass, Pigeon Grass, Summer Grass and suppression of Common Thornapple ( <i>Datura stramonium</i> ) and Nightshades ( <i>Solanum sarrachoides</i> and <i>S. gigrum</i> )	NSW, Vic only	1.5 L per ha	Apply by low volume boom spray prior to transplanting and before weeds and grasses germinate. The seed bed should be fine and free of large clods. Thoroughly incorporate within 7 days of spraying using a rotary hoe or similar equipment set to work to a depth of 7.5 cm. DO NOT use disc cultivating implements. An even coverage of the herbicide is essential. The herbicide needs to be activated by moisture through to the top 3 to 4 cm of soil at the time of application or within 4 days of incorporation. This can be achieved by applying to moist soil, irrigating soon after incorporation or irrigating immediately after transplanting. Avoid boom overlaps as double treatment may cause growth suppression. Temporary crop retardation may be observed when soil conditions remain dry after application or where Black Root Rot may occur.
<b>Wheat, Barley, Oats, Triticale</b>	Toad Rush	NSW, ACT, Vic, Tas, SA, WA only	150 to 250 mL per ha	Apply at or immediately after planting and before crop and weeds emerge. Application should be made to a moist seed bed and sufficient rain to thoroughly wet the top 3 to 4 cm of soil should occur within 10 days after spraying. Use rates toward the higher end of the range in wet areas or when longer residual control is required or in situations where crop competition is minimal. When mixed with Ken-gran 750 WG apply pre-sowing and incorporate by sowing operation (for wheat only). <b>See General Instructions for use in zero-tillage systems.</b>

**NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

#### WITHHOLDING PERIODS

Canola, Mustard: DO NOT GRAZE TREATED FIELDS FOR 10 WEEKS AFTER APPLICATION.

Green beans and Navy beans: DO NOT HARVEST BEANS FOR 8 WEEKS AFTER APPLICATION.

DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 10 WEEKS AFTER APPLICATION.

Pastures: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 13 WEEKS AFTER APPLICATION.

Sweet Potatoes: DO NOT HARVEST, GRAZE OR CUT FOR STOCK FOOD FOR 23 WEEKS AFTER APPLICATION.

Wheat, Barley, Oats, Triticale: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 8 WEEKS AFTER APPLICATION.

Brassica leafy vegetables, Rhubarb: NOT REQUIRED WHEN USED AS DIRECTED.

Spinach and silverbeet: (HARVEST) NOT REQUIRED WHEN USED AS DIRECTED.

DO NOT GRAZE OR USE FOR STOCKFEED FOR 8 WEEKS AFTER APPLICATION.

Spring onions and shallots: (HARVEST) NOT REQUIRED WHEN USED AS DIRECTED.

Herbs, chervil, mizuna, rucola, lemon verbena: DO NOT HARVEST FOR AT LEAST 8 WEEKS AFTER APPLICATION.

Galangal, turmeric: DO NOT HARVEST FOR AT LEAST 23 WEEKS AFTER APPLICATION.

HARVEST with-hold for all other crops: NOT REQUIRED WHEN USED AS DIRECTED

FORAGE with-hold for all other crops (except Sweet Potatoes): DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 13 WEEKS AFTER APPLICATION.

#### GENERAL INSTRUCTIONS

### GROUP 15 HERBICIDE

#### RESISTANCE WEEDS WARNING

Kenso Agcare Metoken Gold Herbicide is a member of the amide group of herbicides. The product has the herbicides with diverse sites of action mode of action (Chloroacetamides Inhibitors of cell division / Inhibitors of very long chain fatty acids (VLCFA inhibitors). For weed resistance management, the product is a Group 15 herbicide.

Some naturally-occurring weed biotypes resistant to the product and other Group 15 herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by this product or other Group 15 herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, Kenso Corporation (M) Sdn Bhd accepts no liability for any losses that may result from the failure of this product to control the resistant weeds.

Advice as to strategies and alternative treatments that can be used should be obtained from your local supplier, consultant, local Department of Agriculture, Primary Industries Department or a Kenso Agcare representative.

#### Uses

The product is a short residual, pre-emergent herbicide for the control of many important annual grasses and broadleaf weeds in the crops nominated. The product does not control the important broadleaf weeds in maize, sweet corn and Concep II treated sorghum but, when mixed with Flowable Gesaprim®, control of many important annual weeds and grasses is obtained without the problem of carryover of herbicides affecting succeeding crops. The product acts mainly through shoot (coleoptile) absorption. Its effectiveness depends on the occurrence of rainfall or irrigation to move it down into the top 3 to 4 cm of soil, where it must be present at the time of weed germination.

#### Application

**Ground Application:** Apply in a minimum of 60 L/ha of water.

#### Mixing

Fill the spray tank to one quarter full. Then add Kenso Agcare Metoken Gold Herbicide while adding water to make up the final spray volume. Pour Kenso Agcare Metoken Gold Herbicide into the stream of incoming water. Good agitation at the time Kenso Agcare Metoken Gold Herbicide is added to the tank will ensure good mixing.

**When tank mixing:** Wettable powder or water dispersible granule formulations should be added to the tank first, followed by suspension concentrates (flowables), water soluble salts then Kenso Agcare Metoken Gold Herbicide or other emulsifiable concentrate formulations. Maintain thorough agitation during mixing and application. Agitate tank-mixes vigorously if allowed to stand.

NB: Tank-mix spray solution should NOT be left standing in the vat overnight.

#### Compatibility

Kenso Agcare Metoken Gold Herbicide is compatible with Flowable Cotogard®, Flowable Cotoran®, Flowable Gesagard®, Flowable Gesaprim®, Atrazine 900 WG, Ken-Gran 750 WG, Basagran®, Speedy 250 and Ken-Up.

**WARNING - Cotton:** DO NOT use when planting into dry soil or soil with marginal moisture. Under dry planting conditions heavy rain or flood irrigation soon after planting, together with below average soil temperatures, can result in crop injury due to the inability of the germinating seedling to metabolise above normal amounts of the herbicide that may accumulate in the root zone as a result of accelerated capillary movement. Kenso Agcare Metoken Gold Herbicide should not be used in situations where these conditions are likely to occur.

**WARNING - Brassica Crops:** Crop retardation may occur where Kenso Agcare Metoken Gold Herbicide is used on soils that have a low organic matter content and which contain more than 60% of fine sand and silt. On these soil types, use rates of Kenso Agcare Metoken Gold Herbicide towards the lower end of the rate range and apply onto moist soil. Irrigation after application should be limited to no more than 25 mm. DO NOT use more than once per year.

**Zero-tillage (Canola, Barley, Oats, Wheat, Triticale):** When sowing using knife points and zero-tillage, and leaving a seeding slot, do not apply post-seeding unless harrows (or similar) have filled-in the seeding slot before application. Alternatively, apply Kenso Agcare Metoken Gold Herbicide immediately before seeding.

#### PRECAUTION

##### Re-entry Period

Do not allow entry into treated areas until the spray has dried.

##### To avoid Crop Damage

As metolachlor is known to cause some crop damage in sandy-sandy loam soils with low organic matter, it is recommended that growers test the safety of metolachlor on a small area of their crop before using the product on larger areas. If any adverse crop symptoms are observed within 14 days of application in which irrigation or rainfall has occurred, then the product should not be used.

##### Trade Advice: Export of treated produce

Growers should note that maximum residue limits (MRLs) or import tolerances may not exist in all markets for edible produce treated with this product. If you are growing edible produce for export, please check with Kenso Agcare Pty Ltd for the latest information on MRLs and import tolerances before using this product.