

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

KENSO AGCARE

AVIOR 800 WG FUNGICIDE

ACTIVE CONSTITUENT: 800 g/kg AZOXYSTROBIN

GROUP **11** FUNGICIDE

For the control of various diseases of grapes, potatoes, tomatoes, cucurbits, avocados, mangoes, passionfruit and poppies as per the Directions for Use table.

Kenso Corporation (M) Sdn Bhd
Level 1, 98 Commercial Road,
Teneriffe QLD 4005
Phone (07) 3216 1188
www.kenso.com.au



IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USE

CONTENTS: 1 Kilograms APVMA Approval No.: 67305/ 55736

KENSO AGCARE AVIOR 800 WG FUNGICIDE

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight.

Single rinse before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. Puncture and bury empty bags in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty bags and product should not be burnt.

SAFETY DIRECTIONS

Harmful if inhaled. May irritate the skin. Will irritate the eyes. DO NOT inhale spray mist. Avoid contact with eyes and skin. When opening the container and mixing and loading the product, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), face shield or goggles and elbow-length chemical resistant gloves. When using the prepared spray, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), face shield or goggles and elbow-length chemical resistant gloves. If product in eyes, wash it out immediately with water. After each day's use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia 13 11 26).

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet, which can be obtained from www.kenso.com.au.

CONDITIONS OF SALE

Kenso Corporation (M) Sdn. Bhd. will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Kenso Corporation (M) Sdn. Bhd. is limited to the replacement of the goods or (at the option of Kenso Corporation (M) Sdn. Bhd.) the refund of the price paid and where possible sufficient part of the goods to enable proper examination being returned to Kenso Corporation (M) Sdn. Bhd. within thirty days of delivery.

In a Transport Emergency
Dial **000** Police or Fire Brigade



Additional statements required by Globally Harmonised System of Classification and Labelling of Chemicals (GHS) and Safe Work Australia: **TOXIC IF INHALED.** *Precautionary:* Avoid breathing dust/fume/gas/mist/vapours/spray. Use only outdoors or in a well-ventilated area. IF INHALED: Remove person to fresh air and keep comfortable for breathing. Call a POISON CENTER or doctor/physician. Specific treatment (see FIRST AID on this label). Store in a well-ventilated place. Keep container tightly closed. Store locked up.

Batch No.:

Date of Manufacture:

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

KENSO
agrosol
AVIOR
800 WG

Fungicide
ACTIVE CONSTITUENT: 800 g/kg
AZOXYSTROBIN

GROUP **11** FUNGICIDE

For the control of various diseases of grapes, potatoes, tomatoes, cucurbits, avocados, mangoes, passionfruit and poppies as per the Directions for Use table.

READ THE LEAFLET BEFORE USE

APVMA Approval No.: 67305/55736



Kenso Corporation (M) Sdn. Bhd.
Level 1, 98 Commercial Road,
Teneriffe QLD 4005
Phone (07) 3216 1188

DIRECTIONS FOR USE

RESTRAINTS

DO NOT apply by air, except on potatoes.

TREE AND VINE CROPS

Rate				Critical Comments
In the following table: Tree and Vine Crops all rates given are for dilute spraying. For concentrate spraying, refer to the Application section.				For all uses in the table Tree and Vine Crops: Apply by dilute or concentrate spraying equipment. Apply the same total amount of product to the target crop whether applying this product by dilute or concentrate spraying methods. When applying Kenso Agcare Avior 800 WG Fungicide through low volume application equipment, DO NOT use a concentrate factor greater than 4X. In these cases adequate coverage of all plant surfaces is still required to achieve control of diseases.
Crop	Disease	Rate/100L	WHP	Critical Comments
Avocado	Stem End Rot, Anthracnose	25g	7 days	Apply Kenso Agcare Avior 800 WG Fungicide in a preventative fungicide program containing fungicides from a different chemical group. For best results commence the disease control program with an approved fungicide from an alternative chemical group, then apply 1 application of Kenso Agcare Avior 800 WG Fungicide during early fruit set. Follow with applications of an approved fungicide from a different chemical group. Apply 2 final applications of Kenso Agcare Avior 800 WG Fungicide at 14 to 28 day intervals with the final spray applied 7 days prior to harvest. Ensure thorough spray coverage. Resistance Management Disease control may be reduced if strains of pathogens less sensitive to Kenso Agcare Avior 800 WG Fungicide develop. DO NOT use Kenso Agcare Avior 800 WG Fungicide curatively. If consecutive applications of Kenso Agcare Avior 800 WG Fungicide are used they must be followed by at least the same number of applications of fungicide(s) from a different group(s) before Kenso Agcare Avior 800 WG Fungicide is used again in that crop in the current or following season. Apply a maximum of 3 applications of Kenso Agcare Avior 800 WG Fungicide per season. DO NOT apply more than 2 consecutive applications of Kenso Agcare Avior 800 WG Fungicide. DO NOT start the disease control program with Kenso Agcare Avior 800 WG Fungicide.
Grapes Table, wine, dried	Powdery Mildew (<i>Uncinular necator</i>), Downy Mildew (<i>Plasmopara viticola</i>), Botrytis Bunch Rot † (<i>Botrytis cinerea</i>)	23.5 to 31.5 g	14 days	Application Method and Rate Apply in a sufficient volume of water to achieve thorough coverage of all foliage and fruit. The volume of water required to achieve this will depend on the stage of vine growth and vigour. Ensure thorough coverage. Adjust spray nozzles to direct spray droplets to the canopy present. Apply the higher rate of application in the following circumstances: 1. Where humid conditions favour Powdery Mildew infection, particularly on susceptible varieties. 2. At the start of the season when there has been a heavy carry over of Powdery Mildew infection (flag shoots are present). Spray Timing and Interval Apply 2 or 3 consecutive applications at 10 to 16 day intervals at any time between early shoot growth and 14 days before harvest. Use the recommended shorter interval during periods when climatic conditions are favourable for disease infection. † Botrytis Bunch Rot Kenso Agcare Avior 800 WG Fungicide must not be used alone for Botrytis control at critical times such as 80 to 100% capfall and pre-harvest. It must be tank mixed with or substituted by a specific Botryticide at these critical times. When Kenso Agcare Avior 800 WG Fungicide is used in a seasonal spray programme it will provide control of Botrytis additional to that of ® specific Botryticides such as Bravo . Resistance Management Disease control may be reduced if strains of pathogens less sensitive to Kenso Agcare Avior 800 WG Fungicide develop. DO NOT use Kenso Agcare Avior 800 WG Fungicide curatively. As a precaution, DO NOT apply more than a total of 3 applications of Kenso Agcare Avior 800 WG Fungicide per crop in 1 season. If consecutive applications of Kenso Agcare Avior 800 WG Fungicide are used they must be followed by at least the same number of applications of fungicide(s) from a different group(s) before Kenso Agcare Avior 800 WG Fungicide is used again in that crop in the current or following season. DO NOT use Kenso Agcare Avior 800 WG Fungicide for disease control in grapevine nurseries.
Mango	Stem End Rot, Anthracnose	25g	3 days	Apply Kenso Agcare Avior 800 WG Fungicide in a preventative fungicide program containing fungicides from a different chemical group. For best results apply 1 to 2 applications of Kenso Agcare Avior 800 WG Fungicide at flowering and early fruit set, at no less than 14 day intervals. Follow with applications of an approved fungicide from a different chemical group. Further applications of Kenso Agcare Avior 800 WG Fungicide may be applied at 21 days and 3 to 7 days prior to harvest. Ensure thorough spray coverage. Resistance Management Disease control may be reduced if strains of pathogens less sensitive to Kenso Agcare Avior 800 WG Fungicide develop. DO NOT use Kenso Agcare Avior 800 WG Fungicide curatively. If consecutive applications of Kenso Agcare Avior 800 WG Fungicide are used they must be followed by at least the same number of applications of fungicide(s) from a different group(s) before Kenso Agcare Avior 800 WG Fungicide is used again in that crop in the current or following season. Apply a maximum of 3 applications of Kenso Agcare Avior 800 WG Fungicide per season. DO NOT apply more than 2 consecutive applications of Kenso Agcare Avior 800 WG Fungicide DO NOT start the disease control program with Kenso Agcare Avior 800 WG Fungicide.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolong periods in direct sunlight.

Single rinse before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. Puncture and bury empty bags in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty bags and product should not be burnt.

SAFETY DIRECTIONS

Harmful if inhaled. May irritate the skin. Will irritate the eyes. DO NOT inhale spray mist. Avoid contact with eyes and skin. When opening the container and mixing and loading the product, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), face shield or goggles and elbow-length chemical resistant gloves. When using the prepared spray, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), face shield or goggles and elbow-length chemical resistant gloves. If product in eyes, wash it out immediately with water. After each day's use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia 13 11 26).

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet, which can be obtained from www.kenso.com.au.

CONDITIONS OF SALE

Kenso Corporation (M) Sdn. Bhd. will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Kenso Corporation (M) Sdn. Bhd. is limited to the replacement of the goods or (at the option of Kenso Corporation (M) Sdn. Bhd.) the refund of the price paid and where possible sufficient part of the goods to enable proper examination being returned to Kenso Corporation (M) Sdn. Bhd. within thirty days of delivery.

® Not a trademark of Kenso Corporation (M) Sdn Bhd

In a Transport Emergency
Dial **000**
Police of Fire Brigade



Kenso Corporation (M) Sdn. Bhd.
Level 1, 98 Commercial Road,
Teneriffe QLD 4005
Phone (07) 3216 1188

APVMA Approval No.: 67305/55736

Crop	Disease	Rate/100L	WHP	Critical Comments
Passion fruit	Alternaria, Cladosporium	25g	1 day	Apply Kenso Agcare Avior 800 WG Fungicide in a preventative fungicide program containing fungicides from a different chemical group. For best results apply 2 to 3 applications of Kenso Agcare Avior 800 WG Fungicide at 14 day intervals over flowering. Follow with applications of an approved fungicide from a different chemical group. Apply a further 1 to 2 applications of Kenso Agcare Avior 800 WG Fungicide finishing 1 day prior to harvest. Ensure thorough spray coverage. Resistance Management Disease control may be reduced if strains of pathogens less sensitive to Kenso Agcare Avior 800 WG Fungicide develop. DO NOT use Kenso Agcare Avior 800 WG Fungicide curatively. DO NOT apply more than 1/3 of the total number of fungicide applications. For example, if the total number of fungicides applied to the crop is 9, DO NOT apply more than 3 applications of Kenso Agcare Avior 800 WG Fungicide. DO NOT exceed 5 applications of Kenso Agcare Avior 800 WG Fungicide per crop. DO NOT apply more than 2 consecutive applications of Kenso Agcare Avior 800 WG Fungicide. DO NOT start the disease control program with Kenso Agcare Avior 800 WG Fungicide.
Poppies	Downy Mildew	235 g/ha	6 weeks	Apply Kenso Agcare Avior 800 WG Fungicide in a preventative fungicide program containing fungicides from a different chemical group. Apply Kenso Agcare Avior 800 WG Fungicide preventatively before disease symptoms appear. Ensure thorough spray coverage. Resistance Management Disease control may be reduced if strains of pathogens less sensitive to Kenso Agcare Avior 800 WG Fungicide develop. DO NOT use Kenso Agcare Avior 800 WG Fungicide curatively. Apply a maximum of 2 applications of Kenso Agcare Avior 800 WG Fungicide per crop.
Potatoes	Early Blight (Target Spot) (<i>Alternaria solani</i>) Late Blight (<i>Phytophthora infestans</i>)	95 to 125 g/ha 155 to 190 g/ha	-	Apply the higher rates when climatic conditions favour Early Blight or Late Blight infection and in crops with large canopies. Application Kenso Agcare Avior 800 WG Fungicide may be applied by ground or aerial application equipment in potatoes. Aerial application may be used only for Early Blight (Target Spot) control. Apply in a sufficient volume of water to achieve thorough coverage of all foliage. The volume of water required to achieve this will depend on the stage of growth of the potatoes. For ground application a volume of 200 to 300L/ha is suggested at the start of the season, increasing to 500 to 600L/ha in a vigorous crop at full canopy. For aerial application (Early Blight only) , a volume of 18 to 24L/ha is recommended. Spray Interval Consecutive applications should be applied at 7 to 14 day intervals at any time between early shoot growth and 14 days before harvest. Use the recommended shorter application interval in the following circumstances: 1. Under humid weather conditions which are favourable for Early or Late Blight infection. 2. When there is rapid vegetative growth during the early part of the crop cycle. 3. At the first sign of Late Blight infection. Resistance Management Disease control may be reduced if strains of pathogens less sensitive to Kenso Agcare Avior 800 WG Fungicide develop. Apply Kenso Agcare Avior 800 WG Fungicide in a protective spray program. DO NOT wait until disease levels have built up to make applications as this reduces the effectiveness of control and increases the risk of resistance development. As a precaution DO NOT apply more than 1/3 of the total fungicide sprays per crop as Kenso Agcare Avior 800 WG Fungicide. A maximum of 3 consecutive applications of Kenso Agcare Avior 800 WG Fungicide are to be applied. They must be preceded and followed by at least 3 applications of fungicide(s) ® from a different fungicide group(s) such as Bravo, before Kenso Agcare Avior 800 WG Fungicide is used again in that crop. Where Late Blight infection has occurred it is recommended that single sprays of Kenso Agcare Avior 800 WG Fungicide be alternated with 2 sprays of Bravo or a fungicide(s) from another group(s). Where crops are grown successively alternation should continue between crops.

Crop	Disease	Rate/100L	WHP	Critical Comments
Tomatoes Except greenhouse	Early Blight (Target Spot) (<i>Alternaria solani</i>)	125 g/ha or 12.5 g/100L	1 day	Application Apply in sufficient volume of water to achieve thorough coverage of all foliage. The volume of water required to achieve this will depend on the stage of growth of the tomatoes and the method of trellising which influences canopy volume. In the case of dilute spraying (g/100L) apply in the range of 400 to 500L/ha after transplanting and increase to 800 to 1000L/ha at full canopy. In the case of fully trellised tomatoes at full canopy, application volumes should be increased to 900L/ha to achieve these results with high volume spraying. For Late Blight and Sclerotinia control use the higher rates when climatic conditions are humid and mild, which favours disease infection.
	Late Blight (<i>Phytophthora infestans</i>) Sclerotinia (<i>Sclerotinia minor</i>)	155 to 190 g/ha or 16 to 19 g/100L		Spray Interval Consecutive applications should be applied at 7 to 14 day intervals commencing soon after transplanting and continuing up to fruit maturity. Use the recommended shorter application interval in the following circumstances: 1. Under humid weather conditions which are favourable for disease infection. 2. When there is rapid vegetative growth during the early part of the crop cycle. Resistance Management Disease control may be reduced if strains of pathogens less sensitive to Kenso Agcare Avior 800 WG Fungicide develop. Apply Kenso Agcare Avior 800 WG Fungicide in a protective spray program. DO NOT wait until disease levels have built up to make applications as this reduces the effectiveness of control and increases the risk of resistance development. As a precaution DO NOT apply more than 1/3 of the total fungicide sprays per crop as Kenso Agcare Avior 800 WG Fungicide. A maximum of 3 consecutive applications of Kenso Agcare Avior 800 WG Fungicide are to be applied. They must be preceded and followed by at least 3 applications of fungicide(s) from a different fungicide group(s) ® such as Bravo , before Kenso Agcare Avior 800 WG Fungicide is used again in that crop. Where Late Blight infection has occurred it is recommended that single sprays of Kenso Agcare Avior 800 WG Fungicide be alternated with 2 sprays of Bravo a fungicide(s) from another group(s). Where crops are grown successively alternation should continue between crops.
Cucurbits	Powdery Mildew (<i>Sphaerotheca fuliginea</i>), Downy Mildew (<i>Pseudoperon ospora cubensis</i>)	25 to 38 g/100L	1 day	Apply the higher rate when climatic conditions favour Powdery or Downy Mildew infection and in crops with large canopies. Application Apply in a sufficient volume of water to achieve thorough coverage of all foliage. The volume of water required to achieve this will depend on the stage of growth of the cucurbits. For dilute spraying (g/100L), an application volume of 300L/ha is suggested where sprays are banded in the early part of the season, increasing to 1000L/ha as a broadcast spray in a vigorous crop at full canopy. Spray Interval Consecutive applications should be applied at 7 to 14 day intervals, commencing soon after transplanting and continuing up to fruit maturity. Use the recommended shorter application interval in the following circumstances: 1. Under humid weather conditions which are favourable for Powdery Mildew, Downy Mildew or Gummy Stem Blight infection. 2. When there is rapid vegetative growth during the early part of the crop cycle. Resistance Management Disease control may be reduced if strains of pathogens less sensitive to Kenso Agcare Avior 800 WG Fungicide develop. Apply Kenso Agcare Avior 800 WG Fungicide in a protective spray program. DO NOT wait until disease levels have built up to make applications as this reduces the effectiveness of control and increases the risk of resistance development. As a precaution DO NOT apply more than 1/3 of the total fungicide sprays up to a maximum of 4 sprays of Kenso Agcare Avior 800 WG Fungicide per crop. A maximum of 2 consecutive applications of Kenso Agcare Avior 800 WG Fungicide are to be applied. They must be preceded and followed by at least 2 applications of fungicide(s) from a different fungicide ® group(s) such as Bravo to control Downy Mildew and Gummy Stem Blight and ® Nimrod to control Powdery Mildew, before Kenso Agcare Avior 800 WG Fungicide is used again in that crop. Where crops are grown successively alternation should continue between crops.
	Gummy Stem Blight (<i>Didymella bryoniae</i>)	38 g/100L		

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS

Avocados: DO NOT HARVEST FOR 7 DAYS AFTER APPLICATION

Cucurbits, Passionfruit, Tomatoes: DO NOT HARVEST FOR 1 DAY AFTER APPLICATION

Grapes: DO NOT HARVEST FOR 14 DAYS AFTER APPLICATION

Mangoes: DO NOT HARVEST FOR 3 DAYS AFTER APPLICATION

Poppies: DO NOT HARVEST FOR 6 WEEKS AFTER APPLICATION

Potatoes: NOT REQUIRED WHEN USED AS DIRECTED

FUNGICIDE RESISTANCE WARNING

GROUP 11 FUNGICIDE

Kenso Agcare Avior 800 WG Fungicide is a member of the Strobilurin and related products group of fungicides. For fungicide resistance management the product is a Group **11** Fungicide. Some naturally occurring individual fungi resistant to the product and other Group **11** Fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by this product or other Group **11** Fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, Kenso Agcare Australia Pty Ltd accepts no liability for any losses that may result from the failure of this product to control resistant fungi.

RE-ENTRY PERIODS

DO NOT enter treated areas until the spray has dried, unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use. For all activities in grapes and high exposure activities in all other proposed crops the following re-entry statements apply:

- After treating Tomatoes, **DO NOT** allow entry into treated areas for high exposure activities such as harvesting or pruning for 7 days, unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each days use.
- After treating Cucurbits, **DO NOT** allow entry into treated areas for high exposure activities such as harvesting or pruning for 14 days, unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each days use.

- After treating Potatoes, **DO NOT** allow entry into treated areas for high exposure activities such as harvesting for 16 days, unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each days use.
- After treating Avocadoes, Mangoes and Passionfruit, **DO NOT** allow entry into treated areas for high exposure activities such as harvesting or pruning for 19 days, unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each days use.
- After treating Grapes, **DO NOT** allow entry into treated areas for low exposure activities such as scouting for 9 days, unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each days use.
- After treating Grapes, **DO NOT** allow entry into treated areas for high exposure activities such as harvesting or pruning for 27 days, unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each days use.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

WARNING - Kenso Agcare Avior 800 WG Fungicide is extremely phytotoxic to certain apple varieties.

Drift Warning:

Extreme care must be used to prevent injury to apple trees.

- DO NOT** spray Kenso Agcare Avior 800 WG Fungicide where spray drift may reach apple trees.
- DO NOT** spray under meteorological conditions or from spraying equipment, which could be expected to cause spray to drift onto nearby susceptible plants, adjacent crops, crop lands or pastures. Conditions which may contribute to drift include thermal inversions, excessive wind speed, certain sprayer nozzle/pressure combinations, small spray droplet size (150 microns or less) etc.
- DO NOT** use spray equipment, which has been previously used to apply Kenso Agcare Avior 800 WG Fungicide to spray apple trees. Even trace amounts can cause unacceptable phytotoxicity.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

HIGHLY TOXIC TO AQUATIC LIFE

DO NOT contaminate streams, rivers or waterways with the chemical or used container.

DO NOT apply under weather conditions or from spraying equipment which could be expected to cause spray drift on adjacent areas, particularly wetlands, waterbodies or watercourses.

APPLICATION

DO NOT use concentrations exceeding 4X when applying through low volume application equipment, except when applying by air. In these cases adequate coverage of all plant surfaces is still required to achieve control of diseases.

Mixing

Fill tank 50 to 75% full then start agitation. Slowly pour granules into tank and maintain agitation throughout spraying. **DO NOT** pre-mix granules.

Application in Tree Crops and Vines

Dilute spraying: Use a sprayer designed to apply high volumes of water up to the point of run-off and matched to the crop being sprayed. Set up and operate the sprayer to achieve even coverage throughout the crop canopy. Apply sufficient water to cover the crop to the point of run-off. Avoid excessive run-off.

The required water volume may be determined by applying different test volumes, using different settings on the sprayer, from industry guidelines or expert advice. Add the amount of product specified in the Direction for Use table for each 100L of water. Spray to the point of run-off. The required dilute spray volume will change and the sprayer set up and operation may also need to be changed, as the crop grows.

Concentrate spraying: Use a sprayer designed and set up for concentrate spraying (that is a sprayer which applies water volumes less than those required to reach the point of run-off) and matched to the crop being sprayed. Set up and operate the sprayer to achieve even coverage throughout the crop canopy using your chosen water volume. Determine an appropriate dilute spray volume (see Dilute spraying above) for the crop canopy. This is needed to calculate the concentrate mixing rate. The mixing rate for concentrate spraying can then be calculated in the following way:

Example only

- Dilute spray volume as determined above: for example 1000L/ha
- Your chosen concentrate spray volume: for example 500L/ha
- The concentration factor in this example is: 2 x (ie 1000L ÷ 500L = 2)
- If the dilute label rate is 50 g/100L, then the concentrate rate becomes 2 x 50, that is 100 g/100L of concentrate spray.

The chosen spray volume, amount of product per 100L of water, and the sprayer set up and operation may need to be changed as the crop grows. For further information on concentrate spraying, users are advised to consult relevant industry guidelines, undertake appropriate competency training and follow industry Best Practices.

Compatibility/Tank Mixing

Kenso Agcare Avior 800 WG Fungicide may be mixed in the spray vat with any one of the following products: Ambush®, Bravo®, Captan® WG, copper hydroxide,

Ken-Tac 100, Dipel® DF, Karate® EC, Larvin® 375, Fortress® 500, Talstar® 80SC.

A mixture of Kenso Agcare Avior 800 WG Fungicide with more than one of these products or with any other product may be ineffective or may cause serious damage. The use of such a mixture is not recommended and would therefore be entirely at the user's risk. If tank mixes are to be used observe all directions, precautions and limitations on all products to be used.

Note: On some tomato varieties tank mixtures of and Kensban 500 or Nitofol® or Supracide® or Lebaycid® or Kelthane® MF have been found to be phytotoxic.

DO NOT tank mix these products with Kenso Agcare Avior 800 WG Fungicide.

Export of Treated Produce

Grapes

While Maximum Residue Limits (MRL) have been set in many major wine export destinations, some export destinations have not finalised MRL applications. For further information regarding export tolerances please contact your winery, Kenso representative or the Australian Wine Research Institute.

Other Crops

While Maximum Residue Limits (MRL) have been set in many major export destination, it should be noted that MRL's or Import tolerances may not be established in all export destinations. For further information regarding export tolerances please contact your export organisation or Kenso Agcare representative.